
Plant genes determining compatibility with hyphal pathogens and symbionts

**Dissertation an der Fakultät für Biologie
der Ludwig-Maximilians-Universität München**

vorgelegt von

Aline Banhara Pereira

aus Rio de Janeiro

München, den 11.12.2014

Erstgutachter: Prof. Dr. Martin Parniske
Zweitgutachter: Dr. Thomas Ott

Tag der mündlichen Prüfung: 23.02.2015

Eidesstattliche Versicherung

Ich versichere hiermit an Eides statt, dass die vorliegende Dissertation von mir selbstständig und ohne unerlaubte Hilfe angefertigt ist. Einige der in dieser Dissertation geschilderten Experimente wurden in Zusammenarbeit mit anderen Wissenschaftlern durchgeführt. Diese sind in den entsprechenden Abbildungsunterschriften sowie in einer Aufstellung dieser Beiträge auf den Seiten 100 und 101 detailliert dargestellt.

München, den _____

Aline Banhara Pereira

Erklärung

Hiermit erkläre ich, dass die Dissertation nicht ganz oder in wesentlichen Teilen einer anderen Prüfungskommission vorgelegt worden ist. Ich habe nicht versucht, anderweitig eine Dissertation einzureichen oder mich einer Doktorprüfung zu unterziehen.

München, den _____

Aline Banhara Pereira

For Theo

Table of contents

Abbreviations	6
Species names and affiliations.....	10
List of publications.....	12
1. Summary	13
2. Zusammenfassung	15
3. Aims of the thesis	17
4. Introduction	18
4.1 The arbuscular mycorrhiza symbiosis and the common symbiosis pathway 19	
4.2 The <i>Piriformospora indica</i> symbiosis with plant roots.....	21
4.3 Nutrient exchange in AM and <i>P. indica</i> symbioses	23
4.4 AM symbiosis and <i>P. indica</i> protect plant hosts against environmental challenges.....	25
4.5 Different approaches to explore possible genetic commonalities of biotrophic interactions.....	26
4.6 The plant immune system	28
4.7 Role of compatibility factors in biotrophy	29
5. Results.....	31
5.1 <i>A. thaliana</i> HCSG mutants reduce the reproductive success of the oomycete <i>Hpa</i> 31	
5.2. Haustorial development of <i>Hpa</i> is impaired in <i>A. thaliana</i> HCSG mutants	38
5.3. <i>A. thaliana</i> HCSG mutants do not exhibit constitutive or enhanced defence responses	48
5.4. Growth kinetics of <i>Pseudomonas syringae</i> are not altered on the <i>A. thaliana</i> HCSG mutants.....	55
5.5. <i>P. indica</i> colonization of <i>L. japonicus</i> roots occurs independently of common symbiosis genes	57
5.6. <i>P. indica</i> did not induce β -glucuronidase (GUS) activity in the <i>L. japonicus</i> symbiosis reporter line T90	62
5.7. Intracellular colonization of <i>A. thaliana</i> roots by <i>P. indica</i> is independent of homologs of common symbiosis genes	63
5.8. Plant growth promotion by <i>P. indica</i> is influenced by nutrient availability	66
5.9. <i>A. thaliana</i> homologs of common symbiosis genes are not required for <i>P.</i> <i>indica</i> -induced growth promotion.....	70
6. Discussion	74
6.1 <i>A. thaliana</i> mutants for homologs of common symbiosis genes impair haustorial development and the reproductive success of a biotrophic oomycete 74	
6.2 Symbiosis-related genes are not required for the colonization of <i>L.</i> <i>japonicus</i> and <i>A. thaliana</i> with the root endophytic fungus <i>P. indica</i>	75

6.3. Increased fungal biomass in <i>L. japonicus</i> common symbiosis mutants suggests cross-talk between symbiosis signaling and defence or developmental pathways.....	76
6.4. Growth promotion of <i>A. thaliana</i> by <i>P. indica</i> is independent of homologs of common symbiosis genes but is affected by nutrient availability	77
6.5. <i>A. thaliana</i> homologs of common symbiosis genes: compatibility factors for <i>Hpa</i> accommodation	80
6.6. “Common symbiosis pathway” in <i>A. thaliana</i> ?.....	81
6.7. Evolutionary forces driving the retention of the HSCGs in <i>A. thaliana</i>	82
7. Conclusion	86
8. Materials and Methods	87
8.3. Fungal/plant material and growth conditions	87
8.4. <i>A. thaliana</i> stable transformation	90
8.5. Pathogen assays and phenotypic analyses	90
8.6. Observation of fluorescently labelled haustoria	91
8.7. Analysis of oomycete-associated callose deposition.....	91
8.8. Analyses of plant root colonization by <i>P. indica</i>	92
8.9. Elicitor treatment.....	92
8.10. DNA/RNA extraction and qPCR analyses.....	93
8.11. β -glucuronidase (GUS) staining assays	94
8.12. Gene structure and phylogenetic analyses	95
8.13. Statistics and data visualisation	95
8.14. Cloning	96
9. Declaration of contributions from other researches	101
10. References.....	103
11. List of figures	119
12. List of tables	121
13. Acknowledgements	122
14. Curriculum Vitae.....	124
15. Appendix.....	128
Appendix 1: Pruned alignment of kinase domain of LjSYMRK (amino acid positions 593-870) and homologous stretches of related MLD-LRR-RLK sequences	128
Appendix 2: Pruned alignment of extracytoplasmic region of LjSYMRK (amino acid positions 30-517) and homologous stretches of related MLD-LRR-RLKs.	133
Appendix 3: Host-related metabolic cues affect colonization strategies of a root endophyte	144

Abbreviations

°C	degree(s) Celsius
Δ	delta
μ	micro
aa	amino acid
AM	Arbuscular Mycorrhiza
Amt	ammonium transporter
ANOVA	analysis of variance
bp	base pair(s)
BLAST	Basic Local Alignment Search Tool
Ca ²⁺	calcium
CCaMK	Calcium/Calmodulin dependent Kinase
cDNA	complementary DNA
CFP	Cyan Fluorescent Protein
CLSM	confocal laser scanning microscopy
co	complementation line
Col-0	Columbia-0
CSGs	common symbiosis genes
df	degree(s) of freedom
DMR	Downy Mildew Resistant
DNA	deoxyribonucleic acid
dpi	day(s) post infection/inoculation
FLS2	Flagellin-Sensing 2
fwd	forward
g	gram(s) / gravitational acceleration
g/L	gram(s) per liter
GDPC	glycin-asparagin-prolin-cystein
GFP	Green Fluorescent Protein
GG	golden gate
GPAT	Glycerol-Phosphate Acyl-Transferase
GUS	β-Glucuronidase
h	hour(s)
HCSG	homologs of common symbiosis genes
HO	Hoagland's

<i>Hpa</i>	<i>Hyaloperonospora arabidopsidis</i>
HR	hypersensitive response
HSD	honestly significant difference
IOS	Impaired Oomycete Susceptibility
ISR	induced systemic resistance
JA	jasmonic acid
kg	kilogram
K	potassium
KDRI	Kazusa DNA Research Institute
L	litre(s)
LRR	leucine-rich repeat
m	meter(s) / mili
M	Molar
μ	micro
min	minute(s)
mL	milliliter(s)
MLD	malectin-like domain
mm	millimeter(s)
mRNA	messenger RNA
MLO	Mildew Resistance <i>Locus</i> O
mol	mol(es)
MS	Murashige and Skoog
MST	Monosaccharide Transporter
MYB	myeloblastosis
N	nirogen
n	nano
NASC	The Nottingham Arabidopsis Stock Centre
NH ₄ ⁺	ammonium
NIN	Nodule Inception
NO ₃ ⁻	nitrate
NUP	nucleoporin
OXI	Oxidative Signal Inducible
p	p-value
<i>P</i>	probability
P	phosphate

PAD4	Phytoalexin Deficient 4
PAMP	pathogen-associated molecular pattern
PCR	polymerase chain reaction
PDK	Phosphoinositide-Dependent protein Kinase
Pi	inorganic orthophosphate
PI	propidium iodide
PI-LTP	Phosphatidylinositol-Lipid Transfer Protein
PLD	Phospholipase D
PMR	Powdery Mildew Resistant
PPA	prepenetration apparatus
PR1	Pathogenesis-Related Gene 1
PSKR1	Phytosulfokine Receptor 1
PT	phosphate transporter
qRT-PCR	quantitative real-time PCR
RAM2	Required for Arbuscular Mycorrhiza 2
RLK	receptor-like kinase
rev	reverse
RIC	ROP (RHO-like GTPase of Plants)- Interactive CRIB (CDC42/RAC interactive binding)
rRNA	ribosomal RNA
ROI	region of interest
ROS	reactive oxygen species
rpm	rotations per minute
RT	room temperature
RT-PCR	polymerase chain reaction following reverse transcription
RNA	ribonucleic acid
s	second(s)
SA	salicylic acid
S.D.	standard deviation
SEH	<i>SEC13</i> Homolog
ShRK	SYMRK homologous Receptor-like Kinase
SP	signal peptide
SYM	symbiosis
SYMRK	Symbiosis Receptor-like Kinase

TAIR	The Arabidopsis Information Resource
TEF	Transcriptional Elongation Factor
TM	transmembrane domain
TX	texas red
UBI	Ubiquitin
WGA-AF	wheat germ agglutinin
WT	wild-type
X-Gluc	5-bromo-4-chloro-3-indolyl glucuronide
w/v	weight/volume
YFP	Yellow Fluorescent Protein

Species names and affiliations

Plants (Kingdom Plantae)

Full scientific name	Abbreviation	Order	Common name
<i>Adenostyles alliariae</i> (Gouan) A. Kern.	<i>A. alliariae</i>	Asterales	hedge-leaved Adenostyle
<i>A. thaliana thaliana</i> (L.). Heynh	<i>A. thaliana</i> , <i>A. thaliana</i>	Brassicales	thale cress
<i>Brassica rapa</i> L.	<i>B. rapa</i>	Brassicales	chinese cabbage
<i>Hordeum vulgare</i> L.	<i>H. vulgare</i>	Poales	barley
<i>Lotus japonicus</i> (Regel) K. Larsen	<i>L. japonicus</i>	Fabales	birdsfoot trefoil
<i>Lycopersicon esculentum</i> Mill.	<i>L. esculentum</i>	Sonales	tomato
<i>Medicago truncatula</i> Gaertn.	<i>M. truncatula</i>	Fabales	barrel medic
<i>Oryza sativa</i> L.	<i>O. sativa</i>	Poales	rice
<i>Plantago lanceolata</i> L.	<i>P. lanceolata</i>	Lamiales	narrowleaf plantain
<i>Triticum aestivum</i> L.	<i>T. aestivum</i>	Poales	wheat
<i>Zea mays</i> L.	<i>Z. mays</i>	Poales	maize
<i>Nicotiana tabacum</i> L.	<i>N. tabacum</i>	Solanales	tobacco

Animals (Kingdom Animalia)

Full scientific name	Abbreviation	Order	Common name
<i>Melitaea cinxia</i> L.	<i>M. cinxia</i>	Lepdoptera	Glanville fritillary
<i>Phaedon cochleariae</i> Fabricius.	<i>P. cochleariae</i>	Coleoptera	mustard beetle
<i>Oreina elongata</i> Suffrian.	<i>O. elongata</i>	Coleoptera	alpine leaf beetle
<i>Oreina cacaliae</i> Schrank.	<i>O. cacaliae</i>	Coleoptera	alpine leaf beetle
<i>Spodoptera littoralis</i> Boisduval.	<i>S. littoralis</i>	Lepdoptera	african cotton leafworm

Fungi (Kingdom Fungi)

Full scientific name	Abbreviation	Order	Common name
<i>Alternaria brassicae</i> (Berk.). Sacc	<i>A. brassicae</i>	Pleosporales	Alternaria black spot
<i>Blumeria graminis</i> (DC.) Speer	<i>B. graminis</i>	Erysiphales	barley powdery mildew

<i>Erysiphe orontii</i>	<i>E. orontii</i>	Erysiphales	Brassica powdery mildew
<i>Erysiphe cruciferarum</i>	<i>E. cruciferarum</i>	Erysiphales	Brassica powdery
Opiz ex L. Junell mildew			
<i>Fusarium graminearum</i>	<i>F. graminearum</i>	Hypocreales	Fusarium head blight
Schwabe			
<i>Glomus intraradices</i>	<i>G. intraradices</i>	Glomerales	AM fungus
Schenck & Smith			
<i>Golovinomyces orontii</i>	<i>G. orontii</i>	Erysiphales	Brassica powdery mildew
(Castagne) V.P. Heluta			
<i>Piriformospora indica</i>	<i>P. indica</i>	Sebacinales	
Sav. Verma, Aj. Varma, Rexer, G. Kost & P. Franken			
<i>Piriformospora williamsii</i>	<i>P. williamsii</i>	Sebacinales	
Zuccaro & M. Weiss			
<i>Podosphaera plantaginis</i>	<i>P. plantaginis</i>	Erysiphales	powdery mildew
(Castagne) U. Braun & S. Takam.			
<i>Rhizophagus irregularis</i>	<i>R. irregularis</i>	Glomerales	AM fungus
Blaszk, Wubet, Renker & Buscot			
<i>Uromyces cacaliae</i> (DC) Unger.	<i>U. cacaliae</i>	Erysiphales	.rust fungus

Oomycetes (Kingdom Chromalveolata)

Full scientific name	Abbreviation	Order	Common name
<i>Hyaloperonospora arabidopsidis</i>	<i>H. arabidopsidis</i> , <i>Hpa</i>	Peronosporales	downy mildew
(Gäum.) Göker, Riethm., Voglmayr, Weiß & Oberw.			
<i>Phytophthora infestans</i>	<i>P. infestans</i>	Peronosporales	late blight
(Mont.) de Bary			
<i>Phytophthora palmivora</i> Butler.	<i>P. palmivora</i>	Peronosporales	bud-rot of palms

Bacteria (Kingdom Eubacteria)

Full scientific name	Abbreviation	Order
<i>Mesorhizobium loti</i>	<i>M. loti</i>	<i>Rhizobiales</i>
<i>Pseudomonas syringae</i>	<i>P. syringae</i>	Pseudomonales

List of publications

Research papers

Banhara A*, Ried MK*, Binder A, Gust AA, Höfler C, Hückelhoven R, Nürnberger T, and Parniske M (2015) Symbiosis-related genes sustain the development of a downy mildew pathogen on *A. thaliana*. Under review.

* These authors contributed equally to the work.

Banhara A, D Yi, Kühner R, Zuccaro A and Parniske M (2015) Biotrophic colonization of host roots by *Piriformospora indica* occurs independently of plant common symbiosis genes. Under review.

Lahrmann U, Ding Y, **Banhara A**, Rath M, Hajirezaei MR, Döhlemann S, von Wirén N, Parniske M, Zuccaro A (2013) Host-related metabolic cues affect colonization strategies of a root endophyte. *Proc Natl Acad Sci USA*. 110(34): 13965-70.

Alves-Ferreira M, Wellmer F, **Banhara A**, Kumar V, Riechmann JL, Meyerowitz EM (2007) Global expression profiling applied to the analysis of *A. thaliana* stamen development. *Plant Physiol*. 145: 747–762.

1. Summary

The majority of land plants are able to engage in root symbiosis with arbuscular mycorrhizal (AM) fungi. As member of the Brassicales, the model plant *Arabidopsis thaliana* is unable to form AM symbiosis, a fact that is correlated with a partial loss of the archaic genetic program for the intracellular accommodation of AM fungi that constitutes the common symbiosis pathway. Despite this partial loss, *A. thaliana* retained homologs of common symbiosis genes (HCSGs), raising questions on evolutionary forces that drive the maintenance of this core program on an AM-asympbiotic plant. Biotrophic microorganisms like AM fungi associates with living plant cells and often colonize them by forming specialized intracellular feeding structures that are surrounded by a plant-derived membrane, named arbuscules in symbiosis, or haustoria in pathogenic interactions. Based on structural and functional similarities between those accommodation structures, we speculated that the *A. thaliana* HCSGs might be utilized during the interaction with other biotrophic microorganisms other than AM fungi. In our work, we used two main approaches to test this hypothesis, that is, the interactions of *A. thaliana* with the biotrophic root symbiont *Piriformospora indica*, and with two biotrophic foliar pathogens: the oomycete *Hyaloperonospora arabidopsidis* (*Hpa*), and the powdery mildew fungus *Erysiphe cruciferarum*. We show that root colonization by *P. indica* is independent of the common symbiosis genes (CSGs) in the legume *Lotus japonicus* or its homologs in *A. thaliana*. Interestingly, though, the CSGs seem to have a negative role on fungal proliferation in *L. japonicus* roots, suggesting an interconnection between symbiosis signalling and plant defence or developmental programs. Colonization of *A. thaliana* leaves by *E. cruciferarum* is also independent of the HCSGs. We discovered, however, that these genes are required for haustoria development and full reproductive success of *Hpa* on *A. thaliana*, and the decreased susceptibility to the oomycete observed in the HCSG mutants occurs without constitutive or exacerbated activation of known defence responses. Together our findings reveal novel genetic commonalities in the plant program for the intracellular accommodation of biotrophic organisms in symbiosis and disease. The retention of HCSGs in the *A. thaliana* lineage calls for a selective advantage conferred by these genes, despite them supporting the colonization of an oomycete pathogen. We were therefore prompted to

investigate whether ecological conditions exist, under which *Hpa* improves host fitness.

2. Zusammenfassung

Ein Großteil der terrestrischen Pflanzen ist in der Lage Wurzelsymbiosen mit arbuskulären Mykorrhizapilzen (AM Pilze) einzugehen. Als Mitglied der Brassicales kann die Modellpflanze *A. thaliana* keine AM Symbiose ausbilden. Dies steht in Zusammenhang mit dem teilweisen Verlust eines archaischen genetischen Programms, des “common symbiosis” Signalwegs, welcher für die intrazelluläre Einnistung von AM Pilzen nötig ist. Trotz dieses teilweisen Verlusts, finden sich noch Homologe von “common symbiosis” Genen (HCSGs) in *A. thaliana*, was Fragen nach den evolutionären Triebkräften aufwirft, die für den Erhalt dieses Programmes in einer AM asymbiotischen Pflanze verantwortlich sind. Biotrophe Organismen wie AM Pilze kolonisieren Pflanzenzellen üblicherweise durch die Ausbildung von speziellen Ernährungsstrukturen, die von einer pflanzlichen Membran umschlossen sind. In der AM Symbiose werden diese Strukturen als Arbuskel bezeichnet; bei pathogenen Interaktionen spricht man von Haustorien. Aufgrund von strukturellen und funktionellen Gemeinsamkeiten zwischen diesen Strukturen, vermuteten wir, dass die *A. thaliana* HCSGs in biotrophen Interaktionen mit anderen Mikroorganismen als AM Pilzen eine Rolle spielen. In der vorliegenden Arbeit nutzten wir zwei Hauptansätze um diese Hypothese zu testen, nämlich die Untersuchung der Interaktion von *A. thaliana* mit dem biotrophen Wurzelsymbionten *Piriformospora indica* und mit zwei biotrophen Blattpathogenen: dem Oomyceten *Hyaloperonospora arabidopsidis* (*Hpa*) und dem Mehltaupilz *Erysiphe cruciferarum*. Wir zeigen, dass die Besiedlung durch *P. indica* in der Leguminose *Lotus japonicus* und in *A. thaliana* unabhängig von den “common symbiosis” Genen (CSGs) bzw. den HCSGs ist. Interessanterweise scheinen die CSGs jedoch eine negative Rolle bei der Vermehrung des Pilzes in *L. Japonicus* Wurzeln auszuüben, was auf eine Verbindung zwischen dem symbiotischen Signalweg und den pflanzlichem Abwehr- oder Entwicklungsprogrammen schließen lässt. Die Kolonisierung von *A. thaliana* Blättern durch *E. cruciferarum* ist ebenfalls unabhängig von den HCSGs. Wir stellen jedoch fest, dass diese Gene für die Haustorienentwicklung und für den maximaler Fortpflanzungserfolg von *Hpa* in *A. thaliana* nötig sind. Die beobachtete verminderte Suszeptibilität gegenüber dem Oomyceten in HCSG Mutanten geht dabei nicht mit einer konstitutiven oder verstärkten Aktivierung von Abwehrreaktionen einher.

Zusammengenommen offenbaren diese Ergebnisse neuartige genetische Gemeinsamkeiten im pflanzlichen Programm bei der Einnistung von symbiotischen und krankheitsauslösenden biotrophen Organismen. Die Tatsache, dass HCSGs im Genom von *A. thaliana* behalten wurden/konserviert sind, legt nahe, dass diese Gene nicht nur die Kolonisierung durch einen pathogenen Oomyceten unterstützen, sondern außerdem selektive Vorteile für die Pflanze bieten. Aus diesem Grund untersuchen wir, ob es ökologische Bedingungen gibt, unter denen *Hpa* die Fitness der Wirtspflanze verbessert.

3. Aims of the thesis

The main objective of this thesis was to elucidate the role of symbiosis-related genes of *Arabidopsis thaliana* and *Lotus japonicus* in biotrophic interactions with hyphal pathogens and symbionts. With that purpose we addressed the following:

1. Investigate the influence of plant genotype on root cell colonization and/or on the growth-promoting effect of the fungal endophyte *Piriformospora indica* on *A. thaliana*;
 - 1.1. Explore the existence of overlaps between the root-*P. indica* association and classical root symbioses with rhizobia and arbuscular mycorrhizal fungi;
 - 1.2. Investigate the role of HCSGs in the intracellular colonization of *A. thaliana* roots;
 - 1.3. Examine whether the CSGs are required for the colonization of *L. japonicus* roots by *P. indica*;
 - 1.4. Confirm growth promotion of *A. thaliana* (Col-0) by *P. indica* and to what extent this effect is dependent on the HCSGs.
2. Evaluate the requirement of the HCSGs for the intracellular accommodation and reproduction of the oomycetal pathogen *Hyaloperonospora arabidopsidis* (*Hpa*) in *A. thaliana* leaf epidermal cells;
3. Analyse if the HCSGs are important for colonization of *A. thaliana* leaves by the fungal pathogen *Erysiphe cruciferarum*;
4. Test whether the HCSGs are necessary for the infection of *A. thaliana* with the extracellular bacterial pathogen *Pseudomonas syringe*.
5. Exclude the role of the *A. thaliana* HCSGs as potential negative regulators of plant immunity by looking for altered defence responses on HCSG mutants upon infection with the pathogenic oomycete *Hpa*.

4. Introduction

Plants engage in biotrophic relationships with fungi from diverse taxonomic groups as well as with oomycetes (Voegelé & Mendgen, 2003; Schulze-Lefert, 2004), fungal-like eukaryotic microorganisms (Margulis & Schwartz, 2002). Obligate biotrophs depend on living cells to obtain their nutrients, grow and complete their life cycle (Spanu *et al.*, 2010). The interaction of plants with symbiotic biotrophs such as arbuscular mycorrhiza (AM) fungi is an important acquisition of higher plants that is thought to have assisted them in the colonization of land (Schüßler & Walker, 2011). On the other hand, pathogenic biotrophic organisms like oomycetes, rust fungi, and powdery mildew fungi are agents of devastating diseases of crop plants (Voegelé & Mendgen, 2003; Schulze-Lefert, 2004). Even though symbiotic and pathogenic relationships have opposite outcomes for the plants, their establishments share features such as overall suppression of or evasion from host immune responses (Okmen & Doehlemann, 2014; van Schie & Takken, 2014). Moreover, both pathogenic and symbiotic biotrophs form specialized intracellular infection structures (Yi & Valent, 2013) known as arbuscules in the AM symbiosis and haustoria in pathogenic interactions. Both structures are surrounded by plant-derived membranes called, respectively, periarbuscular membrane and perih Haustorial membrane, which isolate the microorganisms from the plant cytoplasm. These membranes delimitate the respective periarbuscular space or the extrahaustorial matrix (Mims *et al.*, 2004; Pumplun & Harrison, 2009), which likely constitute the site of nutrient and signal exchange between the microorganisms and the plant host (Mendgen & Hahn, 2002; Voegelé & Mendgen, 2003; Parniske, 2008; Bonfante & Genre, 2010; Harrison, 2012). The structural and functional similarities (Harrison, 1999; Parniske, 2000) between accommodation organs for microbes in symbiosis and disease (Figure 1) raised the hypothesis that both types of interactions rely on a shared genetic program (Parniske, 2000).




Figure 1 – Schematic representation of intracellular accommodation organs in symbiosis and disease. All accommodation organs are surrounded by a plant-derived membrane that separates them from the plant-cell cytoplasm and likely creates an environment for exchange of substances between plant and microorganisms. The highly specialized infection cell known as appressorium (arrow) (Deising *et al.*, 2000), and the callose neck (purple) that typically forms upon cell penetration during pathogenic interactions are shown. (A) The arbuscule of *Glomus sp.* inside a *Lotus japonicus* root cortical cell; (B) Hyphae of *Piriformospora indica* within a root epidermal cell from *Arabidopsis thaliana*; (C) The haustoria of an oomycete (*Hyaloperonospora parasitica*) inside an *A. thaliana* leaf mesophyll cell; (D) The haustoria of the powdery mildew fungus *Erysiphe cruciferarum* within a leaf epidermal cell of *A. thaliana*.

4.1 The arbuscular mycorrhiza symbiosis and the common symbiosis pathway

AM symbiosis is one of the most widespread types of symbiosis between plants and microorganisms. It has existed for more than 400 million years and is present in circa 80% of all terrestrial plant species (Parniske, 2008; Smith & Read, 2008). In AM

symbiosis, plants provide carbon sources to fungi of the phylum Glomeromycota, which in turn deliver water, phosphate, nitrogen, and other mineral nutrients via finely branched, tree-shaped hyphae, called arbuscules (Gutjahr & Parniske, 2013; Gutjahr, 2014). The intracellular accommodation of fungal symbionts requires extensive reprogramming of the plant host cell. This process starts with the first contact between fungal hyphae and the plant roots, followed by the formation of the hyphopodium, a structure that attaches to the root surface at the fungal entry point preceding cell invasion (Gutjahr & Parniske, 2013). Even before cell penetration, hyphopodium formation activates cellular processes in which the plant cell nucleus shifts towards the point of contact with the AM fungus before moving through the plant cell vacuole. This initiates the formation of the prepenetration apparatus, a transcellular tubular cytoplasmic column that connects the nucleus of the plant cell with the fungal entry point. This course of events, which occurs in every cell preceding fungal penetration, guides the way of the fungus through the rhizodermal and the outer cortical cells (Genre *et al.*, 2005; Genre *et al.*, 2008). When the inner cortex is reached, the hypha enters the cell, where it finally branches to form the arbuscule (Gutjahr & Parniske, 2013).

The establishment of the AM symbiosis requires an ancient genetic program that is conserved among angiosperms and comprises the “common symbiosis genes” (CSGs) (Kistner & Parniske, 2002; Markmann *et al.*, 2008). In four orders of eurosids, the CSGs are also required for root nodule symbiosis with nitrogen-fixing bacteria (Kistner *et al.*, 2005; Gherbi *et al.*, 2008). In the order Fabales (legumes), the interaction occurs with bacteria generally known as rhizobia, while in the Rosales, Cucurbitales, and Fagales the symbiosis is termed Actinorhiza and characterizes the interaction with *Frankia* bacteria (Pawlowski & Bisseling, 1996). The proteins encoded by the CSGs participate in a signal transduction pathway that leads from the perception of microbe signals at the plasma membrane to the activation of symbiosis-related genes in the nucleus (Gutjahr & Parniske, 2013).

AM symbiosis formation requires calcium-spiking, periodic calcium oscillations in the nucleus and in the peri-nuclear region that constitute one of the earliest measurable plant responses during symbiotic stimulation (Ehrhardt *et al.*, 1996; Miwa *et al.*, 2006; Oldroyd, 2013). In the legume *Lotus japonicus*, the products of the CSGs Symbiosis Receptor-like Kinase (SYMRK; (Stracke *et al.*, 2002; Markmann *et al.*,

2008), the nucleoporins of the NUP107-160/NUP84 sub-complex (Alber *et al.*, 2007) NUP85, NUP133 and the SEC13 HOMOLOG1 (SEH1) NENA (Kanamori *et al.*, 2006; Saito *et al.*, 2007; Groth *et al.*, 2010), as well as the nuclear-envelope localised cation channels CASTOR and POLLUX (Adam *et al.*, 1999; Charpentier *et al.*, 2008; Venkateshwaran *et al.*, 2012), are all required for calcium-spiking generation, while the nuclear complex Calcium and Calmodulin dependent protein Kinase (CCaMK)-CYCLOPS is responsible for decoding these signals (Singh & Parniske, 2012; Singh *et al.*, 2014).

It has been speculated that the symbiotic phenotypes of *L. japonicus* mutants defective in the CSGs coding for the nucleoporins *seh1*, *nup133*, or *nup85* could be related to a decrease of POLLUX levels in the inner nuclear membrane due to impaired import caused by structural defects in the NUP107-160 subcomplex (Capoen *et al.*, 2011; Binder & Parniske, 2013). Many nucleoporins of this complex show a high rate of evolution (Baptiste *et al.*, 2005), potentially allowing for distinct functional adaptations while keeping the overall complex structure intact. This rapid evolution may be facilitated by the structural modularity of alpha-solenoid and beta-propeller domains shared by many of these nucleoporins (Hoelz *et al.*, 2011). Agreeing with this, phenotypes of individual NUP107-160 subcomplex mutants vary both in occurrence and severity depending on the organism (Gonzalez-Aguilera & Askjaer, 2012; Binder & Parniske, 2013).

The model plant *A. thaliana* lost the ability to undergo root symbiosis with AM fungi. This loss, which happened after the divergence of the Brassicales, is correlated with the absence of a set of CSGs genes, including *CYCLOPS* and *CCaMK* (Delaux *et al.*, 2014).

4.2 The *Piriformospora indica* symbiosis with plant roots

Besides AM symbiosis, plants are able to undergo root symbiosis with sebacinoid fungi, which belong to the phylum Basidiomycota and are widespread root associated filamentous fungi able to engage into ecto- and endomycorrhizal interactions (Selosse *et al.*, 2009). Members of the order Sebaciniales frequently form endophytic interactions with various taxonomically unrelated hosts, displaying a generalist lifestyle (Weiss *et al.*, 2004). The experimental model for this group of fungi is the

root endophyte *Piriformospora indica*, whose association with plant roots results in a series of benefits to the host. Apart from growth promotion (Peškan-Berghöfer *et al.*, 2004; Shahollari *et al.*, 2007; Sherameti *et al.*, 2008; Camehl *et al.*, 2010; Camehl *et al.*, 2011; Hilbert *et al.*, 2012; Nongbri *et al.*, 2012; Lahrmann *et al.*, 2013; Venus & Oelmüller, 2013), plants grown in the presence of the fungus are more tolerant to abiotic (Waller *et al.*, 2005; Sherameti *et al.*, 2008; Kumar *et al.*, 2009; Ghabooli *et al.*, 2013; Jogawat *et al.*, 2013) and biotic stresses (Waller *et al.*, 2005; Deshmukh & Kogel, 2007; Stein *et al.*, 2008; Molitor *et al.*, 2011), and show increased uptake of nutrients such as phosphorus and nitrogen (Sherameti *et al.*, 2005; Kumar *et al.*, 2011).

Like AM fungi, *P. indica* is able to undergo at least part of its life cycle as a biotroph. However, in plants such as barley and *A. thaliana*, *P. indica* has a distinct, biphasic lifestyle (Jacobs *et al.*, 2011; Zuccaro *et al.*, 2011; Lahrmann & Zuccaro, 2012; Lahrmann *et al.*, 2013). After the initial biotrophic stage, during which the fungus grows within metabolically active root cells, a phase characterized by cell death and necrotrophy or saprotrophy follows. At this point, *P. indica* grows and feeds from dead root cells even though the plant host displays no obvious signs of disease (Deshmukh *et al.*, 2006; Zuccaro *et al.*, 2011; Lahrmann & Zuccaro, 2012; Qiang *et al.*, 2012). Despite the fact that *P. indica* possesses characteristics of both biotrophic and saprotrophic fungi, its lifestyle contrasts with the lifestyle of hemibiotrophic pathogens, whose saprotrophy growth is normally associated with the appearance of disease symptoms in the host (Zuccaro *et al.*, 2011).

To date, little is known about plant genes that directly contribute to the intracellular accommodation of *P. indica* in plant roots during the biotrophic phase, with exception of the tubby-like proteins, which are necessary for normal colonization of *A. thaliana* roots by *P. indica* during this colonization stage (Reitz *et al.*, 2012; Reitz *et al.*, 2013). Tubby-like proteins participate in vesicle trafficking in mammals (Mukhopadhyay & Jackson, 2011) and, in plants, they are targeted to the plasma membrane and released upon abiotic stresses to activate intracellular signalling (Santagata *et al.*, 2001). In regards to host growth promoted by *P. indica*, a few molecules have been shown to participate in this process, such as components of the OXI1 kinase pathway (Camehl *et al.*, 2011), which is involved in plant responses to pathogens such as the oomycete *Hyaloperonospora parasitica* (Rentel *et al.*, 2004), or the *CYP79B2*, *CYP79B3*,

CYP71A13, *PAD3*, and *WRKY33* genes, all required for the synthesis of indole-3-acetaldoxime (IAOx)-derived compounds in *A. thaliana* roots colonized by *P. indica* (Nongbri *et al.*, 2012). In *A. thaliana*, IAOx is an intermediate in the biosynthesis of the hormone IAA and the phytoalexin camalexin (Mikkelsen *et al.*, 2009; Burow *et al.*, 2010), which is usually produced in response to plant pathogens (Glawischnig, 2007; Rauhut & Glawischnig, 2009). In the *cyp79b3*, *cyp79b3*, *cyp71a13*, *pad3*, and *wrky33* mutants, the fungus grows in an uncontrolled manner and the host growth promotion effect is no longer present.

4.3 Nutrient exchange in AM and *P. indica* symbioses

Even though phosphorus is indispensable for plant growth and development, it is mostly available in the soil as inorganic orthophosphate (Pi), which is readily sequestered by cations under acid conditions (Tinker & Nye, 2000), resulting in low availability of free Pi to the plants (Bielecki, 1973). In this context, AM fungi play a crucial role in plant nutrition, as they are able to take up Pi from the soil and release it to the plant (Jakobsen, 1995; Harrison, 1997; Smith & Smith, 2011). Phosphorus is quantitatively the most important mineral nutrient transported from the fungus to the plant in AM symbiosis (Smith & Read, 2008). Its uptake by the host plant requires first the assimilation of Pi by the fungus via high-affinity phosphate transporters expressed exclusively in the extraradical mycelium (Harrison & van Buuren, 1995; Maldonado-Mendoza *et al.*, 2001; Fiorilli *et al.*, 2013), where it is stored in vacuoles as polyphosphate (poly-Pi) before being transported to the intraradical hyphae (Callow *et al.*, 1978; Solaiman *et al.*, 1999; Ezawa *et al.*, 2005). Pi is then hydrolysed and translocated to the periarbuscular space, where it is taken up by cortical root cells (Javot *et al.*, 2007b). This last step involves plant transporters located at the periphery of arbuscules (Bapaume & Reinhardt, 2012). The most studied phosphate transporter specifically induced during AM symbiosis and expressed at the periarbuscular membrane is the *Medicago truncatula* MtPT4 (Harrison *et al.*, 2002). Nevertheless, phosphate transporters that are strongly upregulated upon mycorrhization and expressed in arbuscule-containing cells have been identified in other plant species as well, including potato, tomato, *L. japonicus*, and rice (Rosewarne *et al.*, 1997; Rausch *et al.*, 2001; Paszkowski *et al.*, 2002; Nagy *et al.*, 2005; Maeda *et al.*, 2006; Glassop *et al.*, 2007; Gutjahr *et al.*, 2008). The establishment of AM symbiosis depends on

nutrient availability, and high Pi concentrations in the soil can inhibit the symbiotic interaction (Breuillin *et al.*, 2010; Balzergue *et al.*, 2013; Bonneau *et al.*, 2013; Gosling *et al.*, 2013). Phosphate availability also determines the amount of carbon transferred from the plant to the fungus, which is the “carbon fee” the plant host needs to pay in exchange for the mineral nutrients obtained from the fungus and corresponds to up to 20% of the total carbon photosynthetically fixed (Bago *et al.*, 2003). In the situation of high Pi availability, the symbiosis turns out to be no longer necessary, and the carbon transfer towards the fungus, which occurs in the form of monosaccharides, is reduced (Olsson *et al.*, 2010). The sugar transporter Monosaccharide Transporter2 (MST2) from *Rhizophagus irregularis* (previously known as *Glomus intraradices* (Krüger *et al.*, 2012)) and *Glomus* sp., the largest genus of AM fungi (Schwarzott *et al.*, 2001; Schüßler & Walker, 2011), has been recently described. It participates in the transport of glucose and other hexoses to the fungus and, interestingly, has a temporal expression pattern that coincides with that of MtPT4 (Helber *et al.*, 2011).

While phosphorus nutrition during AM symbiosis has been well studied over the last decades, fewer is known about nitrogen transport to the plant, even though AM fungi improve nitrogen acquisition by the plant as well, mainly as ammonium (Villegas *et al.*, 1996; Smith & Read, 1997; Hawkins *et al.*, 2000; Toussaint *et al.*, 2004; Smith & Smith, 2011). After nitrogen is assimilated over the glutamine synthetase/glutamate synthase pathway (Jin *et al.*, 2005; Cruz *et al.*, 2007), it is transported as arginine along the fungal hyphae (Govindarajulu *et al.*, 2005; Tian *et al.*, 2010) and cleaved by arginase in the arbuscules, where it is released to the symbiotic interface and delivered to the host as ammonium (NH₄⁺) (Bago *et al.*, 2003; Govindarajulu *et al.*, 2005). In support to this model, the *R. irregularis* NH₄⁺ transporter LjAMT2;2 was identified as preferentially expressed in arbuscule-containing cells (Bonfante & Genre, 2010). In addition, two *R. irregularis* NH₄⁺ transporters were shown to act in synchrony to promote NH₄⁺ assimilation and amino acid incorporation by the plant (Lopez-Pedrosa *et al.*, 2006; Perez-Tienda *et al.*, 2011). In *Glomus mosseae*, the NH₄⁺ transporter gene *GmAMT 4.1* was specifically upregulated in arbusculated cells during NH₄⁺ transfer across the periarbuscular membrane to the host cells (Kobae *et al.*, 2010). Corroborating the importance of nitrogen levels for the maintenance of a functional symbiosis, Javot and colleagues showed that, in a *M. truncatula* mutant defective in PT4, nitrogen limitation suppressed premature arbuscule degeneration and root

colonization by AM fungi occurred normally (Javot *et al.*, 2011). Furthermore, similar to high Pi levels in the soil surrounding the roots, elevated nitrogen concentrations can reduce carbon allocation to the fungus (Olsson *et al.*, 2005). Limitation of both phosphorus and nitrogen has a cumulative effect on root colonization, highlighting the importance of these nutrients for AM symbiosis formation (Bonneau *et al.*, 2013).

As in AM symbiosis, the presence of *P. indica* in the plant root also seems to improve nitrogen and phosphorus uptake by the plant host. Recently, it has been shown that phosphate transfer is a major player in the symbiosis between *P. indica* and maize plants. A high affinity phosphate transporter from the fungus (PiPT) was upregulated in *P. indica*-inoculated plants, especially under low phosphate availability, and phosphate was transported from the fungus to the host plants, which had increased biomass compared to non-inoculated controls (Yadav *et al.*, 2010; Pedersen *et al.*, 2013). These results are in accordance to previous findings in *A. thaliana*, where *P. indica* increased host phosphate uptake by up to three-fold (Kumar *et al.*, 2011).

The role of nitrogen uptake in the *A. thaliana*-*P. indica* association has not been conclusively established, even though it was suggested that, different than AM symbiosis, in the *P. indica* symbiosis nitrogen is taken up by the plants mainly in the form of nitrate (Sherameti *et al.*, 2008). It has been revealed that inoculated *A. thaliana* and tobacco seedlings show a transfer of nitrogen from the medium (with nitrate as nitrogen source) to the aerial part, with a concomitant induction of plant enzymes related to nitrate and starch metabolism. These findings are contrary to the observation that the preferred sources of nitrogen in *P. indica* are ammonium and glutamine instead of nitrate (Zuccaro *et al.*, 2011). Supporting the latter finding, we recently identified a NH_4^+ transporter from the fungus, PiAmt1, which is responsible for sensing nitrogen levels in barley and leading to the switch from the biotrophic to the saprotrophic phase upon nitrogen starvation (Lahrmann *et al.*, 2013).

4.4 AM fungi and *P. indica* protect plant hosts against environmental challenges

There is evidence that colonization of plant roots by *P. indica* leads to improved plant tolerance against abiotic stresses such as drought (Sherameti *et al.*, 2008; Ghabooli *et al.*, 2013), high concentrations of salt (Waller *et al.*, 2005; Jogawat *et al.*, 2013), and

reactive oxygen species (Kumar *et al.*, 2009), and heavy metals (Oelmüller *et al.*, 2009). It also protects plants against pathogens like head blight (*Fusarium graminearum*) (Deshmukh & Kogel, 2007) and the powdery mildew fungi *Blumeria graminis* f. sp. *hordei* (Waller *et al.*, 2005; Molitor *et al.*, 2011) in barley, and *Golovinomyces orontii* in *A. thaliana* (Stein *et al.*, 2008). The improved resistance of plants colonized by *P. indica* against pathogens occurs regardless of the fact that the fungus induces a widespread suppression of pathogen-associated molecular pattern (PAMP)-induced host innate immunity (Jacobs *et al.*, 2011). The resistance of *P. indica*-colonized plants against powdery mildew fungi is thought to occur by a different defence mechanism, in which the pathogen reportedly leads to induced systemic resistance (ISR) both in barley (Waller *et al.*, 2005; Stein *et al.*, 2008; Waller *et al.*, 2008) and in *A. thaliana*. Improved tolerance to abiotic stresses (Mena-Violante *et al.*, 2006; Miransari, 2010; Aloui *et al.*, 2011) such as drought stress (Aroca *et al.*, 2007) and to biotic stresses (Smith & Read, 2008) have been described for plants undergoing associations with Glomeromycota fungi as well. Besides, similar to *P. indica*, AM fungi temporarily induce plant immunity at initial stages of root colonization, while the later stages of the symbiosis are characterized by localized suppression of defence responses (Kapulnik *et al.*, 1996), thought to be mediated by effector molecules that suppress plant immunity to allow the establishment of a successful infection (de Jonge *et al.*, 2011; Kloppeholz *et al.*, 2011). *P. indica* effectors were also described as key players in the formation of a thriving interaction with the plant host (Lahrmann & Zuccaro, 2012; Lahrmann *et al.*, 2013).

4.5 Different approaches to explore possible genetic commonalities of biotrophic interactions

Despite the loss of symbiosis-related genes, *A. thaliana* retained vestiges of the symbiosis program as homologs of CSGs (HCSGs), such as *POLLUX*, *SYMRK-homologous Receptor-like Kinases (ShRKs)*, and members of the NUP107-160 subcomplex, including *SEC13* and *NUP133*.

This fact raises questions regarding the evolutionary forces driving the retention of this core program and suggests that, despite differences among *P. indica* and AM symbioses, their genetic requirements could be overlapping. The *POLLUX* ortholog in

A. thaliana has already been shown not to be required for the *P. indica*-induced growth promotion (Shahollari *et al.*, 2007). Nevertheless, the study did not exclude that the gene is required for other aspects of the *P. indica* symbiosis, such as intraradical colonization *per se*. Furthermore, the transgenic complementation of a symbiosis-defective legume mutant by expression of *AtPOLLUX* variants suggested an overall functional conservation (Venkateshwaran *et al.*, 2012). This finding, together with the fact that HCSGs have been retained in the *A. thaliana* genome, and considering the structural and functional commonalities between the intracellular accommodation organs of biotrophic microorganisms in symbiosis and disease, led us to speculate that the partial retention of an ancestral symbiotic program in *A. thaliana* could be related to a selection pressure imposed by biotrophic interactions other than AM symbiosis, such as the interaction with *P. indica* itself. The fungus grows inside living plant cells without forming any characteristic accommodation organ. Nevertheless, a plant-derived membrane envelops its hyphae during the biotrophic phase (Jacobs *et al.*, 2011; Lahrman & Zuccaro, 2012; Lahrman *et al.*, 2013) and possibly creates an interface between fungal and plant membrane comparable to the peri-fungal space surrounding transcellular hyphae of AM fungi, where signal and nutrient exchange is likely to take place (Bonfante & Genre, 2010).

The oomycetal pathogen *Hpa* is responsible for the downy mildew of *A. thaliana* (reviewed in (Slusarenko & Schlaich, 2003)). Though it causes neither a highly destructive disease on the plants nor is economically important, the *Hpa-A. thaliana* association is a particularly well-studied plant-oomycete pathogen system that was established in the laboratories in the 1990's, and has been of great importance for the identification and cloning of disease resistance genes (Slusarenko & Schlaich, 2003; Heidel & Dong, 2006; Baxter *et al.*, 2010; Coates & Beynon, 2010; Fabro *et al.*, 2011; Stassen & Van den Ackerveken, 2011), and, more recently, of compatibility factors, which facilitate the colonization of the plant host or promote disease susceptibility by biotrophs without leading to exacerbated activation of known defence responses (Huibers *et al.*, 2009). Such factors have also been shown to play a role in the association between plants (including *A. thaliana*) and powdery mildew fungi, another group of biotrophic pathogens (Adam *et al.*, 1999).

4.6 The plant immune system

Different from mammals, which present both an innate immune system as well as a somatic adaptive immune system and mobile defender cells (Litman *et al.*, 2010), plants depend exclusively on their innate immune system to defend themselves against invading pathogens. The plant innate immune system that has been classically divided into two branches (Jones & Dangl, 2006). The first layer of defence is known as PAMP (pathogen associated molecular pattern)-triggered immunity (PTI), in which transmembrane pattern recognition receptors recognize PAMPs and trigger the defence responses. However, many pathogens have evolved the so-called effector proteins, which interfere with host responses and enables them to evade PTI (Dou & Zhou, 2012). To neutralize the activity of effectors, plants evolved mechanisms to recognize effectors using resistance (*R*) genes, most of which encode nucleotide-binding leucine-rich repeat (NB-LRR) proteins (Jones & Dangl, 2006; van Schie & Takken, 2014). The *R*-gene-mediated immunity consists of a more rapid and amplified layer of defence known as ETI (effector triggered immunity) (Jones & Dangl, 2006). This arms race between plant and pathogens results in a continuous process of co-evolution: while pathogens are constantly evolving new strategies, such as new effectors, to evade or suppress plant immune responses, the plants continuously expand their receptor repertoire to recognize molecules associated with pathogen invasion (Jones & Dangl, 2006).

Even though PTI and ETI are triggered by different mechanisms, their outcomes vary mostly in the intensity of the responses (van Schie & Takken, 2014), which are characterized by a variety of symptoms, the most extreme being cell death induction during the hypersensitive response (HR). Other symptoms include accumulation of reactive oxygen species (ROS), production of defence molecules such as salicylic acid (SA) and jasmonate (JA), cell wall strengthening, and activation of defence gene expression (Jones & Dangl, 2006; Muthamilarasan & Prasad, 2013).

A few *A. thaliana* mutants with improved pathogen resistance, like *edr1*, *cpr1*, *sncl*, *ssi4*, *rin4*, and *cpr5*, show deregulated, often constitutively activated immune responses, usually presenting disease symptoms even in the absence of any trigger (Bowling *et al.*, 1994; Adam *et al.*, 1999; Li *et al.*, 2001; Shirano *et al.*, 2002; Lorrain *et al.*, 2003; Belkhadir *et al.*, 2004; Landoni *et al.*, 2013). Another class of more

resistant mutants, which includes *pskr1*, shows enhanced induction of defence responses after PAMP treatment or pathogen challenge (Frye & Innes, 1998; Igarashi *et al.*, 2012; Mosher *et al.*, 2013).

Interestingly, in *A. thaliana*, components of the NUP107-160/NUP84 sub-complex, which comprise some of the HCSGs, have been shown to play roles in plant defence; mutations in NUP96, NUP160 and SEH1 impair innate immune responses and resistance-gene mediated defence (Zhang & Li, 2005; Wiermer *et al.*, 2012).

4.7 Role of compatibility factors in biotrophy

When a pathogen or, for that matter, any other microorganism, is able to overcome plant defence responses and undergo long-lasting interactions with their host, which in the case of biotrophic partners generally involve the formation of specialized feeding structures, a situation of compatibility takes place (reviewed in (van Schie & Takken, 2014)). Compatibility (also referred to as susceptibility) factor candidates have been described to have specific roles in the plant host, such as in cell wall composition, cell cycle progression, vesicle trafficking, cytoskeleton functionality, and detoxification. These factors are often manipulated by the pathogens to ultimately allow for a successful colonization, and in favour of its own growth (Vogel & Somerville, 2000; Vogel *et al.*, 2004; Haga *et al.*, 2007; Pareja-Jaime *et al.*, 2008). Based on the stage of the interaction the compatibility factors are involved in, three main mechanisms of action have been described: genes involved in facilitating host recognition and cell penetration, genes associated with the negative regulation of plant immunity, and genes required to maintain the interaction, allowing the pathogen to grow and proliferate (reviewed in (van Schie & Takken, 2014)). The first described compatibility factor was the membrane-anchored protein Mildew Resistance *Locus O* (MLO), discovered in barley (Freisleben & Lein, 1942) and shown to be required for susceptibility to powdery mildew in, among others, *A. thaliana*, tomato, and wheat (Consonni *et al.*, 2006; Bai *et al.*, 2008; Varallyay *et al.*, 2012; Zheng *et al.*, 2013). MLO-dependent susceptibility requires vesicle trafficking as well as actin polarization (Consonni *et al.*, 2006; Miklis *et al.*, 2007). In barley, the protein HvRIC171 is recruited by the powdery mildew fungus *B. graminis* to the site of infection and supports fungal penetration (Schultheiss *et al.*, 2008). The powdery mildew fungus

Golovinomyces cichoracearum has been shown to modify plant cell wall components encoded by the genes *POWDERY MILDEW RESISTANT 4* (*PMR4*, a callose synthase), *PMR5* (unknown function), and *PMR6* (pectate lyase) to facilitate entrance into the cell without activating defence responses, and the corresponding mutants are therefore less susceptible to the infection (Vogel & Somerville, 2000; Vogel *et al.*, 2002; Vogel *et al.*, 2004). The mutant *pmr4* is less susceptible to *Hpa* as well (Vogel & Somerville, 2000). Several genes involved in amino acid metabolism have been reported to have a role in the *Hpa*-*A. thaliana* compatibility; *DOWNY MILDEW RESISTANT 1* (*DMR1*), for instance, encodes a homoserine kinase whose absence results in reduced susceptibility to *Hpa* (Van Damme *et al.*, 2005; van Damme *et al.*, 2009). A recent analysis of the *A. thaliana* susceptibility transcriptome was performed on laser-dissected *G. orontii* infection sites, revealing that the fungus induces genetic reprogramming of the host cells to turn them into sources of nutrients (Chandran *et al.*, 2010). In this study, the expression of *MYB3R4*, encoding a transcription factor involved in cell cycle progression, was upregulated at the infection sites, with the cells undergoing endoreduplication events that result in an increased metabolic capacity.

In this work, we investigated the role of symbiosis-related genes in the interaction of hyphal microorganisms with *L. japonicus* and *A. thaliana*. In *L. japonicus*, the CSGs were assessed for involvement in the intracellular colonization by the root endophyte *P. indica*, while in *A. thaliana* the involvement of HCSGs in the fungal-mediated host growth promotion or their roles as possible compatibility factors in biotrophic interactions with *P. indica*, the oomycete *Hpa* and the powdery mildew fungus *E. cruciferarum* were investigated. To test for specificity to biotrophic associations that involve the formation of intracellular accommodation structures in *A. thaliana*, we included in our analysis the extracellular bacterial pathogen *Pseudomonas syringae* pv. tomato *DC3000*, which, contrary to *Hpa* and *E. cruciferarum*, grows in the apoplast without entering living plant cells.

5. Results

5.1 *A. thaliana* HCSG mutants reduce the reproductive success of the oomycete *Hpa*

We inspected the *A. thaliana* genome for the presence of HCSG and identified candidate orthologs of *POLLUX* and of the nucleoporins of the NUP107-160 subcomplex (Figures 2 + 3). A direct ortholog of *SYMRK*, a malectin-like domain leucine-rich repeat receptor-like kinase (MLD-LRR-RLK) (Antolin-Llovera *et al.*, 2014; Ried *et al.*, 2014) was deleted from the syntenic position in the *A. thaliana* genome (Kevei *et al.*, 2005). Nevertheless, the MLD-LRR-RLK protein subfamily is also present in *A. thaliana*. In fact, a specific expansion of this gene family took place in this plant (Shiu & Bleecker, 2003). We performed phylogenetic analyses with members of the *A. thaliana* MLD-LRR-RLK subfamily, *LjSYMRK*, and two additional MLD-LRR-RLKs from *L. japonicus* using the protein sequences of their kinase domain alone or only the extracytoplasmatic region. The clustering of the kinase domain sequences was similar to that of the whole (pruned) amino acid sequences previously published (Hok *et al.*, 2011). However, an analysis of the extracytoplasmatic region resulted in different tree topologies. For instance, while in the first tree the At5g48740 protein (which has an additional LRR domain) was identified as the closest related to *LjSYMRK*, in the second it clustered in a separate group. However, in both cases, the products of two *A. thaliana* genes, which we thus named *SYMRK-homologous Receptor-like Kinase 1* (*ShRK1*) and *ShRK2*, were identified as the most closely related to *LjSYMRK* (this work and (Markmann *et al.*, 2008)).

All investigated *A. thaliana* HCSGs have identical exon-intron structure and domain organization to their CSG counterparts (Figure 2). By sequencing a PCR product amplified from *A. thaliana* Col-0 cDNA, we demonstrated that, contrary to the TAIR prediction, this was also the case for *NUP133*. The curated sequence has been submitted to the GenBank (accession number KM269292).




Figure 2: Comparison of gene structures and protein domains of *L. japonicus* common symbiosis genes (CSGs) and their closest homologs (HCSGs) in *A. thaliana*. Complete annotated genomic sequences were obtained from The Arabidopsis Information Resource (TAIR – www.arabidopsis.org) for *A. thaliana*, and from the KDRI website (Kazusa DNA Research Institute, Japan; <http://www.kazusa.or.jp/lotus/>) and the GenBank for *L. japonicus*. TAIR/GenBank protein identifiers are shown; dashed lines, positions of the introns in the original gene sequence; red triangles, positions of the T-DNA insertion in the respective mutants; SP, signal peptide; MLD, malectin-like domain; LRRs, leucine-rich repeats; TM, transmembrane domain; KD, kinase domain.

Gene	<i>L. japonicus</i> homolog (NCBI mRNA/protein IDs)	Gene ID	NASC ID	Protein Identity / Similarity To <i>L. japonicus</i>
<i>POLLUX</i>	AB162158 / BAD89022	At5g49960	N566135	71 / 82
<i>SEC13</i>	AB506697 / BAJ10727 - AB506698 / BAJ10728	At3g01340	N662322	80-82 / 92-91
<i>NUP133</i>	AJ890251 / CAI64810	At2g05120	N565761	55 / 71
<i>SEH1</i>	AB506696 / BAJ10726	At1g64350	N653094	64 / 78
<i>NUP43</i>	n.d.	At4g30840	N803490	n.d.
<i>NUP85</i>	AB284835 / BAF45348	At4g32910	N613274	61 / 75
<i>NUP160</i>	n.d.	At1g33410	N624418	n.d.
<i>ShRK1</i>	(<i>SYMRK</i>)	At1g67720	N467036	33 (48KD) / 49 (68KD)
<i>ShRK2</i>	AF492655 / AAM67418	At2g37050	N643700	34 (48KD) / 50 (68KD)

Figure 3: *A. thaliana* HCSGs with encoded proteins and their respective identities/similarities to their *L. japonicus* homologs. Sequence identifiers and identities / similarities shared between the protein sequences of *A. thaliana* HCSGs and those of their respective *L. japonicus* counterparts. The numbers for AtSEC13 indicate the identity/similarity of its amino acid sequence to each of the two predicted LjSEC13 proteins. For ShRK1 and ShRK2, numbers in brackets refer to their kinase domain (KD) only. n.d., not detected. NASC ID: insertion mutant identifier.




Figure 4: Maximum likelihood phylogenetic trees of MLD-LRR-RLKs from *A. thaliana* and *L. japonicus*. Maximum likelihood phylogenetic trees were constructed using the highly conserved kinase domains (upper tree) or the extracytoplasmatic regions minus signal peptide (lower tree) of MLD-LRR-RLK proteins from *A. thaliana*, where this family underwent a recent expansion (Shiu & Bleecker, 2003), and *L. japonicus*. Numbers on each node represent the respective bootstrap values. Scale bar = relative genetic distance (arbitrary unit).

Phylogenetic analyses were performed by Aline Banhara Pereira and Martina Katharina Ried. Aline Banhara Pereira prepared the figure.

Insertion mutant lines for *pollux*, *sec13*, *nup133*, *shrk1*, *shrk2*, and the double mutants *sec13* x *nup133* and *shrk1* x *shrk2* were analysed for their phenotype in the interaction with *Hpa*. In *pollux*, *shrk1*, *shrk2*, the double mutant *shrk1* x *shrk2*, and the disease resistant reference mutant *pskr1* (Mosher *et al.*, 2013), the reproductive success of *Hpa* isolate NoCo2, measured as sporangiophore number per cotyledon 4 days post infection (dpi), was significantly reduced and was restored in the available complementation lines (Figures 5 + 6).

To capture the potential structural or functional shifts of nucleoporins during evolution, we included a range of NUP107-160 subcomplex members in our analysis. The *sec13* and *nup133* single mutants and the double mutant *sec13* x *nup133* impaired *Hpa* reproductive success (Figures 5 + 6), while *seh1*, *nup43*, *nup85* and *nup160* did not (Figure 7). This pattern is not congruent with the observation in *L. japonicus*, in which *seh1*, *nup85* and *nup133* impaired symbiosis (Kanamori *et al.*, 2006; Alber *et al.*, 2007; Saito *et al.*, 2007), and may be explained by species-specific adaptations of the NUP107-160 subcomplex.




Figure 5: Mutation of *A. thaliana* HCSGs reduces the reproductive success of *Hpa*. Plots show the number of sporangiophores per infected cotyledon on ca. 50 cotyledons of *A. thaliana* wild-type (Col-0), the indicated mutants, and transgenic complementation lines (co) 4 dpi with *Hpa* isolate NoCo2. Dots: outliers. Numbers above upper whiskers indicate the values of individual outliers outside of the plotting area. Stars indicate significant differences to Col-0 (Wilcoxon–Mann–Whitney test with Bonferroni-Holm correction). ∞ , $p = 0.067$; *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$.

Constructs *pollux co*, *shrkl co*, and *shrk2 co* were produced by Andreas Binder. Experiments were designed, performed, and analysed by Aline Banhara Pereira and Martina Katharina Ried. Martina Katharina Ried did the statistical analyses and prepared the figure.




Figure 6: Reproductive success of *Hpa* is reduced in the HSCG mutants. Plots represent the number of sporangiophores per infected cotyledon on ca. 50 cotyledons of *A. thaliana* wild-type (Col-0) and the HSCGs mutants 4 dpi with *Hpa* isolate NoCo2. Two replicates are shown for each mutant set. Dots: outliers. Numbers above upper whiskers indicate the values of individual outliers outside of the plotting area. Significant differences to the wild-type (Col-0) were detected at the 5% significance level (Wilcoxon–Mann–Whitney with Bonferroni–Holm correction), except for *sec13* in the lower set, for which p was 0,065 (indicated with ∞). $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$.

Experiments were designed, performed, and analysed by Aline Banhara Pereira and Martina Katharina Ried. Martina Katharina Ried did the statistical analyses and prepared the figure.




Figure 7: Reproductive success of *Hpa* is not affected in the *seh1*, *nup43*, *nup85*, and *nup160* mutants. Plots represent the number of sporangiophores per infected cotyledon on ca. 50 cotyledons of *A. thaliana* wild-type (Col-0) and the indicated mutants 4 dpi with *Hpa* isolate NoCo2. Open circle: outlier. No significant differences to the wild-type (Col-0) were detected at the 5% significance level (Wilcoxon–Mann–Whitney test with Bonferroni–Holm correction).

Experiment was designed and performed by Aline Banhara Pereira, and analysed by Aline Banhara Pereira and Martina Katharina Ried. Martina Katharina Ried did the statistical analyses and prepared the figure.

5.2. Haustorial development of *Hpa* is impaired in *A. thaliana* HCSG mutants

The reduced *Hpa* reproductive success could not be explained by a decreased frequency of haustoria formation. Other than a slight decrease in *pollux*, this frequency in the other mutants was indistinguishable from the wild-type (Figure 8). However, all *A. thaliana* HCSG mutants exhibited strikingly altered haustoria morphology. At 5 dpi the majority of haustoria in the wild-type had a globular, single-lobed appearance. Deviations from such morphology, which we generally called multilobed, were observed as well. The amount of multilobed haustoria in the HCSG mutants was significantly increased, a phenomenon that was alleviated in the available complementation lines (Figures 9 +10 + 11 + 12). In contrast to the HCSG mutants, the reference disease resistant mutant *pskr1*, which presents resistance to

disease due to constitutively activated defence responses (Mosher *et al.*, 2013), did not show any signs of altered haustorial development (Figure 9).




Figure 8: *Hpa* haustoria formation on *A. thaliana* HCSG mutants. Plots represent the mean percentage of haustoria-containing cells per cells contacted by hyphae on 5 leaves of *A. thaliana* wild-type (Col-0), the indicated mutants and transgenic complementation lines (co) at 5 dpi with *Hpa* isolate NoCo2. Ten independent hyphal strands were analysed on each leaf. Black circles: outliers. Stars indicate significant differences to Col-0 (Dunnett's Test with Bonferroni correction). *** $p < 0.001$.

Construct *pollux* co was produced by Andreas Binder. Experiment was designed, performed, and analysed by Aline Banhara Pereira and Martina Katharina Ried. Martina Katharina Ried did the statistical analyses and prepared the figure.





Figure 9: *Hpa* haustorium morphology is altered in *A. thaliana* HCSG mutants. *A. thaliana* wild-type (Col-0), the indicated HCSG mutants, and transgenic complementation lines (co) were analysed 5 dpi with *Hpa* isolate NoCo2. Upper panel: Plots show the percentage of multilobed haustoria among total haustoria. For each genotype, 5 leaves were analysed, and 10 hyphal strands with 15-30 haustoria each were counted on each leaf. Dots: outliers. Stars indicate significant differences to Col-0 (Wilcoxon–Mann–Whitney test with Bonferroni–Holm correction). *, $p < 0.05$. Lower panel: Representative pictures of hyphal strands that grow intercellularly in the mesophyll and intracellular haustoria. Stars indicate multilobed haustoria. Scale bars = 25 μm .

Constructs *pollux co*, *shrkl co*, and *shrk2 co* were produced by Andreas Binder. Experiments were designed, performed, and analysed by Aline Banhara Pereira and Martina Katharina Ried. Martina Katharina Ried did the statistical analyses and prepared the figure.




Figure 10: Frequency of multilobed haustoria on *A. thaliana* wild-type and on the HSCG mutants. *A. thaliana* wild-type (Col-0) and the HSCG mutants were analysed 5 dpi with *Hpa* isolate NoCo2. Plots show the percentage of multilobed haustoria among total haustoria. For each genotype, 5 leaves were analysed, and 10 hyphal strands with 15-30 haustoria each were counted on each leaf. Black dots: outliers. Stars indicate significant differences to Col-0 (Wilcoxon–Mann–Whitney test with Bonferroni-Holm correction). For *nup133*: $p = 0,151$. *, $p < 0.05$.

Experiment was designed, performed, and analysed by Aline Banhara Pereira and Martina Katharina Ried. Martina Katharina Ried did the statistical analyses and Aline Banhara Pereira prepared the figure.




Figure 11: *Hpa* haustorium morphology is altered in *A. thaliana* HCSG mutants. Differential interference contrast microscopy of leaves of *A. thaliana* wild-type (Col-0) and HCSG mutants stained with trypan-blue lactophenol 5 dpi with *Hpa* isolate NoCo2. Scale bar = 25 μM.

Experiments was designed, performed, and analysed by Aline Banhara Pereira and Martina Katharina Ried. Aline Banhara Pereira prepared the figure.




Figure 12: *Hpa* haustoria morphology. *Hpa* haustoria on wild-type *A. thaliana* (Col-0) leaves (upper row) 5 dpi, with the regular globular or pear-like morphology, and examples of multilobed haustoria observed in the *shrk1 x shrk2* double mutant (lower row). Leaves were stained with anilin-blue and visualized with a CLSM. For every haustorium, the entry point can be identified by the formation of the usually bright callose neck. In the double mutant, multiple lobes are visualized forming in individual haustoria. Scale bar: 25 μ M.

Experiment was designed, performed, and analysed by Aline Banhara Pereira and Martina Katharina Ried. Aline Banhara Pereira prepared the figure.

In both the wild-type and in *shrk1 x shrk2*, the percentage of multilobed haustoria increased over time, but was significantly higher in *shrk1 x shrk2* at each analysed time point (Figure 13), suggesting early haustoria aging in the double mutant. An altered morphology was also observed for *Hpa* haustoria in leaves of transgenic *A. thaliana* plants expressing the N-terminally YFP-tagged remorin AtREM1.2 (At3g61260) under the control of its native promoter (ProAt3g61260:YFP-At3g61260, (Jarsch *et al.*, 2014)). Similar to its homolog AtREM1.3 (Bozkurt *et al.*,

2014), AtREM1.2 localizes to the perihastorial membrane. We observed that the loss of ProAt3g61260:YFP-At3g61260 fluorescence was associated with changes in the haustorium morphology, perhaps indicating an age-related alteration of the protein composition of the perihastorial membrane (Figure 14). The impaired haustorial development in the HCSG mutants may decrease nutrient availability to the oomycete, with a consequent reduction in sporangiophore production. Collectively, the remarkably specific phenotype of altered haustorial development on all tested *A. thaliana* HCSG mutants pinpoints these genes as contributing to a program for the intracellular accommodation of this oomycetal pathogen. Interestingly, on the HCSG mutants we did not observe a consistent reduction in reproductive success of the powdery mildew fungus *Erysiphe cruciferarum* (Figure 15), which forms haustoria exclusively in epidermal cells. A morphological comparison of haustoria shape was not possible because of the highly variable haustorial morphology already in the wild-type interaction. It is therefore possible that the HCSGs do not play a role in this interaction, due to alternative pathways in the epidermis or distinct genetic requirements for the colonization between fungal and oomycetal intracellular pathogens. The *MLO* gene, for instance, is an epidermal compatibility factor required for powdery mildew fungus penetration (Consonni *et al.*, 2006), with no role in the *Hpa* infection reported to date.




Figure 13: Time-course of haustoria development in *A. thaliana* wild-type and in *shrk1 x shrk2*. Plots represent the mean percentage of multilobed haustoria per total haustoria on 5 leaves of *A. thaliana* wild-type (Col-0) and the *shrk1 x shrk2* double mutant 4, 5 and 7 dpi with *Hpa* isolate NoCo2. On each leaf, 10 independent hyphal strands were analysed. Black circle: outlier. Stars represent significant differences (Wilcoxon–Mann–Whitney test with Bonferroni-Holm correction). **, $p < 0.01$.

Experiment was designed, performed, and analysed by Aline Banhara Pereira and Martina Katharina Ried. Martina Katharina Ried did the statistical analyses and prepared the figure.





Figure 14: Quantification of haustoria with fluorescence on *ProAt3g61260:YFP-At3g61260* plants.

A. thaliana plants expressing N-terminally YFP-tagged remorin AtREM1.2 under the control of its native promoter were infected with *Hpa* and inspected 8 dpi with an epifluorescence microscope. Quantification of haustoria with fluorescence on the perihaustral membrane region was performed in triplicate. For each replicate consisting of 6 leaves, 100 single-lobed and 100 multilobed haustoria were counted and scored for detectable fluorescence. Exemplary images (differential interference contrast, fluorescent, merge) of labelled and non-labelled (completely dark) haustoria are shown. Yellow dashed lines delimitate the haustoria in the bright field images. Graph represents the percentage of haustoria with fluorescence as a function of morphology; distinct symbols refer to different replicates (1st replicate: 49% single-lobed/19% multilobed; 2nd replicate: 56% single-lobed/17% multilobed; 3rd replicate: 68% single-lobed/14% multilobed). Bar: 25 μ M.




Figure 15: Mean number of conidiophores per leaf on *A. thaliana* HCSG mutants relative to the wild-type. Box-plots represent a compilation from four independent replicates and show mean number of conidiophore on HCSG mutant relative to the wild-type (Col-0) leaves 5 dai with 3-4 *E. cruciferarum* spores/mm². For each replicate, conidiophores/colony were counted on 10 colonies per leaf, on 5-10 leaves/genotype. No significant differences to the wild-type (Col-0) were detected at the 5% significance level (Dunnett's Test with Bonferroni-correction).

Experiment was designed and performed by Caroline Höfle, and analysed by Aline Banhara Pereira and Martina Katharina Ried. Martina Katharina Ried did the statistical analyses and prepared the figure.

5.3. *A. thaliana* HCSG mutants do not exhibit constitutive or enhanced defence responses

To investigate whether the reduced oomycete fitness on the HCSG mutants resulted from constitutively activated immune responses or from elevated defence responses upon pathogen perception, we first analysed the transcript levels of PAMP-induced marker genes under non-challenged conditions and 6 hours after flg22 treatment (Figures 16 + 17). The basal expression of *FLG22-INDUCED RECEPTOR-LIKE KINASE 1 (FRK1)* (Asai *et al.*, 2002), *GLUTATHIONE S-TRANSFERASE 1 (GST1)* (Grant *et al.*, 2000), *ETHYLENE RESPONSE FACTOR 1 (ERF1)* (Solano *et al.*, 1998), and *PHOSPHATYDILINOSITOL-LIPID TRANSFER PROTEIN (PI-LTP)* (Denoux *et al.*, 2008) in the HCSG mutants did not differ from that detected in the wild-type or in the *FLAGELLIN SENSING 2 (FLS2)*; (Gomez-Gomez & Boller, 2000)) mutant (Figures 16 + 17). Moreover, 6 hours after flg22 treatment these genes were all upregulated in the HCSG mutants to values similar to the wild-type (Figures 16 + 17).




Figure 16: Basal expression levels and flg22-mediated induction of PAMP-responsive genes in *A. thaliana* HCSG mutants are not different from the wild-type. The relative expression of *FRK1* and *GST1* in mock-treated samples or in samples treated with 1 μ M flg22 for 6h (+) was determined in three biological replicates for each genotype by qRT-PCR. The *fls2* mutant was the negative control. Transcript levels for every plant genotype and each treatment were determined with technical duplicates. Closed circle, mock-treated; open circle, flg22 treated. Stars label datasets significantly different from Col-0 (Dunnett's Test with Bonferroni correction). *, $p < 0.01$; **, $p < 0.001$.

Experiments were designed, performed and analysed by Aline Banhara Pereira and Martina Katharina Ried. Martina Katharina Ried did the statistical analyses and prepared the figure.




Figure 17: Basal expression levels and flg22-mediated induction of PAMP-responsive genes in *A. thaliana* HCSG mutants are not different from the wild-type. Relative expression of *ERF1* and *PI-LTP* in mock-treated samples (-) or in samples treated with 1 mM flg22 for 6h (+) were determined in three biological replicates for each genotype via qRT-PCR. The *fls2* mutant was used as negative control. Transcript levels for every plant genotype and each treatment were determined through technical duplicates. Black circle, mock-treated; open circle, flg22 treated. Stars indicate significant differences to Col-0 (Dunnett's Test with Bonferroni correction). *, p < 0.01.

Experiments were designed, performed and analysed by Aline Banhara Pereira and Martina Katharina Ried. Martina Katharina Ried did the statistical analyses and prepared the figure.

Furthermore, the constitutive transcript levels of *PATHOGENESIS-RELATED GENE 1 (PR1)*, a marker gene for salicylic acid (SA)-mediated resistance (Ryals *et al.*, 1996), which is activated upon infection by biotrophic pathogens (Glazebrook, 2005), were not significantly increased in the HCSG mutants (Figure 18). In all cases, deviations observed for individual mutants were not consistent through the gene set and are thus unlikely responsible for the increased pathogen resistance.




Figure 18: Transcript levels of defence marker gene *PR1* in *A. thaliana* HCSG mutants. Relative expression of *PR1* was determined in three mock-treated biological replicates for each genotype via qRT-PCR. Transcript levels for every plant genotype were determined through technical duplicates. No statistical differences to the wild-type (Col-0) were obtained at the 1% significance level (Dunnett's Test with Bonferroni correction).

Experiments were designed, performed and analysed by Aline Banhara Pereira and Martina Katharina Ried. Martina Katharina Ried did the statistical analyses and prepared the figure.

Symptoms typically associated with deregulated immune responses, such as constitutive or pathogen-induced hypersensitive response (HR) or defects in growth and development due to the hyper-activation of the SA-dependent defence pathway (Bowling *et al.*, 1997), were absent in the *A. thaliana* HCSG mutants (Figure 19 + 20). In addition, the ability of *Hpa* to suppress callose deposition around the haustorial neck region (Sohn *et al.*, 2007) was not disturbed (Figure 21). The NUP107-160 subcomplex has also been implicated in plant defence; mutations in *NUP96*, *NUP160* and *SEHI* impair basal and resistance-gene mediated immunity (Zhang & Li, 2005; Wiermer *et al.*, 2012). However, our mutants did not show such deregulation, possibly due to distinct roles of individual subcomplex components in compatibility and defence.




Figure 19: *A. thaliana* HCSG mutants do not show increased levels of spontaneous or pathogen-induced cell death. Upper panel: differential interference contrast microscopy of representative *A. thaliana* wild-type (Col-0) and HCSG leaves 4 dpi with *Hpa* isolate NoCo2. Most of the non-infected leaves display no sign of cell death (first column), but dark-blue stained dead cells are sporadically observed in non-infected leaves of both wild-type and HCSG mutants (arrows, second column). In infected leaves of the HCSG mutants, cell death is occasionally detected randomly on the leaf surface (third column) and in, or adjacent to, haustoria-containing cells in a frequency indistinguishable from the wild-type (fourth column). In all genotypes, infected leaves contain hyphal strands growing in the absence of any cell death (fifth column). Scale bar = 25 μ m. Lower panel: plots show the mean number of random (left) or *Hpa*-associated (right) cell death spots per leaf on ca. 50 leaves per genotype 5 dpi with *Hpa* isolate NoCo2. Open circles: outliers. For statistical analysis, a one-way ANOVA followed by a Tukey's HSD was performed. Different letters indicate samples that are significantly different at the 5% significance level. Results were obtained from two independent experiments.

Experiments were designed, performed, and analysed by Aline Banhara Pereira and Martina Katharina Ried. Statistical analyses were done by Martina Katharina Ried. Aline Banhara Pereira prepared upper panel and Martina Katharina Ried prepared lower panel.



Figure 20: *A. thaliana* HCSG mutants do not show developmental or growth defects. 3-week-old *A. thaliana* wild-type (Col-0) plants grown alongside the indicated mutant lines under long day conditions (16h light). The dwarf phenotype of the mutant suppressor of *npr1-1*, *constitutive 1* (*snc1*; (Li *et al.*, 2001)) is included on the far right for comparison.




Figure 21: *Hpa* haustoria-associated callose deposition is not altered in the *A. thaliana* HCSG mutants. Plot shows the mean intensity of callose deposition on a total of ca. 50 oomycete penetration sites from 5 different leaves of *A. thaliana* wild-type (Col-0) and the HCSG mutants 4 dpi with *Hpa* isolate NoCo2. Black circles: outliers. For statistical analysis, a one-way ANOVA followed by a Tukey's HSD was performed. No significant differences were obtained at the 5% significance level. Representative pictures of haustoria-associated callose deposition on *Hpa* hyphae growing on the indicated mutants are shown on the right. Scale bar = 25 μ m.

Experiment was designed and performed by Aline Banhara Pereira, and analysed by Aline Banhara Pereira and Martina Katharina Ried. Statistical analyses were done by Martina Katharina Ried. Aline Banhara Pereira prepared the right panel, and Martina Katharina prepared the left panel.

5.4. Growth kinetics of *Pseudomonas syringae* are not altered on the *A. thaliana* HCSG mutants

The bacterial pathogen *Pseudomonas syringae* DC3000 induces the activation of the SA- and the jasmonic acid (JA)-dependent defence signaling in the host (Farmer *et al.*, 2003; Thaler & Bostock, 2004), and overshooting activation of those pathways leads to increased *P. syringae* resistance. The growth of DC3000 wild-type or the avirulent Δ *AvrPto/PtoB* strain (Figure 22) was unaltered on the *A. thaliana* HCSG mutants, providing further evidence that they do not exhibit constitutive or enhanced activation of SA- and JA-dependent defences.




Figure 22: *A. thaliana* HCSG mutants support similar bacterial growth curves as the wild-type. Growth of *Pto* DC3000 (left) or *Pto* DC3000 $\Delta AvrPto/AvrPtoB$ (right) in *A. thaliana* wild-type (Col-0) or the HCSG mutants *pollux*, *shr1 x shr2* and *nup133 x sec13* was determined 2 or 4 days post infiltration of 10^4 colony forming units ml^{-1} (cfu/cm²). Data represent means \pm S.D. of six replicate measurements/genotype/data point. Results from one representative of at least four independent experiments are shown.

Experiments were designed, performed, and analysed by Andrea A. Gust and Thorsten Nürnberger. Figure was prepared by Andrea A. Gust.

5.5. *P. indica* colonization of *L. japonicus* roots occurs independently of CSGs

We investigated whether the classical CSGs are involved in the interaction between roots of the legume *L. japonicus* and *P. indica*. Gifu wild-type plants and common symbiosis mutants derived from this ecotype were analysed with regard to the ability of the fungus to penetrate root epidermal cells, undergo a biotrophic phase, and grow and sporulate inside these cells. For that, we inoculated *L. japonicus* roots with *P. indica* chlamydospores and performed microscopic analyses to assess colonization at 7 dpi and the presence of fungal spores at 14 dpi within cleared roots. Although fungal hyphae were clearly detected within the boundaries of root epidermal cells (Figure 23), it was not clear whether this was the result of biotrophic or necrotrophic growth. We therefore used the vital stain FM4-64 to label plant membranes as a proxy for the vitality status of the root cells. Using this technique we obtained clear evidence for *P. indica* hyphae within living root epidermal cells at 3 dpi. Intracellular biotrophic hyphae were detected in the wild-type and in the common symbiosis mutants (Figure 24). Moreover, intracellular sporulation was observed in roots of all genotypes (Figure 25), evidencing that the *P. indica* successfully completed its life cycle within the roots of the common symbiosis mutants. The fact that intracellular uptake and maintenance of *P. indica* hyphae occurred independent of the common symbiosis genes indicates that alternative pathways must exist for the biotrophic accommodation of *P. indica*.




Figure 23: Intracellular growth of *P. indica* in root cells of *L. japonicus*. Wild-type and the indicated common symbiosis mutants were analysed 7 dpi with *P. indica* chlamydospores. Intracellular hyphae were present in all genotypes (examples are indicated by asterisks). Roots were cleared and double stained with propidium iodide (red), for cell wall visualization, and WGA-AF488 (green), for fungal structures. Scale bar: 25 μm .




Figure 24: Biotrophic growth of *P. indica* in root cells of *L. japonicus*. Microscopy of wild-type and the indicated common symbiosis mutants was performed 3 dpi with *P. indica* chlamydospores. FM4-64-stained endomembranes (red) inside root cells indicates cell viability. Examples of biotrophic hyphae (delimited by dotted lines on the second and fourth columns) are shown within rhizodermal cells. In some cells, the FM4-64-stained membrane surrounding the invading hyphae is well detectable. While extracellular hyphae are stained with WGA-AF488 (green), biotrophic invasive hyphae are not or weakly fluorescent probably due to limited access of the reagent to the fungal cell wall in the fresh root cells. Confocal microscopic pictures were taken 3 dpi with *P. indica* chlamydospores with a Leica TCS-SP5 CLSM. Scale bar: 10 μ m.




Figure 25: *P. indica* sporulation inside root cells of *L. japonicus*. Single and multiple spores are visible inside individual cleared root cells or root hairs of *L. japonicus* wild-type (Gifu) and in the indicated common symbiosis mutants at 14 dpi. Left column: differential interference contrast (DIC) microscopy; right column: WGA-AF488 (WGA) detected by fluorescence microscopy. Scale bar: 25 μ m.

We analysed the relative content of fungal hyphae in roots by determining the ratio of fungal DNA to plant DNA. Interestingly, we detected a tendency for an increased relative amount of fungal DNA in most common symbiosis mutants, with a significant difference to the wild-type in *nup85-1*, *ccamk-12*, and *cyclops-3*, indicating that the fungal biomass in these mutant roots was higher than in the wild-type (Figure 26).




Figure 26: *P. indica* colonization in *A. thaliana* and *L. japonicus* wild-type and mutant roots. Real-time qPCR analysis were performed with DNA from surface-washed *P. indica* colonized roots using primers for the fungal gene *Transcription Elongation Factor (TEF)* and for *A. thaliana* and *L. japonicus* *Ubiquitin (UBI)* genes. Bars show ratios of fungal to plant DNA in 200 mg (ca. 10 plants) per sample of *L. japonicus* and *A. thaliana* roots colonized by *P. indica* at 14 dpi grown on modified HO medium. Error bars represent standard errors of the mean from three independent biological repetitions (except *pollux-5*, for which only one replicate was available). Data were analysed with a two-sided, unpaired t-test at the 5% significance level. *, $P < 0.05$; **, $P < 0.01$.

Experiments were designed by Alga Zuccaro and performed by Aline Banhara Pereira and Alga Zuccaro. Aline Banhara Pereira performed the statistical analyses and prepared the figure.

5.6. *P. indica* did not induce β -glucuronidase (GUS) activity in the *L. japonicus* symbiosis reporter line T90

The *L. japonicus* symbiosis-reporter line T90 was isolated in a screen for symbiosis-specific GUS expression from a promoter-tagging population (Webb *et al.*, 2000). The T90 reporter line is activated both in roots treated with nodulation factors and in roots inoculated with *Mesorhizobium loti*. It also shows GUS expression in the interaction with AM fungi (Radutoiu *et al.*, 2003; Kistner *et al.*, 2005). In order to check whether *P. indica* was able to induce the expression of GUS in the T90 line, we inoculated roots with either the *M. loti* DsRed or with *P. indica* chlamydospores, or mock-inoculated with Tween water before histochemical staining with 5-bromo-4-chloro-3-indolyl glucuronide at 3, 7, and 14 dpi. No GUS activity (blue staining) was observed in *P. indica* or mock-inoculated roots. In roots inoculated with *M. loti* DsRed, blue staining was detected in the root at all time points (Figure 27).




Figure 27: GUS expression in the *L. japonicus* symbiosis reporter line T90 inoculated with *P. indica*. At all evaluated time-points, GUS activity was not observed in T90 plants inoculated with *P. indica* chlamydospores or mock-treated with Tween water. *Mesorhizobium loti* DsRed was used as a positive control and activated GUS expression (blue) in the root or root nodules at all time-points analysed. Scale bar: 2.5 mm.

Experiment was designed and performed by Regina Kühner. Aline Banhara Pereira analysed the experiment and prepared the figure.

5.7. Intracellular colonization of *A. thaliana* roots by *P. indica* is independent of HCSGs

Our results from the mutant analysis in *L. japonicus* suggested that alternative plant genes support the infection by *P. indica*. Although the Brassicaceae, including *A. thaliana*, do not form AM and have lost several key CSGs during their evolutionary history, their genomes harbour homologs of CSGs (Delaux *et al.*, 2013; Delaux *et al.*, 2014; Banhara *et al.* unpublished). In order to determine the contribution of these genes to the interaction with *P. indica*, specially during the biotrophic phase, which presents more structural similarities to the root colonization by AM fungi, we carried out cytological analyses in roots of wild-type plants and of the HCSG mutants *pollux*, *sec13*, *nup133*, and the double mutants *sec13* x *nup133* and *shrk1* x *shrk2*. Hyphae growing within cells filled with FM4-64-stained endomembrane structures (Figure 28) were observed in all genotypes, suggesting that the penetrated cells were alive and consequently that *P. indica* is able to establish biotrophic growth in these HCSG mutants. Furthermore, the mutants did not differ in the relative colonization of their roots by *P. indica* at 14 dpi (Figure 26). In addition, we detected intracellular sporulation in rhizodermal cells of all genotypes (Fig. 29), indicating that the fungus could successfully complete its life cycle in the roots of the tested HCSG mutants.




Figure 28: Biotrophic growth of *P. indica* within *A. thaliana* root cells. Intracellular hyphae (indicated by asterisks) are visible within rhizodermal cells of wild-type and the indicated mutants. Extracellular hyphae are stained with WGA-AF488 (green). FM4-64-stained plant membranes (red) are visible inside the plant cells, and in some situations enveloping invading hyphae (arrowheads). Biotrophic intracellular hyphae are not or weakly fluorescent, probably due to limited access of WGA-AF488 to the fungal cell wall within root cells. Confocal microscopic pictures were taken with a Leica TCS-SP5 CLSM 14 dpi with *P. indica* chlamydospores. Lengths of scale bars (in μm) are shown on each image.

Experiments were designed, performed, and analysed by Aline Banhara Pereira, Yi Ding, and Alga Zuccaro. Aline Banhara Pereira prepared the figure.




Figure 29: Intracellular sporulation of *P. indica* in *A. thaliana* roots. Multiple spores per root cell are present in the *A. thaliana* wild-type (Col-0) as well as in the indicated HCSG mutants at 14 dpi. Left column: differential interference contrast (DIC) microscopy. Right column: WGA-AF488 (WGA) detected by fluorescence microscopy. Scale bar: 25 μ m.

5.8. Plant growth promotion by *P. indica* is influenced by nutrient availability

We previously reported growth promotion of *A. thaliana* plants upon co-culture with *P. indica* when grown on modified HO medium (Lahrmann *et al.*, 2013). To further explore the growth conditions supportive for *P. indica*-mediated growth promotion, we evaluated the effect of *P. indica* on *L. japonicus* and *A. thaliana* grown in soil with two different nutrient regimes. In both plant species, co-cultivation with *P. indica* in soil with high nutrient concentrations had little or no effect on the mean stem height. However, in soil with lower nutrient contents, *P. indica* inoculation roughly doubled the mean stem height of *A. thaliana* plants by 21 dpi, whereas there was only a small however significant effect on *L. japonicus* (Figure 30, Table 1). In general, the positive effect of co-cultivation with *P. indica* in low-nutrient soil was stronger on *A. thaliana* than in *L. japonicus*.




Figure 30: Effect of *P. indica* on the growth of *L. japonicus* and *A. thaliana* grown in soil with different nutrient availabilities. Box-plots represent stem height of plants (ca. 20/treatment) at the indicated dpi. On high-nutrient soil (500 mg l⁻¹ N, 500 mg l⁻¹ P, 500 mg l⁻¹ K), inoculation with *P. indica* did not change plant stem height at 21 dpi ($p > 0.05$). On low-nutrient soil (50-100 mg l⁻¹ N, 50-100 mg l⁻¹ P, 100-150 mg l⁻¹ K), *P. indica* inoculation led to an increase in the mean stem height at 21 dpi ($p < 0.05$). Statistical analyses were performed with a Kruskal-Wallis test followed by a Bonferroni-Holm correction using the mock-inoculated samples as control group. For each mock/*P. indica*-inoculated pair, box-plots that share the same letter do not significantly differ (at the 5% significance level). White boxes: mock; grey boxes: *P. indica*; open circles: outliers. Exemplary pictures of *P. indica*- and mock-inoculated plants grown in low nutrient soil. Experiments were performed three times with similar results.

When *A. thaliana* plants were grown on agar plates with $\frac{1}{2}$ MS medium and no sugar, a growth promoting effect by inoculation with *P. indica* chlamydo spores was observed. In contrast, when 0.05% sucrose was added to the medium, the plants were generally bigger and no increase was observed in the mean fresh weight of *P. indica*-inoculated plants compared to mock-treated plants (Figure 31, Table 1).




Figure 31: Effect of sucrose on the growth promotion of *A. thaliana* plants by *P. indica*. Seedlings (Col-0) (ca. 15/treatment) were grown on $\frac{1}{2}$ MS with or without sucrose for 10 days before being inoculated with *P. indica* chlamydo spores (grey boxes) or mock-inoculated with Tween water (white boxes). Box plots represent the fresh weight of plants at 7 dpi. Without sucrose addition, the mean fresh weight of *P. indica*-inoculated plants was significantly higher than that of mock-treated plants ($p < 0.05$). In the presence of sucrose, the mean fresh weight of the plants grown in the presence of *P. indica* was not significantly different from that of mock-treated plants ($p > 0.05$). Statistical analyses were performed with a Kruskal-Wallis test followed by a Bonferroni-Holm correction using the mock-inoculated samples as control group. Plots sharing a letter do not significantly differ (significance level: 5%). Open circles: outliers. Results from two independent experiments are shown.

Since *A. thaliana* plants inoculated with *P. indica* exhibit increased uptake of nitrogen and phosphate (Shahollari *et al.*, 2007; Kumar *et al.*, 2011; Das *et al.*, 2014), we investigated whether the concentration of nitrate [supplied as $\text{Ca}(\text{NO}_3)_2$ and/or KNO_3], phosphate (KH_2PO_4), or ammonium (NH_4Cl) on modified HO medium influenced the growth promotion of *A. thaliana* plants by *P. indica*. Co-cultivation with the fungus had a positive effect on the mean fresh weight of the plants under all nutrient conditions tested, except on medium depleted of any nitrogen source, a condition that limited plant growth due to nitrogen starvation (Figure 32, Table 1).




Figure 32: Influence of $\text{Ca}(\text{NO}_3)_2$, KNO_3 , KH_2PO_4 or NH_4Cl concentrations on the growth promotion of *A. thaliana* by *P. indica*. Bars represent the fresh weight of *P. indica*-inoculated plants (ca. 10 plants/treatment) relative to the respective mock control at 7 dpi, and error bars refer to the standard deviation of the mean of the fresh weight values. Overall, co-cultivation with *P. indica* chlamydo spores had a positive effect on the mean fresh weight of the plants, independent of nutrient concentration, except for medium depleted of any nitrogen source, on which the presence of the fungus did not significantly change plant biomass ($p > 0.05$). Plants were grown on the modified HO medium with the indicated salt concentrations. White box: mock; grey box: *P. indica*. Statistical analyses were performed with a Kruskal-Wallis test followed by a Bonferroni-Holm correction using the mock-inoculated samples as control group, and P was calculated on the absolute values. Each pair mock/*P. indica*-inoculated was compared individually. Stars indicate a significant difference (at the 5% significance level: 5%) relative to the non-inoculated control. *, $P < 0.05$; ***, $P < 0.001$.

Experiment was designed and performed by Aline Banhara Pereira. Statistical analyses were done by Aline Banhara Pereira and Alga Zuccaro, and Aline Banhara Pereira prepared the figure.

5.9. *A. thaliana* HCSGs are not required for *P. indica*-induced growth promotion

We investigated the influence of *A. thaliana* HCSGs on the host growth-promoting effect by *P. indica*. For that, we inoculated wild-type *A. thaliana* plants, the HCSG mutants *pollux*, *sec13*, *nup133*, and the double mutant *sec13* x *nup133* with chlamydospores of *P. indica*. As a specificity control, we included the closely related sebacinoid fungus *P. williamsii* (Basiewicz *et al.*, 2012; Lahrman *et al.*, 2013). Plant biomass was evaluated 7 dpi. Plants inoculated with *P. indica* had a significant higher mean fresh weight than plants mock-inoculated with Tween water, whereas treatment with *P. williamsii* chlamydospores resulted in no or only a slight increase (for *sec13* x *nup133*) in plant biomass (Figure 33, Table 2).




Figure 33: Biomass of *A. thaliana* wild-type or HCSG mutants upon inoculation with *P. indica* or *P. williamsii*. Upper panel: box-plots show the fresh weight of the indicated *A. thaliana* genotypes (ca. 15 plants/genotype) grown in the presence or absence of *P. indica* or *P. williamsii*. All plant genotypes accumulated more biomass upon co-cultivation with *P. indica* but not with *P. williamsii*. White box: mock; light grey box: *P. indica*; dark grey box: *P. williamsii*. Open circles: outliers. Statistical analyses were performed using a Kruskal-Wallis test followed by a Bonferroni-Holm correction using the mock-inoculated samples as control group. Boxes that do not share a letter are significantly different (at the 5% significance level). Comparisons among the three treatments were made for each genotype separately. Lower panel: representative plates showing sets of 14-day-old plants grown on modified HO medium are shown seven days after mock-treatment with Tween water (left) or inoculation with either *P. indica* (center) or *P. williamsii* (right) chlamydospores. 1: *pollux*; 2: *sec13 x nup133*; 3: *nup133*; 4: *sec13*; 5: Col-0.

Table 1: Statistical values for pairwise comparisons performed with the Kruskal-Wallis test followed by a Bonferroni-Holm correction using the mock-inoculated samples as control group.

Growth in soil

A. thaliana

Condition	treatment	<i>P</i>
Low nutrient soil	mock x <i>P. indica</i>	0.01
	mock x <i>P. indica</i>	0.037
	mock x <i>P. indica</i>	0.175
	mock x <i>P. indica</i>	0.298
High nutrient soil	mock x <i>P. indica</i>	<0.001
	mock x <i>P. indica</i>	0.0
	mock x <i>P. indica</i>	0.0
	mock x <i>P. indica</i>	0.0

L. japonicus

Condition	treatment	dpi	<i>P</i>
High nutrient soil	<i>P. indica</i>	7	0.110
	<i>P. indica</i>	11	0.167
	<i>P. indica</i>	16	0.270
	<i>P. indica</i>	21	0.930
Low nutrient soil	<i>P. indica</i>	7	0.160
	<i>P. indica</i>	11	0.390
	<i>P. indica</i>	16	0.046
	<i>P. indica</i>	21	0.013

Growth on modified HO

Condition	treatment	<i>P</i>
0 mM KNO ₃	<i>P. indica</i>	0.770
0.5 mM KNO ₃	<i>P. indica</i>	<0.001
5 mM KNO ₃	<i>P. indica</i>	<0.001
10 mM NH ₄ Cl	<i>P. indica</i>	0.025
20 mM NH ₄ Cl	<i>P. indica</i>	<0.001
0 mM KH ₂ PO ₄	<i>P. indica</i>	<0.001
Modified HO (standard)	<i>P. indica</i>	<0.001

Growth on MS

Condition	treatment	<i>P</i>	Fig. 31 upper	Fig. 31 lower
MS0	<i>P. indica</i>	0.0		0.0013
MS (+) sucrose	<i>P. indica</i>	0.3866		0.9211

P: probability at the 5% significance level

P < 0.05: significant biomass increase relative to mock

Table 2: Statistical values for multiple comparisons performed the data obtained with the inoculation of *A. thaliana* HSCS mutants with either *P. indica* or *P. williamsii* with the Kruskal-Wallis test followed by a Bonferroni-Holm correction using the mock-inoculated samples as control group.

Treatment	genotype	Effect	<i>P</i>
<i>P. indica</i>	Col(-)0	(+)	<0.001
<i>P. williamsii</i>	Col(-)0	none	0.110
<i>P. indica</i>	<i>pollux</i>	(+)	0.0086
<i>P. williamsii</i>	<i>pollux</i>	none	1.0
<i>P. indica</i>	<i>sec13</i>	(+)	<0.001
<i>P. williamsii</i>	<i>sec13</i>	none	0.35
<i>P. indica</i>	<i>nup133</i>	(+)	<0.001
<i>P. williamsii</i>	<i>nup133</i>	none	1.0
<i>P. indica</i>	<i>sec13</i> x <i>nup133</i>	(+)	<0.001
<i>P. williamsii</i>	<i>sec13</i> x <i>nup133</i>	(+)	0.042

(+): biomass increase

6. Discussion

6.1 *A. thaliana* mutants for HCSGs impair haustorial development and the reproductive success of a biotrophic oomycete

In this work, we demonstrated that the *A. thaliana* HCSGs *POLLUX*, *ShRK1*, *ShRK2*, *NUP133*, and *SEC13* are required for the full reproductive success of *Hpa* during colonization of *A. thaliana* leaves. This reduced susceptibility was related neither to reduced cell penetration frequency nor to constitutive or exacerbated activation of pathogen-associated molecular pattern (PAMP)-triggered plant immunity (Jones & Dangl, 2006) in the HCSG mutants. Instead, it was probably caused by an alteration in the haustorial development. The HCSG mutants presented a higher frequency of morphologically altered haustoria, and the change in morphology was associated with a loss of fluorescence generated by a perahaustorial marker protein, suggesting a modification in the protein composition of the perahaustorial membrane. We showed that, in the double mutant *shrk1* x *shrk2*, the haustoria seem to age earlier than in the wild-type, indicating a direct effect of the plant receptor kinases in the haustorial architecture of the oomycete. The haustorium was described to be the site of nutrient exchange with the plant host (Mendgen & Hahn, 2002; Voegelé & Mendgen, 2003). Thus, our results suggest that the impaired haustorial development possibly results in a decreased nutrient availability to the oomycete, with a consequent reduction in the sporangiophore production.

Interestingly, the colonization by the extracellular bacterial pathogen *P. syringe* was normal in the *A. thaliana* HCSG mutants, indicating that the phenotype of altered haustorial development in all tested HCSG mutants is specific to biotrophic interactions that involve the formation of typical intracellular accommodation organs. To further test this hypothesis, we analysed the reproductive success of the powdery mildew fungus *E. cruciferarum* on *A. thaliana*, without revealing any consistent difference to the colonization pattern in wild-type leaves. Due to the highly variable morphology of the haustoria formed during colonization of wild-type leaves, we were unable to determine if the haustoria developed normally in the HCSG mutants. *E. cruciferarum* only penetrates epidermal cells (Micali *et al.*, 2008; Huesmann *et al.*,

2011; Weis *et al.*, 2014), while *Hpa* initially forms haustoria in two anticlinal epidermal cells but then proceeds to penetrate the mesophyll cells (Koch & Slusarenko, 1990; Slusarenko & Schlaich, 2003). Assuming that HCSGs do not play a role in this interaction, one could therefore propose that distinct pathways are used for colonization of the leaf epidermis and subjacent mesophyll cells. The *MLO* gene, for example, is a compatibility factor expressed in the epidermis and required for penetration of the powdery mildew fungus *G. orontii* into *A. thaliana* epidermal cells (Consonni *et al.*, 2006), with no reported role in the *Hpa* colonization. Alternatively, different genetic components may be required by intracellular oomycetal and fungal pathogens, with the pathway comprising the HCSGs being specifically recruited by *Hpa*. In this context, it would be interesting to investigate whether these genes are required for the interaction between *A. thaliana* and other biotrophic oomycetes, such as *Phytophthora parasitica*, member of a genus known to cause devastating diseases on economically important crops. *P. infestans*, for instance, is the causal agent of the late blight on potato and responsible for the Great Irish Famine in the 1840s (Fry, 2008; Haas *et al.*, 2009). Apart from penetrating *A. thaliana* leaves, *P. parasitica* is also able to enter root cells (Attard *et al.*, 2010; Wang *et al.*, 2011). This is particularly interesting, since the root is the organ where the CSG of legumes exert their role during AM symbiosis.

6.2 Symbiosis-related genes are not required for the colonization of *L. japonicus* and *A. thaliana* with the root endophytic fungus *P. indica*

The similarities between colonization of the outer cell layers of plant roots by AM fungi and *P. indica* led us to investigate whether symbiosis-related genes were required for *P. indica* colonization of *L. japonicus* and *A. thaliana* roots. In both wild-type plants and in the common symbiosis mutants, *P. indica* was able to successfully complete its life within root cells, where colonization went through an initial biotrophic phase, similar to what has been demonstrated for other species such as barley and *A. thaliana* (Jacobs *et al.*, 2011; Zuccaro *et al.*, 2011; Lahrman & Zuccaro, 2012; Lahrman *et al.*, 2013), and culminated with intracellular sporulation, indicating that the CSGs are not required for the intracellular colonization of *L. japonicus* roots by *P. indica*. This was further corroborated by the observation that the

inoculation of the T90 symbiosis reporter line with *P. indica* did not activate GUS expression, for which the common symbiosis genes are required (Kistner *et al.*, 2005; Gossmann *et al.*, 2012).

A core set of CSGs, including *CCaMK* and *CYCLOPS*, is absent from the *A. thaliana* genome (Delaux *et al.*, 2014). The observations that *P. indica* was able to establish a biotrophic colonization and sporulate in the roots of *A. thaliana* HCGS mutants, and the relative amount of fungal hyphae in these mutants was not altered, suggest that the HCSGs are not required for the intracellular colonization of *A. thaliana* by *P. indica*.

Together, our results imply the existence of compatibility factors in the *P. indica*-plant root that function independently of symbiosis-related genes.

6.3. Increased fungal biomass in *L. japonicus* common symbiosis mutants suggests cross-talk between symbiosis signalling and defence or developmental pathways

Interestingly, we detected a tendency for increased amount of intraradical hyphae in some of the common symbiosis mutants in comparison to the wild-type, with a significant difference for *nup85-1*, *ccamk-12*, and *cyclops-3*. This observation does not imply roles for the CSGs in the direct accommodation of *P. indica* within root cells as seen in the AM symbiosis. Instead, it suggests a negative role for these genes on fungal proliferation in the host roots, and might be related to possible overlaps between signalling pathways involved in symbiotic and pathogenic interactions, as already reported for a number of genes (reviewed in (Evangelisti *et al.*, 2014)). However, such effect was detected on *L. japonicus* but not on *A. thaliana* mutants, implying that the regulatory pathways affected in legumes may not be functional in *A. thaliana*.

In *A. thaliana* and in barley, the establishment or maintenance of the association between *P. indica* and plant roots requires that host defence responses are at some extent suppressed (Schäfer *et al.*, 2009; Jacobs *et al.*, 2011). Therefore, a situation in which the manipulation of the host immune system is not properly regulated could result in an increased susceptibility to the fungus, causing a more dense colonization. Hyper-infection by *P. indica* has been associated with deregulated defence responses at least in *A. thaliana* (Camehl *et al.*, 2010). In legumes, common symbiosis genes

have been connected to plant defence signaling as well. The Calcium and Calmodulin dependent protein Kinase gene *DMI3*, the *CCaMK* ortholog in *M. truncatula*, plays a role in the interaction with the hemi-biotrophic pathogenic fungus *Colletotrichum trifolii*; in the *dmi3* mutant, the switch from biotrophy to necrotrophy, which starts upon fungal penetration, occurred earlier (Genre *et al.*, 2009). In barley, a dense colonization by *P. indica* was shown in dead cells of the root differentiation and maturation zones (Deshmukh *et al.*, 2006). It will be therefore important to determine whether increased relative *P. indica* biomass in the roots of some of the *L. japonicus* common symbiosis mutants is associated with the necrotrophic colonization stage. This would explain the increased amount of intracellular hyphae that we detected for *ccamk-12* or perhaps through the same mechanism for the other common symbiosis mutants *nup85-1* and *cyclops-3*.

In addition, the bacterial PAMP (pathogen-associated molecular pattern) flg22 has been shown to play a negative role on rhizobial infection and on the *CCaMK*-induced nodule organogenesis during root nodule symbiosis between *L. japonicus* and *M. loti* (Lopez-Gomez *et al.*, 2011). In the same study, the nodule inception (NIN) transcription factor, which is a negative regulator of root cell competence to infection by rhizobia (Schauser *et al.*, 1999), was expressed after treatment with flg22. In *A. thaliana*, flg-22 mediated defence responses are suppressed by *P. indica* during the biotrophic phase (Jacobs *et al.*, 2011), implying a general negative effect of flg22 on infection by biotrophic microorganisms, and leaving us to speculate that the tendency for increased *P. indica* colonization in the roots of *L. japonicus* common symbiosis mutants results from alterations in the host immune system and the consequent inability to respond to invading microorganisms.

6.4. Growth promotion of *A. thaliana* by *P. indica* is independent of HCSGs but is affected by nutrient availability

In regard with *P. indica*-mediated biomass increase, considering that this trait is not robust and has not always been reproducible, we explored a range of distinct substrate and nutrient conditions and were able to confirm that the growth-promoting effect of *P. indica* (Peškan-Berghöfer *et al.*, 2004; Shahollari *et al.*, 2007; Sherameti *et al.*,

2008; Camehl *et al.*, 2010; Camehl *et al.*, 2011; Nongbri *et al.*, 2012; Lahrmann *et al.*, 2013; Venus & Oelmüller, 2013) on *A. thaliana* plants was present in all situations.

However, the mechanism by which the fungus promotes host growth is still highly debated. The importance of phytohormones for the establishment of a proper association between *P. indica* and plant roots has been shown in a number of studies. On the one hand, jasmonic acid and gibberellins positively influence root colonization, while salicylic acid has inhibitory effects (Jacobs *et al.*, 2011). On the other hand, ethylene was described as being required for colonization by *P. indica* (Khatabi *et al.*, 2012), and genes involved in the synthesis of indole-3-acetaldoxime-derived compounds restricts and prevents excessive growth of *P. indica* inside *A. thaliana* roots (Nongbri *et al.*, 2012). Importantly, the fungus has been shown to produce the phytohormone auxin (Sirrenberg *et al.*, 2007; Lee *et al.*, 2011), which has been implicated in stimulation of plant growth (Davies, 2004). Nevertheless, it has been claimed this auxin does not directly affect host biomass increase in barley (Hilbert *et al.*, 2012) or *A. thaliana* (Vadassery *et al.*, 2008), and the mechanism by which *P. indica* promotes host growth remains therefore still to be elucidated.

Our results in soil indicate that a nutrient effect at least contributes to *P. indica*-induced growth promotion, since this trait was much more pronounced when the substrate contained lower amounts of nitrogen, phosphate and potassium, suggesting that high nutrient availability could have negative effects on the establishment or maintenance of the *P. indica* symbiosis. Nevertheless, in order to obtain a more clear role of nutrient concentration on the interaction between *P. indica* and plant roots, it will be crucial to investigate whether the elevated nutrient availability relates to a decreased root colonization by *P. indica*, similar to what has been described for as AM symbiosis (Olsson *et al.*, 2005; Breuillin *et al.*, 2010; Balzergue *et al.*, 2013; Bonneau *et al.*, 2013; Gosling *et al.*, 2013).

The influence of *P. indica* on increased uptake of nitrogen and phosphate by the plant host has already been reported (Kumar *et al.*, 2011; Das *et al.*, 2014). Therefore, we investigated whether the concentration of nitrogen and phosphate had any influence on the growth promotion effect of *P. indica* on plates. We observed that, regardless of salt concentration, the biomass of inoculated plants was always higher than that of non-inoculated individuals, except under nitrogen deficiency, which may have prevented the plants from properly growing even when *P. indica* was present. Though

nutrient concentration did not have an obvious influence on the *P. indica*-induced host biomass increase, the presence of one specific compound may have compensated for the absence of another, and do not exclude that an increased uptake of nitrate, ammonium and/or phosphate is involved in the incremented host growth, as indicated in (Sherameti *et al.*, 2005; Kumar *et al.*, 2011; Lahrmann *et al.*, 2013).

Das and colleagues reported that the total content of soluble sugar in rice plants inoculated with *P. indica* is higher than in non-inoculated plants, possibly due to an increased transport of carbohydrates to the fungus (Das *et al.*, 2014). In addition, it has been shown that, in *A. thaliana* and *Nicotiana tabacum* roots, *P. indica* inoculation stimulates the expression of starch degrading enzymes, resulting in higher availability of sugar to the plants and/or the fungus (Sherameti *et al.*, 2005). We observed that addition of sucrose to the culture medium abolished the growth-promoting effect by *P. indica* on *A. thaliana*, indicating that excess sugar may have caused the plants to grow to their maximal potential and masked any fungal effect on internal sucrose mobilization.

Even though our studies indicated no role of known symbioses-related genes in the association between plant roots and *P. indica* in regards with colonization, we explored their roles in the *P. indica*-induced host biomass increase, and showed that it was still present in the *A. thaliana* HCSG mutants *pollux*, *nup133*, and *sec13*. Thus, regardless of the mechanism involved in the growth promotion of *A. thaliana* by *P. indica*, it does not require the HCSG *POLLUX* and the nucleoporins genes *NUP133* and *SEC13*, confirming previous observations on *AtPOLLUX* (Shahollari *et al.*, 2007).

Altogether, our results support the idea that the interaction of plant roots with *P. indica* does not require the common symbiosis pathway in legumes or remnants of this signaling network in *A. thaliana*, and suggests that the fungus relies on alternative plant factors to establish its symbiosis with plant roots. However, it is still not clear whether a specific pre-determined genetic pathway is involved in the intracellular accommodation of *P. indica*, or if it is mainly a consequence of successful manipulation of plant host defence responses and phytohormone balance by the fungus (Evangelisti *et al.*, 2014).

While the presence of both *P. indica* and AM fungi results in increased host growth (Peškan-Berghöfer *et al.*, 2004; Javot *et al.*, 2007a; Shahollari *et al.*, 2007; Sherameti

et al., 2008; Smith & Read, 2008; Camehl *et al.*, 2010; Camehl *et al.*, 2011; Nongbri *et al.*, 2012; Lahrmann *et al.*, 2013; Venus & Oelmuller, 2013; Nouri *et al.*, 2014), only the first undergoes its entire life cycle as a biotroph, whereas in *P. indica* the biotrophic phase of root colonization is followed by a stage characterized by cell death and necrotrophy (Deshmukh *et al.*, 2006; Zuccaro *et al.*, 2011; Lahrmann & Zuccaro, 2012; Qiang *et al.*, 2012). There is evidence that excessive *P. indica* proliferation is associated with necrotrophy and/or with detrimental effects on the growth of the plant host (Deshmukh *et al.*, 2006; Camehl *et al.*, 2010; Nongbri *et al.*, 2012). Therefore, it will be interesting to explore whether the increased relative *P. indica* biomass in the roots of some *L. japonicus* common symbiosis mutants lead to host growth retardation, and whether this uncontrolled fungal proliferation occurs during the necrotrophic colonization stage. Such approach might help us to establish if the host growth-promoting effect by *P. indica* results from events that take place during the biotrophic phase of the interaction.

6.5. *A. thaliana* homologs of common symbiosis genes: compatibility factors for *Hpa* accommodation

Taken together, our results place the *A. thaliana* HCSGs in a pathway that is specifically recruited by the oomycete *Hpa* for its accommodation inside leaf epidermal and mesophyll cells. The phenotype of altered haustorial development was observed exclusively in the HCSG mutants infected with *Hpa*. The *A. thaliana pskr1* mutant, which presents disease resistance resulting from constitutive defence responses (Mosher *et al.*, 2013), showed normal haustoria formation. Considering that the HCSG mutants are less susceptible to the oomycete with no abnormal activation of the plant immune system, we propose that the HCSGs act as compatibility factors (opposed to disease resistance factors) in the association of *Hpa* with *A. thaliana*.

Even though many genes were described to play a role in disease resistance, only a few have been reported to contribute to pathogen colonization of the plant hosts. Compatibility factors are normally part of the plant machinery that is manipulated by the biotrophs for their own growth. In this context, the genes *DMR1* and *IOS1* are particularly interesting, since, like *ShRK1* and *ShRK2*, both encode kinase proteins that play a role in the colonization of *A. thaliana* by *Hpa* (Van Damme *et al.*, 2005;

van Damme *et al.*, 2009; Hok *et al.*, 2011). Moreover, *IOS1* is a member of the malectin-like domain leucine-rich repeat receptor-like kinase (MLD-LRR-RLK) family. *ShRK1* and *ShRK2*, which we reported to be the closest homologs of the CSG *SYMRK*, and *SYMRK* itself, belong to this family as well. The GDCP motif in the extracytoplasmatic portion of *SYMRK*, required for the establishment of the AM symbiosis in the root epidermis (Kosuta *et al.*, 2011; Antolin-Llovera *et al.*, 2014), is present in *IOS1*, *ShRK1* and *ShRK2*, suggesting a possible role for this region in the *A. thaliana* association with *Hpa*. To confirm this hypothesis, it would be necessary to evaluate the oomycete colonization in mutants for genes encoding other members of the MLD-LRR-RLK family that lack the GDCP motif. One such candidate is At5g48740, whose kinase domain is the closest related to that of *SYMRK*, despite the fact that the identity between their extracellular domains is relatively lower.

Our finding that symbiosis-related genes may act as compatibility factors during the interaction of plants with both symbiotic and pathogenic microorganisms is reminiscent of the dual involvement of *RAM2* in AM symbiosis as well as in root colonization by an oomycete. In *M. truncatula*, *RAM2*, which encodes a glycerol-3-phosphate acyl transferase required for the production of cutin monomers, is required for the formation of hyphopodia, structures that attaches to the root surface at the fungal entry point preceding cell invasion during AM symbiosis (Gutjahr & Parniske, 2013), but also for the development of the highly specialized infection cells known as appressoria during root colonization by the oomycetal pathogen *Phytophthora palmivora* (Wang *et al.*, 2012). In the *A. thaliana* HCSG mutants, however, the initial process of cell infection by *Hpa* appears unaffected. Rather, our work shows genetic commonalities in the formation of intracellular accommodation organs later in the development of symbiosis and disease.

6.6. “Common symbiosis pathway” in *A. thaliana*?

Based on the function of the legume CSGs, it is tempting to assume that the HCSGs are similarly involved in a signaling network that promotes oomycetal development. However, it is still not clear whether these genes are directly responsible for the intracellular accommodation of *Hpa* in *A. thaliana*. It is therefore important to investigate whether there are mechanisms commonly used by microorganisms during symbiotic and pathogenic interactions and which of them are controlled by the

HCSGs. In this context, even if the remnants of this ancient pathway play roles similar to those of their legume counterparts, the signaling pathway leading to the successful establishment of the pathogenic interaction would still have gaps, meaning that the branch that involves calcium signaling and decoding in the plant nucleus still needs to be identified. Even though we have not yet proven that nuclear calcium signatures are also required for *Hpa* infection, some evidences support this hypothesis. In legumes and most angiosperms, *POLLUX* seems to play a role exclusively in the generation of calcium spiking during the biotrophic association with AM fungi (Charpentier *et al.*, 2008; Venkateshwaran *et al.*, 2012), agreeing with our findings in *A. thaliana*, where a *pollux* mutant had a phenotype only during the biotrophic interaction with *Hpa*. Besides, the facts that *POLLUX* is conserved among angiosperms (Kistner & Parniske, 2002), and potentially functional in the root nodule symbiosis in *M. truncatula* (Venkateshwaran *et al.*, 2012), support the idea that *POLLUX* and the calcium spiking machinery is conserved in *A. thaliana* as well.

Assuming that nuclear calcium signatures are part of the plant program for the accommodation of *Hpa*, candidates with similar functions to *CCaMK* and *CYCLOPS* in signal decoding and transcriptional activation in the plant nucleus during AM symbiosis (Singh & Parniske, 2012; Singh *et al.*, 2014) must be identified in *A. thaliana*. This would allow us to fill in the gaps in the putative signaling pathway that leads to the establishment of the biotrophic association with an oomycetal pathogen and provide further evidence that, at least in part, an ancient pathway is conserved for the accommodation of biotrophic hyphal microorganisms in symbiosis and disease.

6.7. Evolutionary forces driving the retention of the HSCGs in *A. thaliana*

The finding that vestiges of an ancient pathway for the intracellular accommodation of beneficial microorganisms was retained in a plant that is not able to establish root symbiosis with AM fungi, and the symbiosis with *P. indica* was proven to be independent of HCSGs, suggests that pathogens may take advantage of symbiotic programs. The lack of obvious developmental phenotypes in any of the HCSG mutants indicates that these genes do not fulfil any vital housekeeping functions. This

raises the question why the HCSG genes were kept in the *A. thaliana* genome for no apparent reason other than supporting the lifestyle of an oomycetal pathogen.

Contrary to the fitness improvement for plants that are able to establish symbioses, there is no obvious selective advantage for plants that are infected by *Hpa* that would support the retention of the HCSGs in the *A. thaliana* genome. Considering that it is not likely that plants evolved genes solely to favour pathogen growth, it is important to investigate whether ecological conditions exist, under which plants colonized by the oomycete present selective advantages over non-infected plants.

Plants have evolved distinct signaling pathways to deal with different types of pathogenic agents. In general, they activate salicylic acid (SA)-mediated defence responses against biotrophic pathogens, and jasmonate (JA)-dependent defences in response to wounding associated with interaction with necrotrophic pathogens or with herbivore insects (Kessler & Baldwin, 2002; Glazebrook, 2005). Some necrotrophic microorganisms, such as the bacterial pathogen *P. syringae*, induce both the JA and the SA-mediated defence responses (Stout *et al.*, 1998; Preston *et al.*, 1999; Thaler & Bostock, 2004). In nature, plants are often simultaneously or subsequently faced with different sorts of attackers, and must therefore deal with such stresses by finely regulating and/or integrating the defence signaling pathways. Generally, the SA and the JA-mediated defence responses have antagonistic effects on each other, with the outcome being that, when one pathway is activated, the plants may be primed for an improved defence against organisms of the same type, but become more susceptible to pathogens that activate the other defence pathway (Spoel *et al.*, 2007; Koornneef & Pieterse, 2008).

Herbivore pathogens present themselves as some of the most damaging plagues to plant crops (Stout *et al.*, 2006). Therefore, the activation of one or another defence signaling pathway by pathogenic microorganisms and the effect they have on future attacks by herbivores must be taken into consideration. Reports have shown that phloem-sucking insect herbivores are associated with SA-mediated defence responses, whereas chewing Lepidopteran herbivores normally induce the JA-dependent responses (Moran & Thompson, 2001; Cooper *et al.*, 2004; Kaloshian & Walling, 2005; Musser *et al.*, 2005; Rodriguez-Saona *et al.*, 2005; Thompson & Goggin, 2006). As a result, infection by pathogens that induce either SA or JA-

mediated defences may render the plants more resistant to future attacks by aphids or chewing insects, respectively (Thaler *et al.*, 2010).

Besides the influence of the crosstalk between those defence signaling pathways and the potentially positive or detrimental effects it might have on the fitness of herbivores feeding on plants already infected by pathogens, insect performance can also be affected by certain compounds produced by plants as a general defence response, such as peroxidases (Anfoka & Buchenauer, 1997; Dowd & Lagrimini, 1997; Stout *et al.*, 1999), or directly by the presence of microorganisms colonizing the plant host. Despite reports that pathogen-infected plants may be preferred by certain herbivores as food sources (Cardoza *et al.*, 2003; Mondy & Corio-Costet, 2004), existing infections could also render the affected hosts more resistant to herbivores. This increased resistance is assumed to be caused by the presence of detrimental chemical compounds, alterations in the host metabolism or because pathogen structures themselves act as physical barriers (Stout *et al.*, 2006). Herbivores either avoid colonized leaves or show decreased fitness by feeding on infected, nutrient-depleted leaves. The avoidance behaviour was observed when adult *Phaedon cochleariae* beetles chose healthy chinese cabbage (*Brassica rapa*) leaves over leaves colonized by the fungus *Alternaria brassicae* for oviposition. In the same system, larvae that fed on colonized leaves transformed into pupa with reduced weight (Rostás & Hilker, 2002). When the two species of alpine beetles, *Oreina elongata* and *O. cacaliae*, were offered healthy leaves and leaves of the plant host *Adenostyles alliariae* infected with the rust fungus *Uromyces cacaliae*, they chose the first, and, when offered only infected material, larvae of both species presented lower growth rates, lower maximum weights and prolonged developmental times (Röder *et al.*, 2007). Similar effects were observed with the tree-way-interaction between the plant host *Plantago lanceolata*, the powdery mildew fungus *Podosphaera plantaginis*, and larvae of the lepdoptera *Melitaea cinxia* (Laine, 2004).

The symptoms caused by *Hpa* infection of *A. thaliana* are usually only mild (Slusarenko & Schlaich, 2003). In fact, even increased seed production solely due to infection by *Hpa* has been described in low fecund *A. thaliana* ecotypes (Salvaudon *et al.*, 2008). Despite the lack of severe symptoms, *Hpa* colonization still results in the induction of defence responses associated with biotrophic pathogens, such as SA-dependent defences (Glazebrook, 2005; Caillaud *et al.*, 2013; Asai *et al.*, 2014),

raising the possibility that such induction primes the infected plants to defend themselves more efficiently against more aggressive microbial pathogens. Even though both scenarios would justify the retention of genes facilitating oomycetal accommodation in the *A. thaliana* genome, our future studies will concentrate on a third aspect: the potential “protective” effects of *Hpa* colonization against herbivores. Though the oomycete does not typically activate defence pathways associated with herbivory, it is still fathomable that nutrient depletion on the infected leaves, or reduced physical accessibility deters herbivores from feeding on *Hpa*-colonized plants. In order to test this hypothesis, we are currently studying the responses of the leaf-chewing caterpillar of the Lepidoptera *Spodoptera littoralis* to both healthy wild-type *A. thaliana* leaves and leaves fully colonized by *Hpa* and superficially covered with oomycetal sporangiophores.

7. Conclusion

By investigating the role of the *A. thaliana* HCSGs in the establishment of biotrophic interactions with beneficial and pathogenic hyphal microorganisms, we showed that these genes are required for haustoria development and full reproductive success of the pathogenic oomycete *Hpa*. However, we could not detect an influence on the colonization of *A. thaliana* leaves by the powdery mildew fungus *E. cruciferarum* or the bacterial pathogen *P. syringe*. Likewise, the colonization of *A. thaliana* by *P. indica* or the associated growth promotion occurred independently of the HCSGs, even though the fungus is beneficial and grows inside roots, as do symbiotic AM fungi. Altogether, our results imply that the HCSGs have not been generally opted during evolution for the intracellular accommodation of hyphal pathogens and symbionts, but are required for the interaction with *Hpa* in particular. However, the cellular mechanisms involved in such interaction remain to be elucidated. Interestingly, though in *L. japonicus* the CSGs are not required for the biotrophic colonization of roots by *P. indica*, it appears that these genes have a negative role on fungal proliferation, suggesting an interconnection between symbiosis and defence signalling or plant developmental programs.

8. Materials and Methods

8.3. Fungal strains and growth conditions

P. indica ((Savita *et al.*, 1998), DSM11827, from Deutsche Sammlung von Mikroorganismen und Zellkulturen, Braunschweig, Germany) and *P. williamsii* (Basiewicz *et al.*, 2012) were grown at 28°C, in the dark, on plates with solid (1.5% agar) complete medium (CM) (Pham *et al.*, 2004). For studies with plants grown in soil fungal mycelia were propagated in liquid CM medium, in the dark, at RT and 120 rpm shaking. Mycelium was washed three times with sterile distilled water directly prior to mixing with soil substrate.

For experiments on plates, fungal chlamydospores were obtained from four to six-week-old cultures, as follows: Tween water (0.002% Tween 20) was added to plates containing mycelium, which was rubbed off from the agar surface, and the resulting spore suspension was collected and centrifuged three times at 3.000 *g* for washing. Chlamydospore concentration was estimated using a Fuchs-Rosenthal counting chamber and adjusted to the desired $5 \times 10^5 \text{ ml}^{-1}$ with Tween water.

8.4. Plant genotypes and growth conditions

For co-cultivation of *L. japonicus* with *P. indica*, seeds (Gifu ecotype wild-type and common symbiosis mutants; Table 3) were surface sterilized in a 2% sodium hypochlorite solution for 7 min, washed three times with sterile water for 5 min, and kept in water overnight until the seeds were swollen. Seeds were then put on plates with solid modified Hoagland's medium (5 mM KNO₃; 5 M Ca(NO₃)₂; 2 mM MgSO₄; 4 mM KH₂PO₄; 0.03 g l⁻¹ Sprint 138 iron chelate; 0,1% micronutrients solution containing 2.86 g l⁻¹ H₃BO₃; 1.81 g l⁻¹ MnCl₂.4H₂O; 0.08 g l⁻¹ CuSO₄.5H₂O; 0.02 g l⁻¹ 85% MoO₃.H₂O; based on (Hoagland & Arnon, 1950)) and kept for 4 days at 24°C in the dark for germination.

Table 3: *L. japonicus* common symbiosis mutants screened for phenotypes associated with *P. indica* colonization.

Mutant	Reference
<i>castor-13</i>	(Imaizumi-Anraku <i>et al.</i> , 2005)
<i>ccamk-12</i>	(Perry <i>et al.</i> , 2009)
<i>cyclops-3</i>	(Schauser <i>et al.</i> , 1998; Szczyglowski <i>et al.</i> , 1998; Kistner <i>et al.</i> , 2005)
<i>nup85-1</i>	(Saito <i>et al.</i> , 2007)
<i>nup133-1</i>	(Schauser <i>et al.</i> , 1998; Kanamori <i>et al.</i> , 2006)
<i>pollux-5</i>	(Imaizumi-Anraku <i>et al.</i> , 2005)
<i>symrk-3</i>	(Stracke <i>et al.</i> , 2002; Demchenko <i>et al.</i> , 2004; Kistner <i>et al.</i> , 2005)

A. thaliana seeds (all ecotype Col-0) were obtained from "The Nottingham Arabidopsis Stock Centre" – NASC (55) or the GABI-DUPLO double mutant collection (Scholl *et al.*, 2000) (Table 4). For *in vitro* experiments, *A. thaliana* seeds were sterilized by incubation for 5 min in 70% ethanol/0.05% tween20, followed by 2 min in ethanol 100%. For *Hpa* infection, seeds were directly germinated in soil and grown for two weeks under long day conditions (16h light, 22°C $\mu\text{mol m}^{-2}\text{s}^{-1}$). For *Erysiphe cruciferarum* inoculation, *A. thaliana* plants were grown in a 2:1 soil/sand mixture. Seeds were stratified (4°C for 48h) prior to transfer into a growth chamber (10/14 light/dark cycle with 120 $\mu\text{mol m}^{-2}\text{s}^{-1}$ light, 22°C day, 20°C night, 65% relative humidity). For elicitor (flg22) treatment, seeds were placed on half-strength MS plates (57) stratified for 48h at 4°C in the dark, and grown under long day conditions (16h light, 23°C, 85 $\mu\text{mol m}^{-2}\text{s}^{-1}$) for 8 days.

Table 4: *A. thaliana* HCSG mutants used in the screening for phenotypes associated with *P. indica* symbiosis.

Gene	Gene ID	NASC ID
<i>POLLUX</i>	At5g49960	N566135
<i>SEC13</i>	At3g01340	N662322
<i>NUP133</i>	At2g05120	N565761

For co-cultivation of *A. thaliana* with *P. indica* or *P. williamsii* and tests for plant growth promotion, *A. thaliana* seeds were sterilized as described above, left to dry and stratified for 48h at 4°C in the dark. Plants were then grown under long day conditions (16 h : 8 h, light : dark, at 23°C, 85 $\mu\text{mol m}^{-2}\text{s}^{-1}$), on $\frac{1}{2}$ MS (Murashige & Skoog, 1962) with or without sucrose (0.05%) or on a modified Hoagland's (HO) medium solidified with gelrite 4 g l^{-1} , and four times more phosphate than in the original recipe. *L. japonicus* seeds (Table 3) were surface sterilized in a 2% sodium hypochlorite solution for 7 min, washed three times with sterile distilled water for 5 min, and kept in water overnight until the seeds were swollen. Seeds were then put on plates with solid modified HO and kept for 4 days at 24°C in the dark for germination.

Seven to ten-day-old *A. thaliana* or four-day-old *L. japonicus* plants were inoculated with 1 ml Tween water (0.002% Tween 20) or either *P. indica* or *P. williamsii* chlamydospores. Fungal chlamydospores were obtained from four to six-week-old cultures, as follows: Tween water was added to plates containing mycelium, which was rubbed off before the resulting spore suspension was collected and centrifuged three times at 3.000 g for washing. Chlamydospore concentration was estimated using a Fuchs-Rosenthal counting chamber and adjusted to the desired $5 \times 10^5 \text{ ml}^{-1}$ with Tween water. Non-germinated seeds and unhealthy seedlings were removed from the plates before inoculation. Co-cultures or mock-inoculated plants were grown at 24°C under long day conditions (16 h : 8 h, light : dark, 200 $\mu\text{mol m}^{-2}\text{s}^{-1}$), and inspected for biomass seven days post inoculation (dpi).

For analysis of growth promotion on different nutrient conditions, mock- and *P. indica* inoculated plants were grown on the following media: modified HO with no KNO_3 or $\text{Ca}(\text{NO}_3)_2$, supplemented with 0.05 mM KNO_3 , 0.5 mM KNO_3 , or 5 mM KNO_3 (standard concentration), modified HO depleted of phosphate or with 4 mM KH_2PO_4 (standard concentration); and modified HO depleted of ammonium or with 10 mM NH_4Cl or 20 mM NH_4Cl . When necessary, potassium and calcium concentrations were normalized with KCl and CaCl_2 , respectively.

For growth promotion experiments in soil, plants were grown on either low-nutrient soil (50-100 mg l^{-1} N, 50-100 mg l^{-1} P, 100-150 mg l^{-1} K) or high-nutrient soil (500 mg l^{-1} N, 500 mg l^{-1} P, 500 mg l^{-1} K) mixed with 1 g fresh or autoclaved mycelium per 100 g sterile substrate for inoculation and mock-inoculation, respectively. Plant height was determined 7, 11, 16, and 21 dpi.

8.5. *A. thaliana* stable transformation

Floral dipping was performed as described previously (Clough & Bent, 1998).

8.6. Pathogen assays and phenotypic analyses

Seven days after infection, *A. thaliana* leaves with sporulating *Hpa* isolate NoCo2 were harvested into 15 mL reaction tubes containing 10 mL dH₂O, and vortexed for 2s. The spore solution was then filtered through Miracloth filter and sprayed onto 12-day-old plants using a spraying gun. Subsequently, plants were placed into trays and covered with wet translucent lids. Trays were sealed to maintain high humidity, and plants were grown at 18°C under long day conditions (16h light, 85 $\mu\text{mol m}^{-2}\text{s}^{-1}$). Cotyledons (4 dpi) or leaves (4 or 5 dpi) were harvested and stained in 0.01 % trypan blue-lactophenol (10 mL lactic acid, 10 mL glycerol, 10 g phenol, 10 mg trypan blue dissolved in 10 mL distilled water) for 3 min at 95°C and 5h at room temperature, followed by overnight clearing in saturated chloral hydrate (2.5g/mL) and mounting in glycerol for observation using differential interference contrast microscopy. For sporangiophore counting, a minimum of 50 cotyledons per genotype and replicate were analysed and the number of sporangiophores per infected cotyledon was plotted. For investigation of the haustoria shape and the penetration efficiency, a minimum of five leaves per genotype and replicate were analysed. On each leaf, the percentage of multilobed haustoria per total haustoria or the percentage of haustoria-containing cells per cells contacted by hyphae was calculated for 10 individual strands of hyphae. The mean for each leaf was built and plotted.

Erysiphe cruciferarum was grown on Col-0 to maintain aggressiveness and on susceptible *phytoalexin deficient 4* (*pad4*) mutants (Glazebrook *et al.*, 1996) for elevated conidia production. *A. thaliana* plants were placed under a polyamide net (0.2 mm²) and inoculated at a density of 3–4 conidia mm⁻², by brushing conidia off of *pad4* plants through the net. Two leaves per plant were harvested, cleared and kept in acetic acid (25%) until analysis. Leaves were stained in acetic acid (25%) 1:9 + ink (Königsblau, Pelikan, 4001), washed in water, placed in water added of a few drops of Tween20, washed in water again, and analysed under a bright-field microscope.

Bacterial strains *P. syringae* pv. *tomato* DC3000 or *Pto* DC3000 Δ *AvrPto/AvrPto* were grown and used for infection assays on leaves of 4-5-week-old *A. thaliana* plants as described previously (Lin & Martin, 2005; Kemmerling *et al.*, 2007).

8.7. Observation of fluorescently labelled haustoria

For fluorescent analysis of haustoria morphology, leaves of *A. thaliana* wild-type and the *shrkl* x *shrk2* double mutant were harvested at 5 dpi, cleared in 10M KOH for 5 min, stained with 0.05% anilin blue in 0.067M K₂HPO₄ for 20 min and observed with a CLSM (Leica SP5) using excitation at 360-380 nm and detection at 470-505 nm. Images were edited using imageJ with the “volume viewer” plugin (Schneider *et al.*, 2012).

For quantification of fluorescence associated with the perihaustral membrane, wild-type *A. thaliana* (Col-0) plants transformed with the construct *ProAt3g61260:YFP-At3g61260* were infected with *Hpa* as described above, harvested at 8 dpi and observed with a microscope using differential interference contrast or epifluorescence with YFP filter settings (excitation at 500/20 nm and emission at 535/30 nm). The quantification was performed in triplicate. For each biological replicate consisting of 6 leaves, 100 single-lobed and 100 multilobed haustoria were counted and scored for detectable fluorescence. Haustoria without detectable fluorescence were visualized by differential interference contrast microscopy and scored as non-fluorescent. Leaves from *ProAt3g61260:YFP-At3g61260* plants were harvested at 8 dpi and observed with a CLSM (Leica SP5) using excitation at 510-520 nm and detection at 520-530 nm. 3D reconstructions of labelled haustoria were performed from a z-stack of 15 images (1,0-1,5 microns each) taken from fresh leaves at 5 or 6 dpi, using imageJ with the “3D viewer” plugin (Schneider *et al.*, 2012).

8.8. Analysis of oomycete-associated callose deposition

Oomycete-associated callose deposition was analysed on cotyledons of *A. thaliana* wild-type and HCSG mutants. Leaves were harvested at 4 dpi, cleared in 10M KOH for 5 min, stained with 0.05% anilin blue in 0.067M K₂HPO₄ for 20 min and mounted in glycerol for observation in an epifluorescence microscope (Leica DMI6000B) with CFP filter settings (excitation 436/10 nm and emission 465/30 nm). At least 50

pictures were taken per genotype. Regions of interest (ROIs) were selected in ImageJ (Schneider *et al.*, 2012), mean intensities were calculated from single ROIs and plotted.

8.9. Analyses of plant root colonization by *P. indica*

To observe *P. indica* intracellular sporulation in *A. thaliana* and *L. japonicus* roots, colonized roots at 14 dpi were incubated at 96°C for one (*A. thaliana*) or 10 min (*L. japonicus*) in 10% (w/v) KOH and double-stained in the dark at room temperature for 20 min with 10 µg ml⁻¹ WGA-AF488 (Molecular Probes, Karlsruhe, Germany) to visualize fungal structures, and 10 µg ml⁻¹ propidium iodide (PI) to visualize plant cell walls. Samples were analysed with a Leica DMI6000B microscope using differential interference contrast or epifluorescence (GFP filter set for WGA-AF488: excitation 450-490 nm, emission 500-550 nm; TX2 filter settings for PI: 540-580 nm excitation, 608-683 nm emission).

Root colonization and cell viability were analysed by confocal laser scanning microscopy (CLSM). Colonized roots at 7 dpi were stained for 10 min with 10 µg ml⁻¹ WGA-AF488 (Molecular Probes, Karlsruhe, Germany) to visualize fungal structures. Membranes were stained with 3 µM FM4-64 (Molecular Probes, Karlsruhe, Germany) for 5 min. Root samples were imaged with TCS-SP5 confocal microscopes (Leica, Bensheim, Germany) with excitation at 488 nm for WGA-AF488 and detection at 500–540 nm. FM4-64 was excited at 633 nm and detected at 650–690 nm. Propidium iodide was excited at 540 nm and detected at 600-630 nm.

8.10. Elicitor treatment

For pre-incubation, eight-day-old seedlings were transferred to a 12-well plate (3 seedlings/well represent one biological replicate) with half-strength liquid MS medium (Murashige & Skoog, 1962) supplemented with 1% sucrose and incubated overnight under long-day conditions (16 h light, 22 °C, 100 µmol m⁻² s⁻¹; 8 h dark, 18 °C) and 100 rpm shaking. On the following day the medium was exchanged, half the samples were supplemented with 1 µM flg22, and the other half kept in half-strength MS (Murashige & Skoog, 1962) as the mock controls. Plants were then incubated for

6h at 22°C and 100 rpm shaking. For every genotype, three biological replicates of treated and non-treated samples were harvested and immediately frozen in liquid N₂.

8.11. DNA/RNA extraction and qPCR analyses

For analyses of relative fungal/plant DNA content in roots of *A. thaliana* and *L. japonicus* colonized with *P. indica*, colonized roots (18 dpi) were thoroughly washed to remove fungal hyphae from the root surface. 200 mg of root material were then used for DNA extraction according to the protocol of Doyle and Doyle (Doyle & Doyle, 1987). Real-time qPCR analyses were performed from 10 ng DNA mixed with the appropriate primers in 10 µl SYBRgreen Supermix (BIORAD) using the following amplification protocol: 2'-95 °C; 40 x (30''-95°C; 30''-59°C; 30''-72 °C); melting curve 95°C – 60°C – 95°C. Relative DNA amount was calculated using the 2^{- $\Delta\Delta C_t$} method (Livak & Schmittgen, 2001).

For gene expression analyses before and after flg22 treatment, RNA extraction was performed using the SpectrumTM Plant Total RNA kit (Sigma-Aldrich), followed by DNaseI treatment (amplification grade DNaseI, InvitrogenTM) for removal of genomic DNA. First strand cDNA synthesis was performed from 250 ng total RNA using the SuperScript® III First-Strand Synthesis SuperMix (InvitrogenTM) with oligo(dT) primers. RT-qPCR was performed in 20 µL reactions containing 1x SYBR Green I (InvitrogenTM) in a CFX96 Real-time PCR detection system (Bio-Rad). PCR program: 2'-95°C; 40 x (30''-95°C; 30''-60°C; 20''-72°C); melting curve 95°C – 60°C – 95°C. A primer list can be found in the supplemental material. Expression levels of target genes were normalized against the housekeeping genes *TIP41-like* and *PP2A* (Czechowski *et al.*, 2005) was used as a reference to calculate the relative expression of the target genes. The *fls2* mutant was used as internal control. For every genotype, three biological replicates and two technical duplicates were analysed.

Table 5: Oligonucleotides used in the analyses of relative fungal DNA content in *P. indica* colonized roots.

Expression analysis		
Target	Sequence	Reference
<i>P. indica</i> TEF_fwd	TCGTCGCTGTCAACAAGATG	(Butehorn <i>et al.</i> , 2000)

<i>P. indica</i> TEF_rev	ACCGTCTTGGGGTTGTATC	
AtUBI_fwd	CCAAGCCGAAGAAGATCAAG	(Khatabi <i>et al.</i> ,
AtUBI_rev	ACTCCTTCCTCAAACGCTGA	2012)
LjUBI_fwd	ATGCAGATCTTCGTCAAGACCTTG	(Suzaki <i>et al.</i> ,
LjUBI_rev	ACCTCCCCTCAGACGAAG	2013)

Table 6: Oligonucleotides used to analyse gene expression patterns before and after flg22 treatment.

Expression analysis		
Target	Sequence	Reference
PP2A	TAACGTGGCCAAAATGATGC GTTCTCCACAAC^CGCTTGGT	(Czechowski <i>et al.</i> , 2005)
TIP41-like	GTGAAACTGTTGGAGAGAAGCAA TCAACTGGATACCCTTT^CGCA	(Czechowski <i>et al.</i> , 2005)
FRK1	ATCTTCGCTTGGAGCTTCTC TGCAGCGCAAGGACTAGAG	(He <i>et al.</i> , 2006)
GST1	TAATAAAAGTGGCGATGACC ACATTCAAATCAAACACTCG	(Grant <i>et al.</i> , 2000)
ERF1	TCGGCGATTCTCAATTTTTC ACAACCGGAGAACAACCATC-	(Denoux <i>et al.</i> , 2008)
PI-LTP	CTGGTTCATCCGAAACTGT CCTTCTGTTGCAAACGTTGA	(Denoux <i>et al.</i> , 2008)
PR1	TTCTTCCCTCGAAAGCTCAA AAGGCCACCAGAGTGTGTATG	(Schlicht & Kombrink, 2013)

8.12. β -glucuronidase (GUS) staining assays

GUS staining assays were performed with *L. japonicus* plants stably transformed with the symbiosis-reporter line T90 (Webb *et al.*, 2000). Colonized roots at 3 and 7 dpi with *P. indica* chlamydospores or Tween water were vacuum infiltrated with X-Gluc staining solution (100 mM Na₂HPO₄ pH 7.0; 10 mM EDTA; 0.1 % Triton-X 100; 0.5 mg ml⁻¹ X-Gluc; 1 mM K₃[Fe(CN)₆]; 1 mM K₄[Fe(CN)₆].3 H₂O) three times for 10 min, followed by incubation at 37°C for 18 h in the dark. Root systems were analysed for GUS staining with a Leica MZFLIII stereomicroscope.

8.13. Gene structure and phylogenetic analyses

Analyses of gene structures and protein domain organization were performed using online databases TAIR (<http://www.arabidopsis.org/>) for *A. thaliana*, and the KDRI website (Kazusa DNA Research Institute, Japan; <http://www.kazusa.or.jp/lotus/>) and the GenBank for *L. japonicus*. BLAST searches were performed on TAIR (<http://arabidopsis.org/Blast/index.jsp>) with the *L. japonicus* genomic CSG sequences as query.

For phylogenetic studies, protein sequences of *A. thaliana* and *L. japonicus* MLD-LRR-RLKs (either only the highly conserved kinase domain or only the extracytoplasmic region without signal peptide) were aligned using MAFFT 6.822 (Katoh *et al.*, 2002) with the default settings (alignments are provided as supplemental files). The result of the alignments were used to create phylogenetic trees at the CIPRES web-portal with RAxML 7.2.7 (Stamatakis *et al.*, 2008) for fast maximum likelihood analyses using 100 bootstraps. For RAxML, the JTT PAM matrix for amino acid substitutions was chosen, and the GTRGAMMA model was used for both bootstrapping and tree inference.

8.14. Statistics and data visualisation

All statistical analyses and data plots have been performed and generated with R version 3.0.2 (2013-09-25) "Frisbee Sailing" (Team, 2013).

For analyses with *Hpa* and *E. cruciferarum*, the packages "Hmisc" (Harrell, 2014) "car" (Fox, 2011), "multcompView" (Graves *et al.*, 2012), and "multcomp" (Hothorn *et al.*, 2008) were used. For statistical analysis, data was either subjected to the nonparametric Wilcoxon–Mann–Whitney test with Bonferroni-Holm correction comparing mutant and complementation groups with Col-0, or was power transformed to improve normality and a one-way ANOVA followed by a Tukey's HSD test or a Dunnett's Test with Bonferroni correction was performed using Col-0 samples as control group.

For analyses involving the interaction *A. thaliana*-*P. indica*, we used the package 'agricolae' (de Mendiburu, 2010). Pairwise or multiple comparisons of the different subsets of data were performed with the Kruskal-Wallis test followed by a

Bonferroni-Holm correction using the mock-inoculated samples as control group. Two-sided, unpaired t-tests with equal variance were used to analyse relative amount of fungal DNA within plant roots. A 95% family-wise confidence level was set for all experiments.

8.15. Cloning

Constructs labelled with “GG” were generated via Golden Gate cloning. For details on assembly method, general modules and plasmids (Gxx, BBxx), see (Binder *et al.*, 2014). Golden Gate constructs contain silent mutations to facilitate cloning. Constructs and oligonucleotides used for cloning are summarized on tables 7 and 8.

Table 7: Constructs used for cloning.

Entry clones / Golden Gate Level I & Level II plasmids (LI & LII)

Name	Description
pENTR-pSEC13:SEC13	Phusion PCR product consisting of SEC13 genomic construct amplified from <i>A. thaliana</i> gDNA with sec13co_FW and sec13co_RV, cloned into pENTR/D-TOPO (Invitrogen) via TOPO reaction
pENTR-pNUP133:NUP133 (GG)	NUP133 genomic construct with native promoter region (2158 bp). Introns 1 and 2 of <i>NUP133</i> were omitted for technical reasons. Final construct was assembled from 6 subcloned PCR fragments by Bsal cut ligation into pENTR-Bsal vector
	Fragment: Primers - template
	Fragment 1: N133-pro2_FW & N133-pro3_RV – <i>A. thaliana</i> gDNA
	Fragment 2: N133-pro3_FW & N133_Pro4_RV - <i>A. thaliana</i> gDNA
	Fragment 3: N133_ATG_FW & N133 mut_1b_RV - <i>A. thaliana</i> cDNA
	Fragment 4: N133_mut_1_FW & N133_mut_2_RV - <i>A. thaliana</i> gDNA
	Fragment 5: N133_mut2_FW & N133_e3_RV - <i>A. thaliana</i>

	gDNA
	Fragment 6: N133_e3_FW & N133_3'UTR2_RV- <i>A. thaliana</i> gDNA
LI A-C pPOLLUX (GG)	LI promoter element of POLLUX (1523 bp). Assembled from 2 PCR fragments amplified from <i>A. thaliana</i> gDNA by Bpil cut ligation into LI-Bpil vector Fragment 1: AtPol-Pro1+ & AtPol-Pro2- Fragment 2: AtPol-Pro3+ & AtPol-Pro4-
LI C-D POLLUX (GG)	LI element containing genomic <i>POLLUX</i> . Assembled from 2 PCR fragments amplified from <i>A. thaliana</i> gDNA by Bpil cut ligation into LI-Bpil vector. Fragment 1: AtPol1+ & AtPol2- Fragment 2: AtPol3+ & AtPol4-
LII F 1-2 pPOLLUX:POLLUX (GG)	Assembled by Bsal cut ligation from: LI A-C pPOLLUX + LI C-D POLLUX + LI dy D-E (BB8) + LI E-F 35S-T (G59) + LI F-G neo (G3) + LIIc F1-2 (BB30)
LII R 3-4 p35S-mCherry (GG)	Assembled by Bsal cut ligation from: LI A-B p35S (G5) + LI dy B-C (BB6) + LI C-D mCherry (G23) + LI dy D-E (BB8) + LI E-F nos-T (G6) + LI dy F-G (G9) + LIIc R 3-4 (BB33)
LII R 5-6 p35S-mCherry (GG)	Assembled by Bsal cut ligation from: LI A-B p35S (G5) + LI dy B-C (BB6) + LI C-D mCherry (G23) + LI dy D-E (BB8) + LI E-F HSP-T(G45) + LI dy F-G (G9) + LIIc R 5-6 (BB37)
LI A-C ShRK1 (GG)	Full length genomic DNA from ATG to codon prior to STOP (bases 1-4360); silent mutations introduced to remove type IIS restriction sites; obtained by gene synthesis (GenScript)
LI A-C ShRK2 (GG)	Full length genomic DNA from ATG to codon prior to STOP (bases 1-4185); silent mutations introduced to remove type IIS restriction sites; obtained by gene synthesis (GenScript)
LII F 1-2 pUBi: ShRK1-YFP (GG)	Assembled by Bsal cut ligation from: LI A-B pUBi (G7) + LI dy B-C (G4) + LI A-C ShRK1 + LI D-E YFP (G12) + LI E-F 35S-T (G59) + LI F-G hygro (G94) + LIIc F

	1-2 (BB30)
	Assembled by Bsal cut ligation from:
LII F 1-2 pUBi: ShRK2-YFP (GG)	LI A-B pUBi (G7) + LI dy B-C (G4) + LI A-C ShRK2 + LI D-E YFP (G12) + LI E-F 35S-T (G59) + LI F-G hygro (G94) + LIIc F 1-2 (BB30)
	Assembled by Bsal cut ligation from:
LII R 3-4 pUBi: ShRK1-CFP (GG)	LI A-B pUBi (G7) + LI dy B-C (G4) + LI A-C ShRK1 + LI D-E Cerulean (G14) + LI E-F HSP-T(G45) + LI dy F-G (G9) + LIIc R 3-4 (BB34)

Plasmids for stable transformation of *A. thaliana*

Name	Description
pSEC13:SEC13 + free mCherry	LR reaction of pENTR-pSEC13:SEC13 and PMDC99 (2); hygromycin resistance
pNUP133:NUP133	LR reaction of pENTR-pNUP133:NUP133 and pMDC99 (2); hygromycin resistance
	Assembled by Bpil cut ligation from:
pPOLLUX:POLLUX + free mCherry (GG)	LII F 1-2 pPOLLUX:POLLUX + LII 2-3 ins (BB43) + LII R 3-4 p35S-mCherry + LII dy 4-6 (BB41) + LIII α fin (BB45); kanamycin resistance
	Assembled by Bpil cut ligation from:
pUBi:ShRK1-YFP + free mCherry (GG)	LII F 1-2 pUBi:ShRK1 + LII 2-3 ins (BB43) + LII R 3-4 p35S-mCherry + LII dy 4-6 (BB41) + LIII α fin (BB45); hygromycin resistance
	Assembled by Bpil cut ligation from:
pUBi:ShRK2-YFP + free mCherry (GG)	LII F 1-2 pUBi:ShRK2 + LII 2-3 ins (BB43) + LII R 3-4 p35S-

mCherry +

LII dy 4-6 (BB41) + LIII α fin (BB45); hygromycin resistance**Table 8: Oligonucleotides used for cloning**

Plasmid construction

Name	Primer sequence (5' - 3')
sec13co_FW	CACCGGGAACACGGGAGAATAG
sec13co_RV	TTTTGCAATCTCTGTTGTCTGA
N133-pro2_FW	AGGGTCTCACACCGTTTTGAAAGACGGCATATTATGG
N133-pro3_RV	AGGGTCTCATACAAGGTCTTTTATTGCTTAAACTCT
N133-pro3_FW	AGGGTCTCATGTACATTTATTTGTTTTTCATTGATTG
N133_Pro4_RV	AGGGTCTCAACATTTTAAACCAGGAAGAGAGCGA
N133_ATG_FW	AGGGTCTCAATGTTCTCTCCATTGACGAAGA
N133_mut_1b_RV	AGGGTCTCATTTCTTTATCCATTCCACCGGA
N133_mut_1_FW	GGGGTCTCAGAAACCTGTCTTTCTTGTTTATT
N133_mut_2_RV	GGGGTCTCAGCGACCGAGAAGCCCT
N133_mut_2_FW	GGGGTCTCATCGCGTAGTCCTGTTGGTGT
N133_e3_RV	AGGGTCTCACTCTCTGCAGTTGAGTTCCTAGTG
N133_e3_FW	AGGGTCTCAAGAGCCTGCGAACTCTCAA
N133_3'UTR2_RV	AGGGTCTCACCTTGGTAGATTCGATACATCATAAAGAGG
AtPol-Pro1+	ATGAAGACTTTACGGGTCTCAGCGGAGCCCAATGACTTCCCAC AC
AtPol-Pro2-	TAGAAGACAAATGACTACAGTTTCATGCCACCA
AtPol-Pro3+	ATGAAGACTTTTCATCATTATGCTCATCTTGAATATGT
AtPol-Pro4-	TAGAAGACAACAGAGGTCTCAGGTGCGGGTTGAAGTAAGTAA TTGAGA
AtPol1+	ATGAAGACTTTACGGGTCTCACACCATGCCGATTCATACCCCT AGA
AtPol2-	TAGAAGACAACATCTTCTTTCTTCTGATTTGTTTCGT
AtPol3+	ATGAAGACTTGATGTTCTTTGAAGAAGAGACTAGC

AtPol4-

ATGAAGACTTCAGAGGTCTCACCTTCTGACTTGAGGCGATGAC
AAC

9. Declaration of contributions from other researchers

Aline Banhara Pereira, the author of this thesis, contributed to the experiments displayed in figures 1 to 33 as follows:

Figure 1: designed and prepared by Aline Banhara Pereira;

Figure 2 – 3, 23 – 25, 29 – 31, 33: Experiments were designed, performed and analysed by Aline Banhara Pereira, who prepared the corresponding figures;

Figure 4: Phylogenetic analyses were performed by Aline Banhara Pereira and Martina Katharina Ried. Aline Banhara Pereira prepared the figure;

Figure 5: Constructs *pollux* co, *shrkl* co, and *shrk2* co were produced by Andreas Binder. Experiments were designed, performed, and analysed by Aline Banhara Pereira and Martina Katharina Ried. Martina Katharina Ried did the statistical analyses and prepared the figure;

Figure 6: Experiments were designed, performed, and analysed by Aline Banhara Pereira and Martina Katharina Ried. Martina Katharina Ried did the statistical analyses and prepared the figure;

Figure 7: Experiment was designed and performed by Aline Banhara Pereira, and analysed by Aline Banhara Pereira and Martina Katharina Ried. Martina Katharina Ried did the statistical analyses and prepared the figure;

Figure 8: Construct *pollux* co was produced by Andreas Binder. Experiment was designed, performed, and analysed by Aline Banhara Pereira and Martina Katharina Ried. Martina Katharina Ried did the statistical analyses and prepared the figure;

Figure 9: Constructs *pollux* co, *shrkl* co, and *shrk2* co were produced by Andreas Binder. Experiments were designed, performed, and analysed by Aline Banhara Pereira and Martina Katharina Ried. Martina Katharina Ried did the statistical analyses and prepared the figure;

Figure 10: Experiment was designed, performed, and analysed by Aline Banhara Pereira and Martina Katharina Ried. Martina Katharina Ried did the statistical analyses and Aline Banhara Pereira prepared the figure;

Figure 11: Experiment was designed, performed, and analysed by Aline Banhara Pereira and Martina Katharina Ried. Aline Banhara Pereira prepared the figure;

Figure 12: Experiment was designed, performed, and analysed by Aline Banhara Pereira and Martina Katharina Ried. Aline Banhara Pereira prepared the figure;

Figure 13: Experiments was designed, performed, and analysed by Aline Banhara Pereira and Martina Katharina Ried. Martina Katharina Ried did the statistical analyses and prepared the figure;

Figure 14: Experiment was designed, performed, and analysed by Aline Banhara Pereira. Figure was prepared by Aline Banhara Pereira;

Figure 15: Experiment was designed and performed by Caroline Höfle, and analysed by Aline Banhara Pereira and Martina Katharina Ried. Martina Katharina Ried did the statistical analyses and prepared the figure;

Figure 16 – 18: Experiments were designed, performed and analysed by Aline Banhara Pereira and Martina Katharina Ried. Martina Katharina Ried did the statistical analyses and prepared the figure;

Figure 19 upper panel: Experiments were designed, performed, and analysed by Aline Banhara Pereira and Martina Katharina Ried. Statistical analyses were done by Martina Katharina Ried. Aline Banhara Pereira prepared upper panel and Martina Katharina Ried prepared lower panel;

Figure 20: Experiments were designed, performed and analysed by Aline Banhara Pereira. Aline Banhara Pereira prepared the figure;

Figure 21: Experiment was designed and performed by Aline Banhara Pereira, and analysed by Aline Banhara Pereira and Martina Katharina Ried. Statistical analyses were done by Martina Katharina Ried. Aline Banhara Pereira prepared the right panel, and Martina Katharina prepared the left panel;

Figure 22: Experiments were designed, performed, and analysed by Andrea A. Gust and Thorsten Nürnberger. Figure was prepared by Andrea A. Gust;

Figure 26: Experiments were designed by Alga Zuccaro and performed by Aline Banhara Pereira and Alga Zuccaro. Aline Banhara Pereira performed the statistical analyses and prepared the figure;

Figure 27: Experiment was designed and performed by Regina Kühner. Aline Banhara Pereira analysed the experiment and prepared the figure;

Figure 28: Experiments were designed, performed, and analysed by Aline Banhara Pereira, Yi Ding, and Alga Zuccaro. Aline Banhara Pereira prepared the figure;

Figure 32: Experiment was designed and performed by Aline Banhara Pereira. Statistical analyses were done by Aline Banhara Pereira and Alga Zuccaro, and Aline Banhara Pereira prepared the figure.

10. References

- Adam L, Ellwood S, Wilson I, Saenz G, Xiao S, Oliver RP, Turner JG, Somerville S. 1999.** Comparison of *Erysiphe cichoracearum* and *E. cruciferarum* and a survey of 360 *Arabidopsis thaliana* accessions for resistance to these two powdery mildew pathogens. *Mol Plant Microbe Interact* **12**(12): 1031-1043.
- Alber F, Dokudovskaya S, Veenhoff LM, Zhang W, Kipper J, Devos D, Suprpto A, Karni-Schmidt O, Williams R, Chait BT, et al. 2007.** The molecular architecture of the nuclear pore complex. *Nature* **450**(7170): 695-701.
- Aloui A, Recorbet G, Robert F, Schoefs B, Bertrand M, Henry C, Gianinazzi-Pearson V, Dumas-Gaudot E, Aschi-Smiti S. 2011.** Arbuscular mycorrhizal symbiosis elicits shoot proteome changes that are modified during cadmium stress alleviation in *Medicago truncatula*. *BMC Plant Biol* **11**: 75.
- Anfoka G, Buchenauer H. 1997.** Systemic acquired resistance in tomato against *Phytophthora infestans* by pre-inoculation with tobacco necrosis virus. *Physiol Mol Plant P* **50**(2): 85-101.
- Antolin-Llovera M, Ried MK, Parniske M. 2014.** Cleavage of the SYMBIOSIS RECEPTOR-LIKE KINASE ectodomain promotes complex formation with Nod factor receptor 5. *Curr Biol* **24**(4): 422-427.
- Aroca R, Porcel R, Ruiz-Lozano JM. 2007.** How does arbuscular mycorrhizal symbiosis regulate root hydraulic properties and plasma membrane aquaporins in *Phaseolus vulgaris* under drought, cold or salinity stresses? *New Phytol* **173**(4): 808-816.
- Asai S, Rallapalli G, Piquerez SJ, Caillaud MC, Furzer OJ, Ishaque N, Wirthmueller L, Fabro G, Shirasu K, Jones JD. 2014.** Expression profiling during *arabidopsis*/downy mildew interaction reveals a highly-expressed effector that attenuates responses to salicylic acid. *PLoS Pathog* **10**(10): e1004443.
- Asai T, Tena G, Plotnikova J, Willmann MR, Chiu WL, Gomez-Gomez L, Boller T, Ausubel FM, Sheen J. 2002.** MAP kinase signalling cascade in *Arabidopsis* innate immunity. *Nature* **415**(6875): 977-983.
- Attard A, Gourgues M, Callemeyn-Torre N, Keller H. 2010.** The immediate activation of defence responses in *Arabidopsis* roots is not sufficient to prevent *Phytophthora parasitica* infection. *New Phytol* **187**(2): 449-460.
- Bago B, Pfeiffer PE, Abubaker J, Jun J, Allen JW, Brouillette J, Douds DD, Lammers PJ, Shachar-Hill Y. 2003.** Carbon export from arbuscular mycorrhizal roots involves the translocation of carbohydrate as well as lipid. *Plant Physiol* **131**(3): 1496-1507.
- Bai Y, Pavan S, Zheng Z, Zappel NF, Reinstadler A, Lotti C, De Giovanni C, Ricciardi L, Lindhout P, Visser R, et al. 2008.** Naturally occurring broad-spectrum powdery mildew resistance in a Central American tomato accession is caused by loss of mlo function. *Mol Plant Microbe Interact* **21**(1): 30-39.
- Balergue C, Chabaud M, Barker DG, Becard G, Rochange SF. 2013.** High phosphate reduces host ability to develop arbuscular mycorrhizal symbiosis without affecting root calcium spiking responses to the fungus. *Front Plant Sci* **4**: 426.
- Bapaume L, Reinhardt D. 2012.** How membranes shape plant symbioses: signaling and transport in nodulation and arbuscular mycorrhiza. *Front Plant Sci* **3**: 223.
- Baptiste E, Charlebois RL, MacLeod D, Brochier C. 2005.** The two tempos of nuclear pore complex evolution: highly adapting proteins in an ancient frozen structure. *Genome Biol* **6**(10): R85.
- Basiewicz M, Weiss M, Kogel KH, Langen G, Zorn H, Zuccaro A. 2012.** Molecular and phenotypic characterization of *Sebacina vermifera* strains associated with

- orchids, and the description of *Piriformospora williamsii* sp. nov. *Fungal Biol* **116**(2): 204-213.
- Baxter L, Tripathy S, Ishaque N, Boot N, Cabral A, Kemen E, Thines M, Ah-Fong A, Anderson R, Badejoko W, et al. 2010.** Signatures of adaptation to obligate biotrophy in the *Hyaloperonospora arabidopsidis* genome. *Science* **330**(6010): 1549-1551.
- Belkhadir Y, Nimchuk Z, Hubert DA, Mackey D, Dangl JL. 2004.** *Arabidopsis* RIN4 negatively regulates disease resistance mediated by RPS2 and RPM1 downstream or independent of the NDR1 signal modulator and is not required for the virulence functions of bacterial type III effectors AvrRpt2 or AvrRpm1. *Plant Cell* **16**(10): 2822-2835.
- Bieleski RL. 1973.** Phosphate Pools, Phosphate Transport, and Phosphate Availability. *Ann Rev of Plant Physio* **24**(1): 225-252.
- Binder A, Lambert J, Morbitzer R, Popp C, Ott T, Lahaye T, Parniske M. 2014.** A modular plasmid assembly kit for multigene expression, gene silencing and silencing rescue in plants. *PLoS One* **9**(2): e88218.
- Binder A, Parniske M. 2013.** Analysis of the *Lotus japonicus* nuclear pore NUP107-160 subcomplex reveals pronounced structural plasticity and functional redundancy. *Front Plant Sci* **4**: 552.
- Bonfante P, Genre A. 2010.** Mechanisms underlying beneficial plant-fungus interactions in mycorrhizal symbiosis. *Nat Commun* **1**: 48.
- Bonneau L, Huguet S, Wipf D, Pauly N, Truong HN. 2013.** Combined phosphate and nitrogen limitation generates a nutrient stress transcriptome favorable for arbuscular mycorrhizal symbiosis in *Medicago truncatula*. *New Phytol* **199**(1): 188-202.
- Bowling SA, Clarke JD, Liu Y, Klessig DF, Dong X. 1997.** The *cpr5* mutant of *Arabidopsis* expresses both NPR1-dependent and NPR1-independent resistance. *Plant Cell* **9**(9): 1573-1584.
- Bowling SA, Guo A, Cao H, Gordon AS, Klessig DF, Dong X. 1994.** A mutation in *Arabidopsis* that leads to constitutive expression of systemic acquired resistance. *Plant Cell* **6**(12): 1845-1857.
- Bozkurt TO, Richardson A, Dagdas YF, Mongrand S, Kamoun S, Raffaele S. 2014.** The Plant Membrane-Associated REMORIN1.3 Accumulates in Discrete Perihyphal Domains and Enhances Susceptibility to *Phytophthora infestans*. *Plant Physiol* **165**(3): 1005-1018.
- Breullin F, Schramm J, Hajirezaei M, Ahkami A, Favre P, Druege U, Hause B, Bucher M, Kretschmar T, Bossolini E, et al. 2010.** Phosphate systemically inhibits development of arbuscular mycorrhiza in *Petunia hybrida* and represses genes involved in mycorrhizal functioning. *Plant J* **64**(6): 1002-1017.
- Burow M, Halkier BA, Kliebenstein DJ. 2010.** Regulatory networks of glucosinolates shape *Arabidopsis thaliana* fitness. *Curr Opin Plant Biol* **13**(3): 348-353.
- Butehorn B, Rhody D, Franken P. 2000.** Isolation and characterisation of *Pitef1* encoding the translation elongation factor EF-1 alpha of the root endophyte *Piriformospora indica*. *Plant Biol (Stuttg)* **2**(6): 687-692.
- Caillaud MC, Asai S, Rallapalli G, Piquerez S, Fabro G, Jones JD. 2013.** A downy mildew effector attenuates salicylic acid-triggered immunity in *Arabidopsis* by interacting with the host mediator complex. *PLoS Biol* **11**(12): e1001732.
- Callow JA, Capaccio LCM, Parish G, Tinker PB. 1978.** Detection and estimation of polyphosphate in vesicular-arbuscular mycorrhizas. *New Phytol* **80**(1): 125-134.
- Camehl I, Drzewiecki C, Vadassery J, Shahollari B, Sherameti I, Forzani C, Munnik T, Hirt H, Oelmüller R. 2011.** The OX11 kinase pathway mediates *Piriformospora indica*-induced growth promotion in *Arabidopsis*. *PLoS Pathog* **7**(5): e1002051.

- Camehl I, Sherameti I, Venus Y, Bethke G, Varma A, Lee J, Oelmuller R. 2010.** Ethylene signalling and ethylene-targeted transcription factors are required to balance beneficial and nonbeneficial traits in the symbiosis between the endophytic fungus *Piriformospora indica* and *Arabidopsis thaliana*. *New Phytol* **185**(4): 1062-1073.
- Capoen W, Sun J, Wysham D, Otegui MS, Venkateshwaran M, Hirsch S, Miwa H, Downie JA, Morris RJ, Ane JM, et al. 2011.** Nuclear membranes control symbiotic calcium signaling of legumes. *Proc Natl Acad Sci U S A* **108**(34): 14348-14353.
- Cardoza YJ, Lait CG, Schmelz EA, Huang J, Tumlinson JH. 2003.** Fungus-induced biochemical changes in peanut plants and their effect on development of beet armyworm, *Spodoptera exigua* Hübner (Lepidoptera: Noctuidae) larvae. *Environ Entomol* **32**(1): 220-228.
- Chandran D, Inada N, Hather G, Kleindt CK, Wildermuth MC. 2010.** Laser microdissection of *Arabidopsis* cells at the powdery mildew infection site reveals site-specific processes and regulators. *Proc Natl Acad Sci U S A* **107**(1): 460-465.
- Charpentier M, Bredemeier R, Wanner G, Takeda N, Schleiff E, Parniske M. 2008.** *Lotus japonicus* CASTOR and POLLUX are ion channels essential for perinuclear calcium spiking in legume root endosymbiosis. *Plant Cell* **20**(12): 3467-3479.
- Clough SJ, Bent AF. 1998.** Floral dip: a simplified method for *Agrobacterium*-mediated transformation of *Arabidopsis thaliana*. *Plant J* **16**(6): 735-743.
- Coates ME, Beynon JL. 2010.** *Hyaloperonospora arabidopsidis* as a pathogen model. *Annu Rev Phytopathol* **48**: 329-345.
- Consonni C, Humphry ME, Hartmann HA, Livaja M, Durner J, Westphal L, Vogel J, Lipka V, Kemmerling B, Schulze-Lefert P, et al. 2006.** Conserved requirement for a plant host cell protein in powdery mildew pathogenesis. *Nat Genet* **38**(6): 716-720.
- Cooper WC, Jia L, Goggin FL. 2004.** Acquired and R-gene-mediated resistance against the potato aphid in tomato. *Journal of Chemical Ecology* **30**(12): 2527-2542.
- Cruz C, Egsgaard H, Trujillo C, Ambus P, Requena N, Martins-Loucao MA, Jakobsen I. 2007.** Enzymatic evidence for the key role of arginine in nitrogen translocation by arbuscular mycorrhizal fungi. *Plant Physiol* **144**(2): 782-792.
- Czechowski T, Stitt M, Altmann T, Udvardi MK, Scheible WR. 2005.** Genome-wide identification and testing of superior reference genes for transcript normalization in *Arabidopsis*. *Plant Physiol* **139**(1): 5-17.
- Das J, Ramesh KV, Maithri U, Mutangana D, Suresh CK. 2014.** Response of aerobic rice to *Piriformospora indica*. *Indian J Exp Biol* **52**(3): 237-251.
- Davies PJ 2004.** The plant hormones: their nature, occurrence and function. In: Davies PJ ed. *Plant Hormones: Biosynthesis, Signal Transduction, Action!* Dordrecht, Netherland: Kluwer Academic Publisher, 1-15.
- de Jonge R, Bolton MD, Thomma BP. 2011.** How filamentous pathogens co-opt plants: the ins and outs of fungal effectors. *Curr Opin Plant Biol* **14**(4): 400-406.
- de Mendiburu F. 2010.** *Agricolae*: statistical procedures for agricultural research.
- Deising HB, Werner S, Wernitz M. 2000.** The role of fungal appressoria in plant infection. *Microbes Infect* **2**(13): 1631-1641.
- Delaux PM, Sejalon-Delmas N, Becard G, Ane JM. 2013.** Evolution of the plant-microbe symbiotic 'toolkit'. *Trends Plant Sci* **18**(6): 298-304.
- Delaux PM, Varala K, Edger PP, Coruzzi GM, Pires JC, Ane JM. 2014.** Comparative phylogenomics uncovers the impact of symbiotic associations on host genome evolution. *PLoS Genet* **10**(7): e1004487.
- Demchenko K, Winzer T, Stougaard J, Parniske M, Pawlowski K. 2004.** Distinct roles of *Lotus japonicus* SYMRK and SYM15 in root colonization and arbuscule formation. *New Phytologist* **163**(2): 381-392.

- Denoux C, Galletti R, Mammarella N, Gopalan S, Werck D, De Lorenzo G, Ferrari S, Ausubel FM, Dewdney J. 2008.** Activation of defence response pathways by OGs and Flg22 elicitors in *Arabidopsis* seedlings. *Mol Plant* **1**(3): 423-445.
- Deshmukh S, Huckelhoven R, Schafer P, Imani J, Sharma M, Weiss M, Waller F, Kogel KH. 2006.** The root endophytic fungus *Piriformospora indica* requires host cell death for proliferation during mutualistic symbiosis with barley. *Proc Natl Acad Sci U S A* **103**(49): 18450-18457.
- Deshmukh SD, Kogel KH. 2007.** *Piriformospora indica* protects barley from root rot caused by *Fusarium graminearum*. *J Plant Dis Protect* **114**(6): 263-268.
- Dou D, Zhou JM. 2012.** Phytopathogen effectors subverting host immunity: different foes, similar battleground. *Cell Host Microbe* **12**(4): 484-495.
- Dowd PF, Lagrimini LM. 1997.** Examination of different tobacco (*Nicotiana* spp.) types under- and overproducing tobacco anionic peroxidase for their leaf resistance to *Helicoverpa zea*. *Journal of Chemical Ecology* **23**(10): 2357-2370.
- Doyle JJ, Doyle JL. 1987.** A rapid DNA isolation procedure for small quantities of fresh leaf tissue. *Phytochem Bull* **19**: 11-15.
- Ehrhardt DW, Wais R, Long SR. 1996.** Calcium spiking in plant root hairs responding to Rhizobium nodulation signals. *Cell* **85**(5): 673-681.
- Evangelisti E, Rey T, Schornack S. 2014.** Cross-interference of plant development and plant-microbe interactions. *Curr Opin Plant Biol* **20C**: 118-126.
- Ezawa T, Hayatsu M, Saito M. 2005.** A new hypothesis on the strategy for acquisition of phosphorus in arbuscular mycorrhiza: up-regulation of secreted acid phosphatase gene in the host plant. *Mol Plant Microbe In* **18**(10): 1046-1053.
- Fabro G, Steinbrenner J, Coates M, Ishaque N, Baxter L, Studholme DJ, Korner E, Allen RL, Piquerez SJ, Rougon-Cardoso A, et al. 2011.** Multiple candidate effectors from the oomycete pathogen *Hyaloperonospora arabidopsidis* suppress host plant immunity. *PLoS Pathog* **7**(11): e1002348.
- Farmer EE, Almeras E, Krishnamurthy V. 2003.** Jasmonates and related oxylipins in plant responses to pathogenesis and herbivory. *Curr Opin Plant Biol* **6**(4): 372-378.
- Fiorilli V, Lanfranco L, Bonfante P. 2013.** The expression of GintPT, the phosphate transporter of *Rhizophagus irregularis*, depends on the symbiotic status and phosphate availability. *Planta* **237**(5): 1267-1277.
- Fox JW, S. 2011.** An {R} Companion to Applied Regression. Thousand Oaks {CA}: Sage.
- Freisleben R, Lein A. 1942.** Über die Auffindung einer mehlttauresistenten Mutante nach Röntgenbestrahlung einer anfälligen reinen Linie von Sommergerste. *Naturwissenschaften* **30**(40): 608-608.
- Fry W. 2008.** *Phytophthora infestans*: the plant (and R gene) destroyer. *Mol Plant Pathol* **9**(3): 385-402.
- Frye CA, Innes RW. 1998.** An *Arabidopsis* mutant with enhanced resistance to powdery mildew. *Plant Cell* **10**(6): 947-956.
- Genre A, Chabaud M, Faccio A, Barker DG, Bonfante P. 2008.** Prepenetration apparatus assembly precedes and predicts the colonization patterns of arbuscular mycorrhizal fungi within the root cortex of both *Medicago truncatula* and *Daucus carota*. *Plant Cell* **20**(5): 1407-1420.
- Genre A, Chabaud M, Timmers T, Bonfante P, Barker DG. 2005.** Arbuscular mycorrhizal fungi elicit a novel intracellular apparatus in *Medicago truncatula* root epidermal cells before infection. *Plant Cell* **17**(12): 3489-3499.
- Genre A, Ortu G, Bertoldo C, Martino E, Bonfante P. 2009.** Biotic and abiotic stimulation of root epidermal cells reveals common and specific responses to arbuscular mycorrhizal fungi. *Plant Physiol* **149**(3): 1424-1434.
- Ghabooli M, Khatabi B, Ahmadi FS, Sepehri M, Mirzaei M, Amirkhani A, Jorin-Novo JV, Salekdeh GH. 2013.** Proteomics study reveals the molecular mechanisms

- underlying water stress tolerance induced by *Piriformospora indica* in barley. *J Proteomics* **94**: 289-301.
- Gherbi H, Markmann K, Svistoonoff S, Estevan J, Autran D, Giczey G, Auguy F, Peret B, Laplaze L, Franche C, et al. 2008.** SymRK defines a common genetic basis for plant root endosymbioses with arbuscular mycorrhiza fungi, rhizobia, and Frankiabacteria. *Proc Natl Acad Sci U S A* **105**(12): 4928-4932.
- Glassop D, Godwin RM, Smith SE, Smith FW. 2007.** Rice phosphate transporters associated with phosphate uptake in rice roots colonised with arbuscular mycorrhizal fungi. *Can J Bot* **85**(7): 644-651.
- Glawischnig E. 2007.** Camalexin. *Phytochemistry* **68**(4): 401-406.
- Glazebrook J. 2005.** Contrasting mechanisms of defence against biotrophic and necrotrophic pathogens. *Annu Rev Phytopathol* **43**: 205-227.
- Glazebrook J, Rogers EE, Ausubel FM. 1996.** Isolation of *Arabidopsis* mutants with enhanced disease susceptibility by direct screening. *Genetics* **143**(2): 973-982.
- Gomez-Gomez L, Boller T. 2000.** FLS2: an LRR receptor-like kinase involved in the perception of the bacterial elicitor flagellin in *Arabidopsis*. *Mol Cell* **5**(6): 1003-1011.
- Gonzalez-Aguilera C, Askjaer P. 2012.** Dissecting the NUP107 complex: multiple components and even more functions. *Nucleus* **3**(4): 340-348.
- Gosling P, Mead A, Proctor M, Hammond JP, Bending GD. 2013.** Contrasting arbuscular mycorrhizal communities colonizing different host plants show a similar response to a soil phosphorus concentration gradient. *New Phytol* **198**(2): 546-556.
- Gossmann JA, Markmann K, Brachmann A, Rose LE, Parniske M. 2012.** Polymorphic infection and organogenesis patterns induced by a *Rhizobium leguminosarum* isolate from Lotus root nodules are determined by the host genotype. *New Phytol* **196**(2): 561-573.
- Govindarajulu M, Pfeffer PE, Jin H, Abubaker J, Douds DD, Allen JW, Bucking H, Lammers PJ, Shachar-Hill Y. 2005.** Nitrogen transfer in the arbuscular mycorrhizal symbiosis. *Nature* **435**(7043): 819-823.
- Grant JJ, Yun BW, Loake GJ. 2000.** Oxidative burst and cognate redox signalling reported by luciferase imaging: identification of a signal network that functions independently of ethylene, SA and Me-JA but is dependent on MAPKK activity. *Plant J* **24**(5): 569-582.
- Graves S, Piepho HP, Selzer L, with help from Dorai-Raj S 2012.** multcompView: Visualizations of paired comparisons.
- Groth M, Takeda N, Perry J, Uchida H, Draxl S, Brachmann A, Sato S, Tabata S, Kawaguchi M, Wang TL, et al. 2010.** *NENA*, a *Lotus japonicus* homolog of *Sec13*, is required for rhizodermal infection by arbuscular mycorrhiza fungi and rhizobia but dispensable for cortical endosymbiotic development. *Plant Cell* **22**(7): 2509-2526.
- Gutjahr C. 2014.** Phytohormone signaling in arbuscular mycorrhiza development. *Curr Opin Plant Biol* **20C**: 26-34.
- Gutjahr C, Banba M, Croset V, An K, Miyao A, An G, Hirochika H, Imaizumi-Anraku H, Paszkowski U. 2008.** Arbuscular mycorrhiza-specific signaling in rice transcends the common symbiosis signaling pathway. *Plant Cell* **20**(11): 2989-3005.
- Gutjahr C, Parniske M. 2013.** Cell and developmental biology of arbuscular mycorrhiza symbiosis. *Annu Rev Cell Dev Biol* **29**: 593-617.
- Haas BJ, Kamoun S, Zody MC, Jiang RH, Handsaker RE, Cano LM, Grabherr M, Kodira CD, Raffaele S, Torto-Alalibo T, et al. 2009.** Genome sequence and analysis of the Irish potato famine pathogen *Phytophthora infestans*. *Nature* **461**(7262): 393-398.

- Haga N, Kato K, Murase M, Araki S, Kubo M, Demura T, Suzuki K, Muller I, Voss U, Jurgens G, et al. 2007.** R1R2R3-Myb proteins positively regulate cytokinesis through activation of KNOLLE transcription in *Arabidopsis thaliana*. *Development* **134**(6): 1101-1110.
- Harrell FEJawcfCDamo 2014.** Hmisc: Harrell Miscellaneous.
- Harrison MJ. 1997.** The arbuscular mycorrhizal symbiosis: an underground association. *Trends Plant Sci* **2**(2): 54-60.
- Harrison MJ. 1999.** Molecular and cellular aspects of the arbuscular mycorrhizal symbiosis. *Annu Rev Plant Physiol Plant Mol Biol* **50**: 361-389.
- Harrison MJ. 2012.** Cellular programs for arbuscular mycorrhizal symbiosis. *Curr Opin Plant Biol* **15**(6): 691-698.
- Harrison MJ, Dewbre GR, Liu J. 2002.** A phosphate transporter from *Medicago truncatula* involved in the acquisition of phosphate released by arbuscular mycorrhizal fungi. *Plant Cell* **14**(10): 2413-2429.
- Harrison MJ, van Buuren ML. 1995.** A phosphate transporter from the mycorrhizal fungus *Glomus versiforme*. *Nature* **378**(6557): 626-629.
- Hawkins H-J, Johansen A, George E. 2000.** Uptake and transport of organic and inorganic nitrogen by arbuscular mycorrhizal fungi. *Plant Soil* **226**(2): 275-285.
- He P, Shan L, Lin NC, Martin GB, Kemmerling B, Nurnberger T, Sheen J. 2006.** Specific bacterial suppressors of MAMP signaling upstream of MAPKKK in *Arabidopsis* innate immunity. *Cell* **125**(3): 563-575.
- Heidel AJ, Dong X. 2006.** Fitness benefits of systemic acquired resistance during *Hyaloperonospora parasitica* infection in *Arabidopsis thaliana*. *Genetics* **173**(3): 1621-1628.
- Helber N, Wippel K, Sauer N, Schaarschmidt S, Hause B, Requena N. 2011.** A versatile monosaccharide transporter that operates in the arbuscular mycorrhizal fungus *Glomus* sp is crucial for the symbiotic relationship with plants. *Plant Cell* **23**(10): 3812-3823.
- Hilbert M, Voll LM, Ding Y, Hofmann J, Sharma M, Zuccaro A. 2012.** Indole derivative production by the root endophyte *Piriformospora indica* is not required for growth promotion but for biotrophic colonization of barley roots. *New Phytol* **196**(2): 520-534.
- Hoagland DR, Arnon DI. 1950.** *The water-culture method for growing plants without soil*. Berkeley, CA: University of California.
- Hoelz A, Debler EW, Blobel G. 2011.** The structure of the nuclear pore complex. *Annu Rev Biochem* **80**: 613-643.
- Hok S, Danchin EG, Allasia V, Panabieres F, Attard A, Keller H. 2011.** An *Arabidopsis* (malectin-like) leucine-rich repeat receptor-like kinase contributes to downy mildew disease. *Plant Cell Environ* **34**(11): 1944-1957.
- Hothorn T, Bretz F, Westfall P. 2008.** Simultaneous inference in general parametric models. *Biometrical Journal* **50**(3): 346-363.
- Huesmann C, Hoefle C, Huckelhoven R. 2011.** ROPGAPs of *Arabidopsis* limit susceptibility to powdery mildew. *Plant Signal Behav* **6**(11): 1691-1694.
- Huibers RP, de Jong M, Dekter RW, Van den Ackerveken G. 2009.** Disease-specific expression of host genes during downy mildew infection of *Arabidopsis*. *Mol Plant Microbe Interact* **22**(9): 1104-1115.
- Igarashi D, Tsuda K, Katagiri F. 2012.** The peptide growth factor, phyto-sulfokine, attenuates pattern-triggered immunity. *Plant J* **71**(2): 194-204.
- Imaizumi-Anraku H, Takeda N, Charpentier M, Perry J, Miwa H, Umehara Y, Kouchi H, Murakami Y, Mulder L, Vickers K, et al. 2005.** Plastid proteins crucial for symbiotic fungal and bacterial entry into plant roots. *Nature* **433**(7025): 527-531.
- Jacobs S, Zechmann B, Molitor A, Trujillo M, Petutschnig E, Lipka V, Kogel KH, Schafer P. 2011.** Broad-spectrum suppression of innate immunity is required

- for colonization of *Arabidopsis* roots by the fungus *Piriformospora indica*. *Plant Physiol* **156**(2): 726-740.
- Jakobsen I 1995.** Transport of phosphorus and carbon in VA mycorrhizas. In: Varma A, Hock B eds. *Mycorrhiza*: Springer Berlin Heidelberg, 297-324.
- Jarsch IK, Konrad SS, Stratil TF, Urbanus SL, Szymanski W, Braun P, Braun KH, Ott T. 2014.** Plasma membranes are subcompartmentalized into a plethora of coexisting and diverse microdomains in *Arabidopsis* and *Nicotiana benthamiana*. *Plant Cell* **26**(4): 1698-1711.
- Javot H, Penmetsa RV, Breuillin F, Bhattarai KK, Noar RD, Gomez SK, Zhang Q, Cook DR, Harrison MJ. 2011.** *Medicago truncatula* *mtpt4* mutants reveal a role for nitrogen in the regulation of arbuscule degeneration in arbuscular mycorrhizal symbiosis. *Plant J* **68**(6): 954-965.
- Javot H, Penmetsa RV, Terzaghi N, Cook DR, Harrison MJ. 2007a.** A *Medicago truncatula* phosphate transporter indispensable for the arbuscular mycorrhizal symbiosis. *Proc Natl Acad Sci U S A* **104**(5): 1720-1725.
- Javot H, Pumplin N, Harrison MJ. 2007b.** Phosphate in the arbuscular mycorrhizal symbiosis: transport properties and regulatory roles. *Plant Cell Environ* **30**(3): 310-322.
- Jin H, Pfeffer PE, Douds DD, Piotrowski E, Lammers PJ, Shachar-Hill Y. 2005.** The uptake, metabolism, transport and transfer of nitrogen in an arbuscular mycorrhizal symbiosis. *New Phytol* **168**(3): 687-696.
- Jogawat A, Saha S, Bakshi M, Dayaman V, Kumar M, Dua M, Varma A, Oelmuller R, Tuteja N, Johri AK. 2013.** *Piriformospora indica* rescues growth diminution of rice seedlings during high salt stress. *Plant Signal Behav* **8**(10): doi: 10.4161/psb.26891.
- Jones JD, Dangl JL. 2006.** The plant immune system. *Nature* **444**(7117): 323-329.
- Kaloshian I, Walling LL. 2005.** Hemipterans as plant pathogens. *Annu Rev Phytopathol* **43**: 491-521.
- Kanamori N, Madsen LH, Radutoiu S, Frantescu M, Quistgaard EM, Miwa H, Downie JA, James EK, Felle HH, Haaning LL, et al. 2006.** A nucleoporin is required for induction of Ca²⁺ spiking in legume nodule development and essential for rhizobial and fungal symbiosis. *Proc Natl Acad Sci U S A* **103**(2): 359-364.
- Kapulnik Y, Volpin H, Itzhaki H, Ganon D, Galili S, David R, Shaul O, Elad Y, Chet I, Okon Y. 1996.** Suppression of defence responses in mycorrhizal alfalfa and tobacco roots. *New Phytol* **133**(1): 59-64.
- Katoh K, Misawa K, Kuma K, Miyata T. 2002.** MAFFT: a novel method for rapid multiple sequence alignment based on fast Fourier transform. *Nucleic Acids Res* **30**(14): 3059-3066.
- Kemmerling B, Schwedt A, Rodriguez P, Mazzotta S, Frank M, Qamar SA, Mengiste T, Betsuyaku S, Parker JE, Mussig C, et al. 2007.** The BRI1-associated kinase 1, BAK1, has a brassinolide-independent role in plant cell-death control. *Curr Biol* **17**(13): 1116-1122.
- Kessler A, Baldwin IT. 2002.** Plant responses to insect herbivory: the emerging molecular analysis. *Annu Rev Plant Biol* **53**: 299-328.
- Kevei Z, Seres A, Kereszt A, Kalo P, Kiss P, Toth G, Endre G, Kiss GB. 2005.** Significant microsynteny with new evolutionary highlights is detected between *Arabidopsis* and legume model plants despite the lack of macrosynteny. *Mol Genet Genomics* **274**(6): 644-657.
- Khatabi B, Molitor A, Lindermayr C, Piffi S, Durner J, von Wettstein D, Kogel KH, Schafer P. 2012.** Ethylene supports colonization of plant roots by the mutualistic fungus *Piriformospora indica*. *PLoS One* **7**(4): e35502.
- Kistner C, Parniske M. 2002.** Evolution of signal transduction in intracellular symbiosis. *Trends Plant Sci* **7**(11): 511-518.

- Kistner C, Winzer T, Pitzschke A, Mulder L, Sato S, Kaneko T, Tabata S, Sandal N, Stougaard J, Webb KJ, et al. 2005.** Seven *Lotus japonicus* genes required for transcriptional reprogramming of the root during fungal and bacterial symbiosis. *Plant Cell* **17**(8): 2217-2229.
- Kloppholz S, Kuhn H, Requena N. 2011.** A secreted fungal effector of *Glomus intraradices* promotes symbiotic biotrophy. *Curr Biol* **21**(14): 1204-1209.
- Kobae Y, Tamura Y, Takai S, Banba M, Hata S. 2010.** Localized expression of arbuscular mycorrhiza-inducible ammonium transporters in soybean. *Plant Cell Physiol* **51**(9): 1411-1415.
- Koch E, Slusarenko A. 1990.** *Arabidopsis* is susceptible to infection by a downy mildew fungus. *Plant Cell* **2**(5): 437-445.
- Koornneef A, Pieterse CM. 2008.** Cross talk in defence signaling. *Plant Physiol* **146**(3): 839-844.
- Kosuta S, Held M, Hossain MS, Morieri G, Macgillivray A, Johansen C, Antolin-Llovera M, Parniske M, Oldroyd GE, Downie AJ, et al. 2011.** *Lotus japonicus* symRK-14 uncouples the cortical and epidermal symbiotic program. *Plant J* **67**(5): 929-940.
- Kruger M, Kruger C, Walker C, Stockinger H, Schussler A. 2012.** Phylogenetic reference data for systematics and phylotaxonomy of arbuscular mycorrhizal fungi from phylum to species level. *New Phytol* **193**(4): 970-984.
- Kumar M, Yadav V, Kumar H, Sharma R, Singh A, Tuteja N, Johri AK. 2011.** *Piriformospora indica* enhances plant growth by transferring phosphate. *Plant Signal Behav* **6**(5): 723-725.
- Kumar M, Yadav V, Tuteja N, Johri AK. 2009.** Antioxidant enzyme activities in maize plants colonized with *Piriformospora indica*. *Microbiology* **155**(Pt 3): 780-790.
- Lahrman U, Ding Y, Banhara A, Rath M, Hajirezaei MR, Dohlemann S, von Wiren N, Parniske M, Zuccaro A. 2013.** Host-related metabolic cues affect colonization strategies of a root endophyte. *Proc Natl Acad Sci U S A* **110**(34): 13965-13970.
- Lahrman U, Zuccaro A. 2012.** *Opprimo ergo sum*--evasion and suppression in the root endophytic fungus *Piriformospora indica*. *Mol Plant Microbe Interact* **25**(6): 727-737.
- Laine A-L. 2004.** A powdery mildew infection on a shared host plant affects the dynamics of the *Glanville fritillaria* butterfly populations. *Oikos* **107**(2): 329-337.
- Landoni M, De Francesco A, Bellatti S, Delledonne M, Ferrarini A, Venturini L, Pulu R, Bononi M, Tonelli C. 2013.** A mutation in the *FZL* gene of *Arabidopsis* causing alteration in chloroplast morphology results in a lesion mimic phenotype. *J Exp Bot* **64**(14): 4313-4328.
- Lee YC, Johnson JM, Chien CT, Sun C, Cai D, Lou B, Oelmuller R, Yeh KW. 2011.** Growth promotion of Chinese cabbage and *Arabidopsis* by *Piriformospora indica* is not stimulated by mycelium-synthesized auxin. *Mol Plant Microbe Interact* **24**(4): 421-431.
- Li X, Clarke JD, Zhang Y, Dong X. 2001.** Activation of an EDS1-mediated R-gene pathway in the *snc1* mutant leads to constitutive, NPR1-independent pathogen resistance. *Mol Plant Microbe Interact* **14**(10): 1131-1139.
- Lin NC, Martin GB. 2005.** An *avrPto/avrPtoB* mutant of *Pseudomonas syringae* pv. *tomato* DC3000 does not elicit Pto-mediated resistance and is less virulent on tomato. *Mol Plant Microbe Interact* **18**(1): 43-51.
- Litman GW, Rast JP, Fugmann SD. 2010.** The origins of vertebrate adaptive immunity. *Nature reviews. Immunology* **10**(8): 543-553.
- Livak KJ, Schmittgen TD. 2001.** Analysis of relative gene expression data using real-time quantitative PCR and the 2⁻($\Delta\Delta C_T$) Method. *Methods* **25**(4): 402-408.

- Lopez-Gomez M, Sandal N, Stougaard J, Boller T. 2011.** Interplay of flg22-induced defence responses and nodulation in *Lotus japonicus*. *J Exp Bot*.
- Lopez-Pedrosa A, Gonzalez-Guerrero M, Valderas A, Azcon-Aguilar C, Ferrol N. 2006.** *GintAMT1* encodes a functional high-affinity ammonium transporter that is expressed in the extraradical mycelium of *Glomus intraradices*. *Fungal Genet Biol* **43**(2): 102-110.
- Lorrain S, Vaillau F, Balague C, Roby D. 2003.** Lesion mimic mutants: keys for deciphering cell death and defence pathways in plants? *Trends Plant Sci* **8**(6): 263-271.
- Maeda D, Ashida K, Iguchi K, Chechetka SA, Hijikata A, Okusako Y, Deguchi Y, Izui K, Hata S. 2006.** Knockdown of an arbuscular mycorrhiza-inducible phosphate transporter gene of *Lotus japonicus* suppresses mutualistic symbiosis. *Plant Cell Physiol* **47**(7): 807-817.
- Maldonado-Mendoza IE, Dewbre GR, Harrison MJ. 2001.** A phosphate transporter gene from the extra-radical mycelium of an arbuscular mycorrhizal fungus *Glomus intraradices* is regulated in response to phosphate in the environment. *Mol Plant Microbe Interact* **14**(10): 1140-1148.
- Margulis L, Schwartz KV. 2002.** *Five kingdoms : an illustrated guide to the phyla of life on earth*. New York: Freeman/Owl Book.
- Markmann K, Giczey G, Parniske M. 2008.** Functional adaptation of a plant receptor-kinase paved the way for the evolution of intracellular root symbioses with bacteria. *PLoS Biol* **6**(3): e68.
- Mena-Violante HG, Ocampo-Jimenez O, Dendooven L, Martinez-Soto G, Gonzalez-Castaneda J, Davies FT, Jr., Olalde-Portugal V. 2006.** Arbuscular mycorrhizal fungi enhance fruit growth and quality of chile ancho (*Capsicum annum* L. cv San Luis) plants exposed to drought. *Mycorrhiza* **16**(4): 261-267.
- Mendgen K, Hahn M. 2002.** Plant infection and the establishment of fungal biotrophy. *Trends Plant Sci* **7**(8): 352-356.
- Micali C, Göllner K, Humphry M, Consonni C, Panstruga R. 2008.** The Powdery Mildew Disease of *Arabidopsis*: A Paradigm for the Interaction between Plants and Biotrophic Fungi. *The Arabidopsis Book / American Society of Plant Biologists* **6**: e0115.
- Mikkelsen MD, Fuller VL, Hansen BG, Nafisi M, Olsen CE, Nielsen HB, Halkier BA. 2009.** Controlled indole-3-acetaldoxime production through ethanol-induced expression of CYP79B2. *Planta* **229**(6): 1209-1217.
- Miklis M, Consonni C, Bhat RA, Lipka V, Schulze-Lefert P, Panstruga R. 2007.** Barley MLO modulates actin-dependent and actin-independent antifungal defence pathways at the cell periphery. *Plant Physiol* **144**(2): 1132-1143.
- Mims CW, Richardson EA, Holt Iii BF, Dangl JL. 2004.** Ultrastructure of the host pathogen interface in *Arabidopsis thaliana* leaves infected by the downy mildew *Hyaloperonospora parasitica*. *Can J Bot* **82**(7): 1001-1008.
- Miransari M. 2010.** Contribution of arbuscular mycorrhizal symbiosis to plant growth under different types of soil stress. *Plant Biol* **12**(4): 563-569.
- Miwa H, Sun J, Oldroyd GE, Downie JA. 2006.** Analysis of Nod-factor-induced calcium signaling in root hairs of symbiotically defective mutants of *Lotus japonicus*. *Mol Plant Microbe Interact* **19**(8): 914-923.
- Molitor A, Zajic D, Voll LM, Pons KHJ, Samans B, Kogel KH, Waller F. 2011.** Barley leaf transcriptome and metabolite analysis reveals new aspects of compatibility and *Piriformospora indica*-mediated systemic induced resistance to powdery mildew. *Mol Plant Microbe Interact* **24**(12): 1427-1439.
- Mondy N, Corio-Costet M-F. 2004.** Feeding insects with a phytopathogenic fungus influences their diapause and population dynamics. *Ecol Entomol* **29**(6): 711-717.

- Moran PJ, Thompson GA. 2001.** Molecular responses to aphid feeding in *Arabidopsis* in relation to plant defence pathways. *Plant Physiol* **125**(2): 1074-1085.
- Mosher S, Seybold H, Rodriguez P, Stahl M, Davies KA, Dayaratne S, Morillo SA, Wierzba M, Favery B, Keller H, et al. 2013.** The tyrosine-sulfated peptide receptors PSKR1 and PSY1R modify the immunity of *Arabidopsis* to biotrophic and necrotrophic pathogens in an antagonistic manner. *Plant J* **73**(3): 469-482.
- Mukhopadhyay S, Jackson PK. 2011.** The tubby family proteins. *Genome Biol* **12**(6): 225.
- Murashige T, Skoog F. 1962.** A revised medium for rapid growth and bio assays with tobacco tissue cultures. *Physiol Plant* **15**(3): 473-497.
- Musser RO, Cipollini DF, Hum-Musser SM, Williams SA, Brown JK, Felton GW. 2005.** Evidence that the caterpillar salivary enzyme glucose oxidase provides herbivore offense in solanaceous plants. *Arch Insect Biochem Physiol* **58**(2): 128-137.
- Muthamilarasan M, Prasad M. 2013.** Plant innate immunity: an updated insight into defence mechanism. *J Biosci* **38**(2): 433-449.
- Nagy R, Karandashov V, Chague V, Kalinkevich K, Tamasloukht M, Xu G, Jakobsen I, Levy AA, Amrhein N, Bucher M. 2005.** The characterization of novel mycorrhiza-specific phosphate transporters from *Lycopersicon esculentum* and *Solanum tuberosum* uncovers functional redundancy in symbiotic phosphate transport in solanaceous species. *Plant J* **42**(2): 236-250.
- Nongbri PL, Johnson JM, Sherameti I, Glawischnig E, Halkier BA, Oelmüller R. 2012.** Indole-3-acetaldoxime-derived compounds restrict root colonization in the beneficial interaction between *Arabidopsis* roots and the endophyte *Piriformospora indica*. *Mol Plant Microbe Interact* **25**(9): 1186-1197.
- Nouri E, Breuillin-Sessoms F, Feller U, Reinhardt D. 2014.** Phosphorus and nitrogen regulate arbuscular mycorrhizal symbiosis in *Petunia hybrida*. *PLoS One* **9**(6): e90841.
- Oelmüller R, Sherameti I, Tripathi S, Varma A. 2009.** *Piriformospora indica*, a cultivable root endophyte with multiple biotechnological applications. *Symbiosis* **49**(1): 1-17.
- Okmen B, Doehlemann G. 2014.** Inside plant: biotrophic strategies to modulate host immunity and metabolism. *Curr Opin Plant Biol* **20C**: 19-25.
- Oldroyd GE. 2013.** Speak, friend, and enter: signalling systems that promote beneficial symbiotic associations in plants. *Nat Rev Microbiol* **11**(4): 252-263.
- Olsson PA, Burleigh SH, van Aarle IM. 2005.** The influence of external nitrogen on carbon allocation to *Glomus intraradices* in monoxenic arbuscular mycorrhiza. *New Phytol* **168**(3): 677-686.
- Olsson PA, Rahm J, Aliasgharzad N. 2010.** Carbon dynamics in mycorrhizal symbioses is linked to carbon costs and phosphorus benefits. *FEMS Microbiol Ecol* **72**(1): 125-131.
- Pareja-Jaime Y, Roncero MI, Ruiz-Roldan MC. 2008.** Tomatinase from *Fusarium oxysporum* f. sp. *lycopersici* is required for full virulence on tomato plants. *Mol Plant Microbe Interact* **21**(6): 728-736.
- Parniske M. 2000.** Intracellular accommodation of microbes by plants: a common developmental program for symbiosis and disease? *Curr Opin Plant Biol* **3**(4): 320-328.
- Parniske M. 2008.** Arbuscular mycorrhiza: the mother of plant root endosymbioses. *Nat Rev Microbiol* **6**(10): 763-775.
- Paszkowski U, Kroken S, Roux C, Briggs SP. 2002.** Rice phosphate transporters include an evolutionarily divergent gene specifically activated in arbuscular mycorrhizal symbiosis. *Proc Natl Acad Sci U S A* **99**(20): 13324-13329.
- Pawlowski K, Bisseling T. 1996.** Rhizobial and actinorhizal symbioses: what are the shared features? *Plant Cell* **8**(10): 1899-1913.

- Pedersen BP, Kumar H, Waight AB, Risenmay AJ, Roe-Zurz Z, Chau BH, Schlessinger A, Bonomi M, Harries W, Sali A, et al. 2013.** Crystal structure of a eukaryotic phosphate transporter. *Nature* **496**(7446): 533-536.
- Perez-Tienda J, Testillano PS, Balestrini R, Fiorilli V, Azcon-Aguilar C, Ferrol N. 2011.** GintAMT2, a new member of the ammonium transporter family in the arbuscular mycorrhizal fungus *Glomus intraradices*. *Fungal Genet Biol* **48**(11): 1044-1055.
- Perry J, Brachmann A, Welham T, Binder A, Charpentier M, Groth M, Haage K, Markmann K, Wang TL, Parniske M. 2009.** TILLING in *Lotus japonicus* identified large allelic series for symbiosis genes and revealed a bias in functionally defective ethyl methanesulfonate alleles toward glycine replacements. *Plant Physiol* **151**(3): 1281-1291.
- Peškan-Berghöfer T, Shahollari B, Giong PH, Hehl S, Markert C, Blanke V, Kost G, Varma A, Oelmüller R. 2004.** Association of *Piriformospora indica* with *Arabidopsis thaliana* roots represents a novel system to study beneficial plant-microbe interactions and involves early plant protein modifications in the endoplasmic reticulum and at the plasma membrane. *Physiologia Plantarum* **122**(4): 465-477.
- Pham G, Kumari R, Singh A, Malla R, Prasad R, Sachdev M, Kaldorf M, Buscot F, Oelmüller R, Hampp R, et al. 2004.** Axenic Culture of Symbiotic Fungus *Piriformospora indica*. In: Varma A, Abbott L, Werner D, Hampp R eds. *Plant Surface Microbiology*: Springer Berlin Heidelberg, 593-613.
- Preston CA, Lewandowski C, Enyedi AJ, Baldwin IT. 1999.** Tobacco mosaic virus inoculation inhibits wound-induced jasmonic acid-mediated responses within but not between plants. *Planta* **209**(1): 87-95.
- Pumplin N, Harrison MJ. 2009.** Live-cell imaging reveals periarbuscular membrane domains and organelle location in *Medicago truncatula* roots during arbuscular mycorrhizal symbiosis. *Plant Physiol* **151**(2): 809-819.
- Qiang X, Zechmann B, Reitz MU, Kogel KH, Schafer P. 2012.** The mutualistic fungus *Piriformospora indica* colonizes *Arabidopsis* roots by inducing an endoplasmic reticulum stress-triggered caspase-dependent cell death. *Plant Cell* **24**(2): 794-809.
- Radutoiu S, Madsen LH, Madsen EB, Felle HH, Umehara Y, Gronlund M, Sato S, Nakamura Y, Tabata S, Sandal N, et al. 2003.** Plant recognition of symbiotic bacteria requires two LysM receptor-like kinases. *Nature* **425**(6958): 585-592.
- Rauhut T, Glawischnig E. 2009.** Evolution of camalexin and structurally related indolic compounds. *Phytochemistry* **70**(15-16): 1638-1644.
- Rausch C, Daram P, Brunner S, Jansa J, Laloi M, Leggewie G, Amrhein N, Bucher M. 2001.** A phosphate transporter expressed in arbuscule-containing cells in potato. *Nature* **414**(6862): 462-470.
- Reitz MU, Bissue JK, Zocher K, Attard A, Huckelhoven R, Becker K, Imani J, Eichmann R, Schafer P. 2012.** The subcellular localization of Tubby-like proteins and participation in stress signaling and root colonization by the mutualist *Piriformospora indica*. *Plant Physiol* **160**(1): 349-364.
- Reitz MU, Pai S, Imani J, Schafer P. 2013.** New insights into the subcellular localization of Tubby-like proteins and their participation in the *Arabidopsis-Piriformospora indica* interaction. *Plant Signal Behav* **8**(8).
- Rentel MC, Lecourieux D, Ouaked F, Usher SL, Petersen L, Okamoto H, Knight H, Peck SC, Grierson CS, Hirt H, et al. 2004.** OXI1 kinase is necessary for oxidative burst-mediated signalling in *Arabidopsis*. *Nature* **427**(6977): 858-861.
- Ried MK, Antolin-Llovera M, Parniske M. 2014.** Spontaneous symbiotic reprogramming of plant roots triggered by receptor-like kinases. **3**.

- Röder G, Rahier M, Naisbit RE. 2007.** Coping with an antagonist: the impact of a phytopathogenic fungus on the development and behaviour of two species of alpine leaf beetle. *Oikos* **116**(9): 1514-1523.
- Rodriguez-Saona C, Chalmers JA, Raj S, Thaler JS. 2005.** Induced plant responses to multiple damagers: differential effects on an herbivore and its parasitoid. *Oecologia* **143**(4): 566-577.
- Rosewarne GM, Barker SJ, Smith SE. 1997.** Production of near-synchronous fungal colonization in tomato for developmental and molecular analyses of mycorrhiza. *Mycol Res* **101**(8): 966-970.
- Rostás M, Hilker M. 2002.** Asymmetric plant-mediated cross-effects between a herbivorous insect and a phytopathogenic fungus. *Agric Forest Entomol* **4**(3): 223-231.
- Ryals JA, Neuenschwander UH, Willits MG, Molina A, Steiner HY, Hunt MD. 1996.** Systemic acquired resistance. *Plant Cell* **8**(10): 1809-1819.
- Saito K, Yoshikawa M, Yano K, Miwa H, Uchida H, Asamizu E, Sato S, Tabata S, Imaizumi-Anraku H, Umehara Y, et al. 2007.** NUCLEOPORIN85 is required for calcium spiking, fungal and bacterial symbioses, and seed production in *Lotus japonicus*. *Plant Cell* **19**(2): 610-624.
- Salvaudon L, Héraudet V, Shykoff JA. 2008.** *Arabidopsis thaliana* and the Robin Hood parasite: a chivalrous oomycete that steals fitness from fecund hosts and benefits the poorest one? *Biology Lett* **4**(5): 526-529.
- Santagata S, Boggon TJ, Baird CL, Gomez CA, Zhao J, Shan WS, Myszka DG, Shapiro L. 2001.** G-protein signaling through tubby proteins. *Science* **292**(5524): 2041-2050.
- Savita V, Ajit V, Karl-Heinz R, Annette H, Gerhard K, Ashok S, Prakash B, Britta B, Philipp F. 1998.** *Piriformospora indica*, gen. et sp. nov., a new root-colonizing fungus. *Mycologia* **90**(5): 896-903.
- Schäfer P, Pffiffi S, Voll LM, Zajic D, Chandler PM, Waller F, Scholz U, Pons-Kühnemann J, Sonnewald S, Sonnewald U, et al. 2009.** Manipulation of plant innate immunity and gibberellin as factor of compatibility in the mutualistic association of barley roots with *Piriformospora indica*. *The Plant Journal* **59**(3): 461-474.
- Schauser L, Handberg K, Sandal N, Stiller J, Thykjaer T, Pajuelo E, Nielsen A, Stougaard J. 1998.** Symbiotic mutants deficient in nodule establishment identified after T-DNA transformation of *Lotus japonicus*. *Mol Gen Genet* **259**(4): 414-423.
- Schauser L, Roussis A, Stiller J, Stougaard J. 1999.** A plant regulator controlling development of symbiotic root nodules. *Nature* **402**(6758): 191-195.
- Schlicht M, Kombrink E. 2013.** The role of nitric oxide in the interaction of *Arabidopsis thaliana* with the biotrophic fungi, *Golovinomyces orontii* and *Erysiphe pisi*. *Front Plant Sci* **4**: 351.
- Schneider CA, Rasband WS, Eliceiri KW. 2012.** NIH Image to ImageJ: 25 years of image analysis. *Nat Methods* **9**(7): 671-675.
- Scholl RL, May ST, Ware DH. 2000.** Seed and molecular resources for *Arabidopsis*. *Plant Physiol* **124**(4): 1477-1480.
- Schultheiss H, Preuss J, Pircher T, Eichmann R, Huckelhoven R. 2008.** Barley RIC171 interacts with RACB in planta and supports entry of the powdery mildew fungus. *Cell Microbiol* **10**(9): 1815-1826.
- Schulze-Lefert P. 2004.** Knocking on the heaven's wall: pathogenesis of and resistance to biotrophic fungi at the cell wall. *Curr Opin Plant Biol* **7**(4): 377-383.
- Schüßler A, Walker C. 2011.** Evolution of the 'plant-symbiotic' fungal phylum, Glomeromycota. In: Pöggeler S, Wöstemeyer J eds. *Evolution of fungi and fungal-like organisms*: Springer Berlin Heidelberg, 163-185.

- Schwarzott D, Walker C, Schussler A. 2001.** *Glomus*, the largest genus of the arbuscular mycorrhizal fungi (Glomales), is nonmonophyletic. *Mol Phylogenet Evol* **21**(2): 190-197.
- Selosse MA, Dubois MP, Alvarez N. 2009.** Do Sebaciales commonly associate with plant roots as endophytes? *Mycol Res* **113**(Pt 10): 1062-1069.
- Shahollari B, Vadassery J, Varma A, Oelmuller R. 2007.** A leucine-rich repeat protein is required for growth promotion and enhanced seed production mediated by the endophytic fungus *Piriformospora indica* in *Arabidopsis thaliana*. *Plant J* **50**(1): 1-13.
- Sherameti I, Shahollari B, Venus Y, Altschmied L, Varma A, Oelmuller R. 2005.** The endophytic fungus *Piriformospora indica* stimulates the expression of nitrate reductase and the starch-degrading enzyme glucan-water dikinase in tobacco and *Arabidopsis* roots through a homeodomain transcription factor that binds to a conserved motif in their promoters. *J Biol Chem* **280**(28): 26241-26247.
- Sherameti I, Tripathi S, Varma A, Oelmuller R. 2008.** The root-colonizing endophyte *Piriformospora indica* confers drought tolerance in *Arabidopsis* by stimulating the expression of drought stress-related genes in leaves. *Mol Plant Microbe Interact* **21**(6): 799-807.
- Shirano Y, Kachroo P, Shah J, Klessig DF. 2002.** A gain-of-function mutation in an *Arabidopsis* Toll Interleukin1 receptor-nucleotide binding site-leucine-rich repeat type R gene triggers defence responses and results in enhanced disease resistance. *Plant Cell* **14**(12): 3149-3162.
- Shiu SH, Bleeker AB. 2003.** Expansion of the receptor-like kinase/Pelle gene family and receptor-like proteins in *Arabidopsis*. *Plant Physiol* **132**(2): 530-543.
- Singh S, Katzer K, Lambert J, Cerri M, Parniske M. 2014.** CYCLOPS, a DNA-binding transcriptional activator, orchestrates symbiotic root nodule development. *Cell Host Microbe* **15**(2): 139-152.
- Singh S, Parniske M. 2012.** Activation of calcium- and calmodulin-dependent protein kinase (CCaMK), the central regulator of plant root endosymbiosis. *Curr Opin Plant Biol* **15**(4): 444-453.
- Sirrenberg A, Gobel C, Grond S, Czempinski N, Ratzinger A, Karlovsky P, Santos P, Feussner I, Pawlowski K. 2007.** *Piriformospora indica* affects plant growth by auxin production. *Physiol Plant* **131**(4): 581-589.
- Slusarenko AJ, Schlaich NL. 2003.** Downy mildew of *Arabidopsis thaliana* caused by *Hyaloperonospora parasitica* (formerly *Peronospora parasitica*). *Mol Plant Pathol* **4**(3): 159-170.
- Smith SE, Read DJ. 1997.** *Mycorrhizal symbiosis*. San Diego; London: Academic Press.
- Smith SE, Read DJ. 2008.** *Mycorrhizal symbiosis*. Amsterdam; London: Academic.
- Smith SE, Smith FA. 2011.** Roles of arbuscular mycorrhizas in plant nutrition and growth: new paradigms from cellular to ecosystem scales. *Annu Rev Plant Biol* **62**: 227-250.
- Sohn KH, Lei R, Nemri A, Jones JD. 2007.** The downy mildew effector proteins ATR1 and ATR13 promote disease susceptibility in *Arabidopsis thaliana*. *Plant Cell* **19**(12): 4077-4090.
- Solaiman MZ, Ezawa T, Kojima T, Saito M. 1999.** Polyphosphates in intraradical and extraradical hyphae of an arbuscular mycorrhizal fungus, *Gigaspora margarita*. *Appl Environ Microbiol* **65**(12): 5604-5606.
- Solano R, Stepanova A, Chao Q, Ecker JR. 1998.** Nuclear events in ethylene signaling: a transcriptional cascade mediated by ETHYLENE-INSENSITIVE3 and ETHYLENE-RESPONSE-FACTOR1. *Genes Dev* **12**(23): 3703-3714.
- Spanu PD, Abbott JC, Amselem J, Burgis TA, Soanes DM, Stuber K, Ver Loren van Themaat E, Brown JK, Butcher SA, Gurr SJ, et al. 2010.** Genome expansion and gene loss in powdery mildew fungi reveal tradeoffs in extreme parasitism. *Science* **330**(6010): 1543-1546.

- Spoel SH, Johnson JS, Dong X. 2007.** Regulation of tradeoffs between plant defences against pathogens with different lifestyles. *Proc Natl Acad Sci U S A* **104**(47): 18842-18847.
- Stamatakis A, Hoover P, Rougemont J. 2008.** A rapid bootstrap algorithm for the RAxML Web servers. *Syst Biol* **57**(5): 758-771.
- Stassen JH, Van den Ackerveken G. 2011.** How do oomycete effectors interfere with plant life? *Curr Opin Plant Biol* **14**(4): 407-414.
- Stein E, Molitor A, Kogel KH, Waller F. 2008.** Systemic resistance in *Arabidopsis* conferred by the mycorrhizal fungus *Piriformospora indica* requires jasmonic acid signaling and the cytoplasmic function of NPR1. *Plant Cell Physiol* **49**(11): 1747-1751.
- Stout MJ, Fidantsef AL, Duffey SS, Bostock RM. 1999.** Signal interactions in pathogen and insect attack: systemic plant-mediated interactions between pathogens and herbivores of the tomato, *Lycopersicon esculentum*. *Physiol Mol Plant P* **54**(3-4): 115-130.
- Stout MJ, Thaler JS, Thomma BP. 2006.** Plant-mediated interactions between pathogenic microorganisms and herbivorous arthropods. *Annu Rev Entomol* **51**: 663-689.
- Stout MJ, Workman KV, Bostock RM, Duffey SS. 1998.** Specificity of induced resistance in the tomato, *Lycopersicon esculentum*. *Oecologia* **113**(1): 74-81.
- Stracke S, Kistner C, Yoshida S, Mulder L, Sato S, Kaneko T, Tabata S, Sandal N, Stougaard J, Szczyglowski K, et al. 2002.** A plant receptor-like kinase required for both bacterial and fungal symbiosis. *Nature* **417**(6892): 959-962.
- Suzaki T, Kim CS, Takeda N, Szczyglowski K, Kawaguchi M. 2013.** *TRICOT* encodes an AMP1-related carboxypeptidase that regulates root nodule development and shoot apical meristem maintenance in *Lotus japonicus*. *Development* **140**(2): 353-361.
- Szczyglowski K, Shaw RS, Wopereis J, Copeland S, Hamburger D, Kasiborski B, Dazzo FB, de Bruijn FJ. 1998.** Nodule organogenesis and symbiotic mutants of the model legume *Lotus japonicus*. *Molecular Plant-Microbe Interactions* **11**(7): 684-697.
- Team RC 2013.** R: A language and environment for statistical computing. Vienna, Austria: R Foundation for Statistical Computing.
- Thaler JS, Agrawal AA, Halitschke R. 2010.** Salicylate-mediated interactions between pathogens and herbivores. *Ecology* **91**(4): 1075-1082.
- Thaler JS, Bostock RM. 2004.** Interaction between abscisic-acid-mediated responses and plant resistance to pathogens and insects *Ecology* **85**(1): 48-58.
- Thompson GA, Goggin FL. 2006.** Transcriptomics and functional genomics of plant defence induction by phloem-feeding insects. *J Exp Bot* **57**(4): 755-766.
- Tian C, Kasiborski B, Koul R, Lammers PJ, Bucking H, Shachar-Hill Y. 2010.** Regulation of the nitrogen transfer pathway in the arbuscular mycorrhizal symbiosis: gene characterization and the coordination of expression with nitrogen flux. *Plant Physiol* **153**(3): 1175-1187.
- Tinker PB, Nye PH. 2000.** *Solute movement in the rhizosphere*. New York; Oxford: Oxford University Press.
- Toussaint JP, St-Arnaud M, Charest C. 2004.** Nitrogen transfer and assimilation between the arbuscular mycorrhizal fungus *Glomus intraradices* Schenck & Smith and Ri T-DNA roots of *Daucus carota* L. in an *in vitro* compartmented system. *Can J Microbiol* **50**(4): 251-260.
- Vadassery J, Ritter C, Venus Y, Camehl I, Varma A, Shahollari B, Novak O, Strnad M, Ludwig-Muller J, Oelmuller R. 2008.** The role of auxins and cytokinins in the mutualistic interaction between *Arabidopsis* and *Piriformospora indica*. *Mol Plant Microbe Interact* **21**(10): 1371-1383.

- Van Damme M, Andel A, Huibers RP, Panstruga R, Weisbeek PJ, Van den Ackerveken G. 2005.** Identification of *Arabidopsis loci* required for susceptibility to the downy mildew pathogen *Hyaloperonospora parasitica*. *Mol Plant Microbe Interact* **18**(6): 583-592.
- van Damme M, Zeilmaier T, Elberse J, Andel A, de Sain-van der Velden M, van den Ackerveken G. 2009.** Downy mildew resistance in *Arabidopsis* by mutation of HOMOSERINE KINASE. *Plant Cell* **21**(7): 2179-2189.
- van Schie CC, Takken FL. 2014.** Susceptibility genes 101: how to be a good host. *Annu Rev Phytopathol* **52**: 551-581.
- Varallyay E, Giczey G, Burgyan J. 2012.** Virus-induced gene silencing of *Mlo* genes induces powdery mildew resistance in *Triticum aestivum*. *Arch Virol* **157**(7): 1345-1350.
- Venkateshwaran M, Cosme A, Han L, Banba M, Satyshur KA, Schleiff E, Parniske M, Imaizumi-Anraku H, Ane JM. 2012.** The recent evolution of a symbiotic ion channel in the legume family altered ion conductance and improved functionality in calcium signaling. *Plant Cell* **24**(6): 2528-2545.
- Venus Y, Oelmüller R. 2013.** *Arabidopsis* ROP1 and ROP6 influence germination time, root morphology, the formation of F-actin bundles, and symbiotic fungal interactions. *Mol Plant* **6**(3): 872-886.
- Villegas J, Williams RD, Nantais L, Archambault J, Fortin JA. 1996.** Effects of N source on pH and nutrient exchange of extramatrical mycelium in a mycorrhizal Ri T-DNA transformed root system. *Mycorrhiza* **6**(4): 247-251.
- Voegelé RT, Mendgen K. 2003.** Rust haustoria: nutrient uptake and beyond. *New Phytol* **159**(1): 93-100.
- Vogel J, Somerville S. 2000.** Isolation and characterization of powdery mildew-resistant *Arabidopsis* mutants. *Proc Natl Acad Sci U S A* **97**(4): 1897-1902.
- Vogel JP, Raab TK, Schiff C, Somerville SC. 2002.** *PMR6*, a pectate lyase-like gene required for powdery mildew susceptibility in *Arabidopsis*. *Plant Cell* **14**(9): 2095-2106.
- Vogel JP, Raab TK, Somerville CR, Somerville SC. 2004.** Mutations in *PMR5* result in powdery mildew resistance and altered cell wall composition. *Plant J* **40**(6): 968-978.
- Waller F, Achatz B, Baltruschat H, Fodor J, Becker K, Fischer M, Heier T, Huckelhoven R, Neumann C, von Wettstein D, et al. 2005.** The endophytic fungus *Piriformospora indica* reprograms barley to salt-stress tolerance, disease resistance, and higher yield. *Proc Natl Acad Sci U S A* **102**(38): 13386-13391.
- Waller F, Mukherjee K, Deshmukh SD, Achatz B, Sharma M, Schafer P, Kogel KH. 2008.** Systemic and local modulation of plant responses by *Piriformospora indica* and related Sebaciniales species. *J Plant Physiol* **165**(1): 60-70.
- Wang Y, Meng Y, Zhang M, Tong X, Wang Q, Sun Y, Quan J, Govers F, Shan W. 2011.** Infection of *Arabidopsis thaliana* by *Phytophthora parasitica* and identification of variation in host specificity. *Mol Plant Pathol* **12**(2): 187-201.
- Webb KJ, Skot L, Nicholson MN, Jørgensen B, Mizen S. 2000.** *Mesorhizobium loti* increases root-specific expression of a calcium-binding protein homologue identified by promoter tagging in *Lotus japonicus*. *Mol Plant Microbe Interact* **13**(6): 606-616.
- Weis C, Hildebrandt U, Hoffmann T, Hemetsberger C, Pfeilmeier S, König C, Schwab W, Eichmann R, Huckelhoven R. 2014.** CYP83A1 is required for metabolic compatibility of *Arabidopsis* with the adapted powdery mildew fungus *Erysiphe cruciferarum*. *New Phytol* **202**(4): 1310-1319.
- Weiss M, Selosse MA, Rexer KH, Urban A, Oberwinkler F. 2004.** Sebaciniales: a hitherto overlooked cosm of heterobasidiomycetes with a broad mycorrhizal potential. *Mycol Res* **108**(Pt 9): 1003-1010.

- Wiermer M, Cheng YT, Imkampe J, Li M, Wang D, Lipka V, Li X. 2012.** Putative members of the *Arabidopsis* Nup107-160 nuclear pore sub-complex contribute to pathogen defence. *Plant J* **70**(5): 796-808.
- Yadav V, Kumar M, Deep DK, Kumar H, Sharma R, Tripathi T, Tuteja N, Saxena AK, Johri AK. 2010.** A phosphate transporter from the root endophytic fungus *Piriformospora indica* plays a role in phosphate transport to the host plant. *J Biol Chem* **285**(34): 26532-26544.
- Yi M, Valent B. 2013.** Communication between filamentous pathogens and plants at the biotrophic interface. *Annu Rev Phytopathol* **51**: 587-611.
- Zhang Y, Li X. 2005.** A putative nucleoporin 96 Is required for both basal defence and constitutive resistance responses mediated by suppressor of npr1-1, constitutive 1. *Plant Cell* **17**(4): 1306-1316.
- Zheng Z, Nonomura T, Appiano M, Pavan S, Matsuda Y, Toyoda H, Wolters AM, Visser RG, Bai Y. 2013.** Loss of function in *Mlo* orthologs reduces susceptibility of pepper and tomato to powdery mildew disease caused by *Leveillula taurica*. *PLoS One* **8**(7): e70723.
- Zuccaro A, Lahrmann U, Guldener U, Langen G, Pfiffi S, Biedenkopf D, Wong P, Samans B, Grimm C, Basiewicz M, et al. 2011.** Endophytic life strategies decoded by genome and transcriptome analyses of the mutualistic root symbiont *Piriformospora indica*. *PLoS Pathog* **7**(10): e1002290.

11. List of figures

Figure 1 – Schematic representation of intracellular accommodation organs in symbiosis and disease	19
Figure 2: Comparison of gene structures and protein domains of <i>L. japonicus</i> common symbiosis genes (CSGs) and their closest homologs (HCSGs) in <i>A. thaliana</i>.....	32
Figure 3: <i>A. thaliana</i> HCSGs with encoded proteins and their respective identities/similarities to their <i>L. japonicus</i> homologs.....	33
Figure 4: Maximum likelihood phylogenetic trees of MLD-LRR-RLKs from <i>A. thaliana</i> and <i>L. japonicus</i>.....	34
Figure 5: Mutation of <i>A. thaliana</i> HCSGs reduces the reproductive success of <i>Hpa</i>	36
Figure 6: Reproductive success of <i>Hpa</i> is reduced in the HSCG mutants.....	37
Figure 7: Reproductive success of <i>Hpa</i> is not affected in the <i>seh1</i>, <i>nup43</i>, <i>nup85</i> and <i>nup160</i> mutants	38
Figure 8: <i>Hpa</i> haustoria formation on <i>A. thaliana</i> HCSG mutants.....	39
Figure 9: <i>Hpa</i> haustorium morphology is altered in <i>A. thaliana</i> HCSG mutants .	40
Figure 10: Frequency of multilobed haustoria on <i>A. thaliana</i> wild-type and on the HSCG mutants.....	41
Figure 11: <i>Hpa</i> haustorium morphology is altered in <i>A. thaliana</i> HCSG mutants.	42
Figure 12: <i>Hpa</i> haustoria morphology	43
Figure 13: Time-course of haustoria development in <i>A. thaliana</i> wild-type and in <i>shrk1</i> x <i>shrk2</i>	45
Figure 14: Quantification of haustoria with fluorescence on <i>ProAt3g61260:YFP-At3g61260</i> plants	46
Figure 15: Mean number of conidiophores per leaf on <i>A. thaliana</i> HCSG mutants relative to the wild-type.....	47
Figure 16: Basal expression levels and flg22-mediated induction of PAMP-responsive genes in <i>A. thaliana</i> HCSG mutants are not different from the wild-type	49

Figure 17: Basal expression levels and flg22-mediated induction of PAMP-responsive genes in <i>A. thaliana</i> HCSG mutants are not different from the wild-type	50
Figure 18: Transcript levels of defence marker gene <i>PR1</i> in <i>A. thaliana</i> HCSG mutants	51
Figure 19: <i>A. thaliana</i> HCSG mutants do not show increased levels of spontaneous or pathogen-induced cell death	53
Figure 20: <i>A. thaliana</i> HCSG mutants do not show developmental or growth defects	54
Figure 21: <i>Hpa</i> haustoria-associated callose deposition is not reduced in the <i>A. thaliana</i> HCSG mutants	55
Figure 22: <i>A. thaliana</i> HCSG mutants support similar bacterial growth curves as the wild-type	56
Figure 23: Intracellular growth of <i>P. indica</i> in root cells of <i>L. japonicus</i>	58
Figure 24: Biotrophic growth of <i>P. indica</i> in root cells of <i>L. japonicus</i>	59
Figure 25: <i>P. indica</i> sporulation inside root cells of <i>L. japonicus</i>	60
Figure 26: <i>P. indica</i> colonization in <i>A. thaliana</i> and <i>L. japonicus</i> wild-type and mutant roots	61
Figure 27: GUS expression in the <i>L. japonicus</i> symbiosis reporter line T90 inoculated with <i>P. indica</i>	62
Figure 28: Biotrophic growth of <i>P. indica</i> within <i>A. thaliana</i> root cells	64
Figure 29: Intracellular sporulation of <i>P. indica</i> in <i>A. thaliana</i> roots	65
Figure 30: Effect of <i>P. indica</i> on the growth of <i>L. japonicus</i> and <i>A. thaliana</i> grown in soil with different nutrient availabilities.	67
Figure 31: Effect of sucrose on the growth promotion of <i>A. thaliana</i> plants by <i>P. indica</i>	68
Figure 32: Influence of Ca(NO₃)₂, KNO₃, KH₂PO₄ or NH₄Cl concentrations on the growth promotion of <i>A. thaliana</i> by <i>P. indica</i>.	69
Figure 33: Biomass of <i>A. thaliana</i> wild-type or HCSG mutants upon inoculation with <i>P. indica</i> or <i>P. williamsii</i>	71

12. List of tables

Table 1: Statistical values for pairwise comparisons performed with the Kruskal-Wallis test followed by a Bonferroni-Holm correction using the mock-inoculated samples as control group.	72
Table 2: Statistical values for multiple comparisons performed the data obtained with the inoculation of <i>A. thaliana</i> HSCS mutants with either <i>P. indica</i> or <i>P. williamsii</i> with the Kruskal-Wallis test followed by a Bonferroni-Holm correction using the mock-inoculated samples as control group.	73
Table 3: <i>L. japonicus</i> common symbiosis mutants screened for phenotypes associated with <i>P. indica</i> colonization.	88
Table 4: <i>A. thaliana</i> HCSG mutants used in the screening for phenotypes associated with <i>P. indica</i> symbiosis.	88
Table 5: Oligonucleotides used in the analyses of relative fungal DNA content in <i>P. indica</i> colonized roots.	93
Table 6: Oligonucleotides used to analyse gene expression patterns before and after <i>flg22</i> treatment.	94

13. Acknowledgements

In the first place I would like to thank my PhD supervisor Martin Parniske for giving me the opportunity to work in his group on many exciting projects. I am very grateful for his invaluable support and guidance along the journey. I also appreciate his encouragement to pursue collaborations and broaden my scientific horizons. Furthermore, I owe him how greatly my ability to critically think and my writing skills improved in the years I have been under his supervision. Thank you.

I felt most welcome in the Parniske group and treasure the time I spent there. I particularly thank Laura Rose, for teaching me all issues related to *Hpa* inoculation on *A. thaliana*, and for always being so kind, supportive, and a pleasure to be around. I would like to thank Martin Groth, Kristina Haage, and Makoto Maekawa, for helping me with *Lotus japonicus* culture and AM inoculation, and Steffanie Kiebart, for helping me with production of AM spore preparations and with microscope issues.

I express my gratitude to Patrick Schäfer, for welcoming me in his laboratory to learn about *P. indica* cultivation and for sending me plates with the fungus, Alga Zuccaro, for *P. indica* strains and our close collaboration on recent manuscripts, as well as her most welcome encouraging words. I thank Arthur Schlüßer, for greatly appreciated help with the confocal microscope.

Many thanks to Iris Jarsch, for helping me with ROS assays and image analyses with imageJ, Axel Strauss, for teaching me a lot of Statistics and how to work with the “R” software, Regina Kühner and Yi Ding, for performing experiments with *P. indica*, and to my student helper Nadia Sinner for her support. I would like to show my appreciation for the administrative support from Barbara Dollrieß and Evi Pretzel.

I am especially thankful to my colleagues that directly worked with me in the lab: Martina Ried, for always being very competent and helpful in our project together, for our discussions and cheerful conversations, and for critically reading my thesis; and Andreas Binder, for designing constructs for complementation of *A. thaliana* HCSG mutants, performing experiments on *P. indica*, and for our discussions and chats, and for reviewing my thesis.

I would like to thank Caroline Gutjahr, for helping me with qPCR issues and for our internal collaboration, as well as our collaborators Andrea Gust and Thorsten

Nürnberg for receiving me in their laboratory to learn about the ROS assay; Corina Weis, Caroline Höfler, Ruth Eichmann, and Ralph Hüchelhoven, for our collaboration on assays with powdery mildew fungi and *A. thaliana*; Erik Kemen and Ulla Neumann, for performing EM on our *Hpa*-inoculated *A. thaliana* samples; and Axel Mithöfer and Andrea Lehr, for our collaboration on ecological assays involving *A. thaliana*, *Hpa* and a herbivore.

Special thanks to Ana Cosme, Meritxell Antolin-Llovera, Jayne Lambert, and Gloria Torres, for always being great company and sharing experiences and tales in and outside the lab.

I would like to thank past and present members of the lab for their helpfulness and all the fruitful discussions, as well as the warm atmosphere during work and at social events: Andreas Brachmann, Caroline Senes, David Chiasson, Gloria Torres, Gisela Brinkmann, Griet den Herder, Jasmin Gossmann, Jayne Lambert, Karl-Heinz Braun, Katharina Markmann, Katja Katzer, Macarena Marin, Manoj Kamble, Marion Cerri, Meritxell Antolin-Llovera, Myriam Charpentier, Petra Winterholler, Pryia Pimprikar, Samy Carbonell, Thomas Ott, Ute Bergmann, and Verena Klingl.

I am most grateful to Joana Bittencourt Silvestre, for our friendship and her continuous support, and for being there for me during difficult times.

I want to thank my family in Brazil, specially my parents, for their unconditional love and support. Despite the distance they have always been there, and they are part of who I am what I have achieved.

Thank you Simon, for your love and support. And above all, thank you Theo, for being my little star.

Supervisor: Prof. Dr. Marcio Alves-Ferreira

2001 – 2005 Federal University of Rio de Janeiro, Genetics: Bachelor in Biological Sciences
 Dissertation: Identification of genes differentially expressed in nodulated cowpea (*Vigna unguiculata*) in response to heat-shock.
 Supervisor: Prof. Dr. Márcia M. A. N. Pinheiro Margis

Research Projects

2007 – present – The role of common symbiosis gene homologs in the interaction of *A. thaliana* with pathogenic and symbiotic microorganisms.

2005 – 2007 – Functional Genomics of plant reproduction: characterization of new genes for use in the Biotechnology.

2003 – 2004 – Identification and cloning of genes involved in heat shock resistance: a strategy for optimization of Biological Nitrogen Fixation in tropical legumes.

Publications

Banhara A.*, Ried, MK*, Binder A, Gust, AA, Höfler C, Hüchelhoven R, Nürnberger T, Parniske M (2014). Symbiosis-related genes sustain the development of a downy mildew pathogen on *A. thaliana thaliana*. Under review.

* These authors contributed equally to the work

Lahrman U, Ding Y, **Banhara A**, Rath M, Hajirezaei MR, Döhlemann S, von Wirén N, Parniske M, Zuccaro A (2013) Host-related metabolic cues affect colonization strategies of a root endophyte. Proc Natl Acad Sci USA. 110(34): 13965-70.

Alves-Ferreira M, Wellmer F, **Banhara A**, Kumar V, Riechmann JL, Meyerowitz EM (2007) Global expression profiling applied to the analysis of *A. thaliana* stamen development. Plant Physiol. 145: 747–762.

Teaching Experience

2001 – Colegio Ph, Rio de Janeiro, Brazil, Biology Tutor for undergraduate students.

2012 – LMU München, Munich, Germany, Tutor in the practical section of the Genetik II course by Prof. Martin Parniske.

Attended Meetings and Workshops

Talk

Banhara A, Alves-Ferreira M. Characterization of an *A. thaliana* Phosphatidylinositol 3- and 4- Kinase (PI3_PI4K) Knockout Mutant with Abnormal Microsporogenesis, XXXV Meeting of Brazilian Biochemistry and Molecular Biology Society, Águas de Lindóia SP, Brazil, 2006.

Posters

Keymer A, Gutjahr C, Banhara A, Hardel S, Parniske M. DISORGANIZED ARBUSCULE is required for arbuscule branching. SFB924 (Plant Biology of the Next Generation) Conference. Freising, Germany, 2013.

Banhara A, Binder A, Groth M, Parniske M. Nucleoporins in plant-microbe symbiosis and defence signaling, 10th European Nitrogen Fixation Meeting and 1st Molecular Mycorrhiza Meeting, Munich, Germany, 2012.

Banhara A, Alves-Ferreira M. Characterization of an *A. thaliana* Phosphatidylinositol 3- and 4- Kinase (PI3_PI4K) Knockout Mutant with Abnormal Microsporogenesis. 1st Brazilian Symposium of Molecular Plant Genetics, 2006, Natal, Brazil.

Alves-Ferreira M, Wellmer F, Banhara A, Kumar V, Riechmann JL, Meyerowitz EM. Identification and characterization of new genes critical to normal *A. thaliana thaliana* stamen development by microarray and functional analysis of flower mutants *apetala-3*, *sporocyteless* and *male-sterile-1*, XIX International Congress on Sexual Plant Reproduction, Budapest, Hungary, 2006.

Banhara A, Alves-Ferreira M. Characterization of two novel genes encoding putative signal transduction components required to normal stamen development in *A. thaliana thaliana*, 51th Brazilian Congress on Genetics, 2005, Águas de Lindóia SP, Brazil, 2005.

Banhara A, de Araujo JLS. Estudo da tolerância da fixação biológica de nitrogênio a altas temperaturas através da análise do padrão de expressão gênica em nódulos de caupi (*Vigna unguiculata*) submetidos a estresse térmico, 51th Brazilian Congress on Genetics, 2005, Águas de Lindóia SP, Brazil, 2005.

Workshop

“Project Management (Natural Sciences and Medicine).” Graduate Center LMU – 24.25/10/2013.

Honors and Awards

2007 – Poster prize at the 1st Brazilian Symposium of Molecular Plant Genetics, Natal, Brazil.

2006 – “Grade A” Student Master Fellowship, Rio de Janeiro Research Council.

Fellowships

2007 – Fellowship for a 3 year PhD in Germany from the National Council for Scientific and Technological Development (CNPq) in collaboration with the Deutsche Akademischer Austauschdienst (DAAD).

*Awarded but not utilized due to funding becoming available in the host research group.

2006 – 2007 – ‘Grade A’ Student Master Fellowship from Rio de Janeiro Research Council.

2005 – 2006 – Master Fellowship. National Council for Scientific and Technological Development, CNPq.

2003 – 2004 – Undergraduate student fellowship, National Council for Scientific and Technological Development, CNPq.

Supervision

Nadia Sinner (2012-2013) – *Hilfswissenschaftler*, HiWi.

Languages

Portuguese (native)

English (fluent)

German (advanced)

15. Appendix

Appendix 1: Pruned alignment of kinase domain of LjSYMRK (amino acid positions 593-870) and homologous stretches of related MLD-LRR-RLK sequences [LjSYMRK, LjShRK1, LjShRK2, ShRK1(AT1G67720), ShRK2(AT2G37050), AT5G48740, AT1G51790, IOS1(AT1g51800), AT1G51910, AT1G51890, AT1G51860, AT1G51880, AT1G07550, AT2G14440, AT2G14510, AT3G46350, AT3G46340, AT3G46370, AT3G46400, AT3G46330, AT5G59670, AT5G59680, AT5G59650, AT5G16900, AT1G07560, AT4G20450, AT2G28960, AT2G29000, AT2G28970, AT2G28990, AT1G491000, AT1G51810, AT1G51805, AT1G51830, AT1G51820, AT1G51850, AT2G04300, AT3G21340, AT1G05700, AT2G19210, AT2G19230, AT2G19190, AT4G29990, AT4G29180].

```

LjSYMRK/1-278      1 ERYKTLIGEGGFGSVYRGTLND--GQEVAVKVRSATST--QGTREFDNELNLLSAIQHENLV
LjShRK1/1-278     1 NNFEEKKIGSGGFGVYVYKGLKD--GKEIAVKVLTNSY--QGKREFSNEVALLSRIHHRNLV
LjShRK2/1-272     1 RNLERIVGKGGFGIVYHGCVDG--IEVAVKMLS--PSA--QGYLQFQAEAKFLAKVHHKCLT
ShRK1/1-276       1 DNFSKVGRGSGVYVYGRMKD--GKEVAVKITADPSS--HLNRQFVTEVALLSRIHHRNLV
ShRK2/1-279       1 KKFEEKRIGSGGFGIVYVYKGTRE--GKEIAVKVLANNSY--QGKREFANEVLLSRIHHRNLV
AT5G48740/1-277   1 RNFKKEVIGRGSFGAVYRGLKLPD--GKQVAVKVRFDRTQ--LGADSFINEVHLLSQIRHQNLV
AT1G51790/1-273   1 NGFDRDQKGVGFRNYLGLKLDG--KEVTVKLVSSLSS--QGYKQLRAEVKHLFRIHKNLI
IOS1/1-274        1 NNFERVLGRGGFGVYVYGVLNN--EPVAVKMLTSTA--LGYKQFKAEVALLLRVHHKDLT
AT1G51910/1-272   1 NNFERVLGKGGYGRVYVYKGLDD--TEVAVKMLFSSAEQDYKHFKAEVALLLRVHHRHLV
AT1G51890/1-274   1 KNFERVLGKGGFGTVYHGNLDD--TQVAVKMLSHSSA--QGYKEFKAEVALLLRVHHRHLV
AT1G51860/1-274   1 NNFERVLGKGGFGTVYHGNLDG--AEVAVKMLSHSSA--QGYKEFKAEVALLLRVHHRHLV
AT1G51880/1-274   1 NNFERVLGKGGFGTVYHGNLDD--TQVAVKMLSHSSA--QGYKEFKAEVALLLRVHHRNLV
AT1G07550/1-273   1 NNFQVVIKGGGFGVYVYQCLN--NEQAAIKVLSHSSA--QGYKEFKTEVALLLRVHHEKLV
AT2G14440/1-274   1 NNFVVLGKGGFGVYHGFNLN--NEQVAVKVLSSST--QGYKEFKTEVALLLRVHVNVLV
AT2G14510/1-274   1 NNFVVLGKGGFGVYHGFNLN--NEQVAVKVLSSST--QGYKEFKTEVALLLRVHVNVLV
AT3G46350/1-275   1 NNFQRALGEGGFGTVYHGDLDL--SQVAVKLLSSST--QGYKEFKAEVALLLRVHHINLV
AT3G46340/1-276   1 KNLQRPLGEGGFGVYHGDINGSSSQVAVKLLSSST--QGYKEFKAEVALLLRVHHINLV
AT3G46370/1-275   1 KNFQKTIGEGGFGTVYHGNLNG--SEQVAVKVLSSSS--QGYKHFKAEVALLLRVHHINLV
AT3G46400/1-275   1 KKFEEKALGEGGFGIVYHGYLKN--VEQVAVKVLSSSS--QGYKHFKAEVALLLRVHHINLV
AT3G46330/1-276   1 KNLQRPLGEGGFGVYHGDNLG--SEQVAVKLLSQTSA--QGYKEFKAEVALLLRVHHINLV
AT5G59670/1-275   1 KNFQRVLGKGGFGMYYHGTVKG--SEQVAVKVLSSST--QGSKEFKAEVALLLRVHHTNLV
AT5G59680/1-275   1 NNFGRVVGEGGFGVVCVCHTVNG--SEQVAVKLLSSST--QGYKEFKAEVALLLRVHHTNLV
AT5G59650/1-275   1 NNFQRVVGEGGFGVVCVCHTING--SEQVAVKVLSSSS--QGYKHFKAEVALLLRVHHTNLV
AT5G16900/1-275   1 NNFERVIKGGGFGVYHGYLND--SEQVAVKVLSPSSS--QGYKEFKAEVALLLRVHHINLV
AT1G07560/1-272   1 KKFERVLGKGGFGMYYHGYING--TEEVAVKLLSPSSA--QGYKEFKTEVALLLRVYHTNLV
AT4G20450/1-275   1 NNFERPLGEGGFGVYHGNVND--NEQVAVKVLSESSA--QGYKQFKAEVALLLRVHHINLV
AT2G28960/1-275   1 DNFERVLGEGGFGVYHGIILNG--TQPIAVKLLSSSV--QGYKEFKAEVALLLRVHVNVLV
AT2G29000/1-275   1 NKFERVIKGGGFGIVYHGHVND--TEQVAVKLLSHSST--QGYKQFKAEVALLLRVHHTNLV
AT2G28970/1-275   1 NNFQRVLGEGGFGVYHGCVNG--TQVAVKLLSSSS--QGYKHFKAEVALLLRVHHKNLV
AT2G28990/1-275   1 NNFDKALGEGGFGVYHGFVNV--IEQVAVKLLSSSS--QGYKHFKAEVALLMRVHHINLV
AT1G49100/1-275   1 NNFQRIKGGGFGMYYHGYVNG--REQVAVKVLSSSS--QGYKQFKAEVALLLRVHHKNLV
AT1G51810/1-275   1 NNFQRILGKGGFGIVYHGFVNG--VEQVAVKILSHSSS--QGYKQFKAEVALLLRVHHKNLV
AT1G51805/1-275   1 NNFQRILGKGGFGIVYHGFVNG--VEQVAVKILSHSSS--QGYKQFKAEVALLLRVHHKNLV
AT1G51830/1-275   1 NNFQRVLGKGGFGIVYHGLVNG--TEQVAIKILSHSSS--QGYKQFKAEVALLLRVHHKNLV
AT1G51820/1-275   1 NNFQRILGKGGFGMYYHGFVNG--TEQVAVKILSHSSS--QGYKQFKAEVALLLRVHHKNLV
AT1G51850/1-275   1 NNFQRILGKGGFGMYYHGFVNG--TEQVAVKILSHSSS--QGYKEFKAEVALLLRVHHKNLV
AT2G04300/1-275   1 NNFERVLGKGGFGMYYHGTVND--AEQVAVKMLSPSSS--QGYKEFKAEVALLLRVHHKNLV
AT3G21340/1-275   1 NNFERVLGKGGFGMYYHGTVNN--TEQVAVKMLSHSSS--QGYKEFKAEVALLLRVHHKNLV
AT1G05700/1-275   1 NNFQVVLGKGGFGTVYHGFYDN--LQVAVKLLSETSA--QGFKEFRSEVFLVRVHVNLT
AT2G19210/1-276   1 NNFERVLGQGGFGKVVYHGVND--DQVAVKILSESSA--QGYKEFKAEVALLLRVHHKNLT
AT2G19230/1-275   1 NNFERVLGQGGFGKVVYGVLRG--EQVAIKMLSKSSA--QGYKEFKAEVALLLRVHHKNLI
AT2G19190/1-273   1 NNFERVIKGGGFGKVVYHGVING--EQVAVKVLSESSA--QGYKEFKAEVALLMRVHHTNLT
AT4G29990/1-273   1 NNFERVLGKGGFGKVVYHGFING--DQVAVKILSEEST--QGYKEFKAEVALLMRVHHTNLT
AT4G29180/1-277   1 NNFNKVIGKGGFGIVYVYGLSLED--GTEIAVKMISSSSSQVSKEFQVEAELLT VHHRNLA
AT4G29450/1-277   1 NNFNKVIGKGGFGIVYVYGLSLED--GTKIAVKMISSSSLSRASNQFQVEAELLT VHHRNLA

```

LjSYMRK/1-278 59 PLLGYCNESDQQLIVYPFMSNGSLQDRLYGEPAKRKILDWPTRLSIALGAARGLAYLHTF
 LjShRK1/1-278 59 QLLGYCREEGNSMLIYEFMHNGLKEHLYGPLTHGRSINWIKRLEIAEDSAKGIEYLLHTG
 LjShRK2/1-272 57 ALIGYCDDGTNMALIYEMANSDLAKHLSGKNE--NILGWNQRLQIAVDAAEGLEYLHHG
 ShRK1/1-276 59 PLIGYCEEADRRILVYEYMHNSLGDHLHGSSDY-KPLDWLTRLQIAQDAAGLEYLHTG
 ShRK2/1-279 59 QFLGYCQEEGKNMLVYEYFMHNGLKEHLYGVVPRDRRISWIKRLEIAEDAARGLEYLHTG
 AT5G48740/1-277 59 SFEGFCYEPKRQILVYEYLSGGSLADHLYGPRSKRHSLNWVSRLKVAVDAAGLDYLNHG
 AT1G51790/1-273 58 TMLGYCNEGDKMAVIYEYMANGNLKQHIS-ENST-TVFSWEDRLGIAVDVAQGLEYLLHTG
 IOS1/1-274 58 CLVGYCEEGDKMSLIYEFMANGDLKEHLSGKRGP-SILTWEGRLRIAAESAQGLEYLHNG
 AT1G51910/1-272 59 GLVGYCDDGNFALIYEYMANGDLKENMSGNRSG-HVLSWENRMQIAEEAAQGLEYLHNG
 AT1G51890/1-274 58 GLVGYCDDGNLALIYEYMEKGDLRENMSGKHSV-NVLSWETRMQIAVEEAAQGLEYLHNG
 AT1G51860/1-274 58 GLVGYCDDGNLALIYEYMANGDLRENMSGKRG-NVLTWENRMQIAVEEAAQGLEYLHNG
 AT1G51880/1-274 58 GLVGYCDDGNLALIYEYMANGDLKENMSGKRG-NVLTWENRMQIAVEEAAQGLEYLHNG
 AT1G07550/1-273 58 SLIGYCDDNGLALIYEYMGKNLKEHLSGKPGC-SVLSWPIRLKIALESAIGIEYLLHTG
 AT2G14440/1-274 58 SLVGYCDKGNDALIYEYFMENGLKEHLSGKRGG-PVLNWPGRLKIAIESALGIEYLLHIG
 SLVGYCDEGIDLALIYEYMENGLKEHLSGKRGG-SVLNWSRLKIAIESALGIEYLLHIG
 AT3G14510/1-274 58 SLVGYCDERDLLALIYEYMSNGLKHHLSGKHGG-SVLKNTRLQIAVDAALGLEYLLHIG
 AT3G46350/1-275 59 NLVGYCDERDLLALIYEYMSNGLKHHLSGEHGG-SVLSNIRLRIAIVDAALGLEYLLHIG
 AT3G46340/1-276 60 SLVGYCDERDLLALIYEYMSNGLKHHLSGKHGG-SVLKNTRLQIAVDAALGLEYLLHIG
 AT3G46370/1-275 59 SLVGYCDERNHLALIYEYCMSNGDLKDHLSGKGN-AVLKWSTRLRIAIVDAALGLEYLLHYG
 AT3G46400/1-275 59 SLVGYCDEKDHLALIYEYMANGDLKDHLSGKQGD-SVLEWTTRLQIAVEEAAQGLEYLLHYG
 AT3G46330/1-276 59 NLVGYCDEQDHFALIYEYMSNGLHQHLSGKHGG-SVLNWGTRLQIAVEEAAQGLEYLLHTG
 AT5G59670/1-275 59 SLVGYCCEGDYLALVYEFLPNGDLKQHLSGKGN-SIINWSIRLRIALEEAAQGLEYLLHIG
 AT5G59680/1-275 59 SLVGYCDEGDHLALIYEYFVPNGDLRQHLSGKGG-PIVNWGTRLRIAEEAAQGLEYLLHIG
 AT5G59650/1-275 59 SLVGYCDERDLLALIYEYFLPKGDLRQHLSGKSGG-SFINWGNRLRIALEEAAQGLEYLLHSG
 AT5G16900/1-275 59 SLVGYCDEQAHLALIYEYMANGDLKSHLSGKHGD-CVLKWENRLSIAVEEAAQGLEYLLHSG
 AT1G07560/1-272 59 SLVGYCDEKDHLALIYQYMVNGDLKKHFSGS---SIISWDRLNIAVDAASGLEYLLHIG
 AT4G20450/1-275 59 TLVGYCDEGQHLVLIEYMSNGLKQHLSGENSR-SPLSWENRLRIAEEAAQGLEYLLHIG
 AT2G28960/1-275 59 SLVGYCDEESNLALLYEYAPNGDLKQHLSGERGG-SPLKWSSRLKIVVETAQGLEYLLHTG
 AT2G29000/1-275 59 NLVGYCNEEDHLALVYEYAANGDLKQHLSGESSS-AALNWASRLGIATETAQGLEYLLHIG
 AT2G28970/1-275 59 SLVGYCDEGDHLALIYEYMPNGLKQHLSGKGG-FVLSWESRLRVAVDAALGLEYLLHTG
 AT2G28990/1-275 59 SLVGYCDEGEHLALIYEYMPNGLKQHLSGKHGG-FVLSWESRLKIVLDAALGLEYLLHTG
 AT1G49100/1-275 59 SLVGYCEKGKELALVYEYMANGDLKEFFSGKRGD-DVLRWETRLQIAVEEAAQGLEYLLHKG
 AT1G51810/1-275 59 GLVGYCEEGDKLALIYEYMANGDLDEHMSGKRGG-SILNWGTRLKIALEEAAQGLEYLLHNG
 AT1G51805/1-275 59 GLVGYCDEGENMALIYEYMANGDLKEHMSGTRNR-FILNWETRLKIVIDSAQGLEYLLHNG
 AT1G51830/1-275 59 GLVGYCDEGENLALIYEYMANGDLKEHMSGTRNR-FILNWGTRLKIVESAQGLEYLLHNG
 AT1G51820/1-275 59 GLVGYCDEGDNLALIYEYMANGDLKEHMSGTRNR-FILNWGTRLKIVIESAQGLEYLLHNG
 AT1G51850/1-275 59 GLVGYCDEGENMALIYEYMANGDLKEHMSGTRNR-FTLNWGTRLKIVESAQGLEYLLHNG
 AT2G04300/1-275 59 GLVGYCDEGENSLIYEYMAKGDLKEHMLGNQGV-SILDWKTRLKIVAESAQGLEYLLHNG
 AT3G21340/1-275 59 GLVGYCDEGENLALIYEYMANGDLREHMSGKRGG-SILNWETRLKIVESAQGLEYLLHNG
 AT1G05700/1-275 58 ALIGYFHEGDQMLIYEYFMANGNMADHLAGKYQ--HTLSWRQRLQIALDAAQGLEYLLHCG
 AT2G19210/1-276 58 ALIGYCHEGKMALIYEYFMANGTLGDYLSGEKS--YVLSWEERLQISLDAAQGLEYLLHNG
 AT2G19230/1-275 58 ALIGYCHEGDQMLIYEYIGNTLGDYLSGKNS--SILSWEERLQISLDAAQGLEYLLHNG
 AT2G19190/1-273 58 SLVGYCNEINHMVLIEYMANENLGDYLAGKRS--FILSWEERLKISLDAAQGLEYLLHNG
 AT4G29990/1-273 58 SLIGYCNEDNHMALIYEYMANGNLGDYLSGKSS--LILSWEERLQISLDAAQGLEYLLHYG
 AT4G29180/1-277 60 SFVGYCDDGRSMALIYEYMANGNLQDYLSSENAE-D-LSWEKRLHIAIDSAQGLEYLLHHG
 AT4G29450/1-277 60 SFVGYCDDDRSMALIYEYMANGNLQAYLSSENAE-D-LSWEKRLHIAIDSAQGLEYLLHDG

LjSYMRK/1-278 119 PGRSVIHRDIKSSNILLDHSMCAKVADFGFSKYAPQEG-DSYVSLEVRGTAGYLDPEYYK
 LjShRK1/1-278 119 CVPAVIHRDLKSSNILLDRQLRAKVSDFGLSK-LAVDG-VSHVSSIVRGTVGYLDPEYYI
 LjShRK2/1-272 115 SNPPIVHRDVKSKNILLNEKFQAKLADFGLSKIFPNEG-DTHVYTVVAGTPGYLDPEYNR
 ShRK1/1-276 118 CNPSTIHRDVKSSNILLDINMRAKVSDFGLSR-QTEED-LTHVSSVAKGTVGYLDPEYYA
 ShRK2/1-279 119 CVPAPIHRDLKTSNILLDKHMRKVSDFGLSK-FAVDG-TSHVSSIVRGTVGYLDPEYYI
 AT5G48740/1-277 119 SEPRIIHRDVKSSNILLDKDMNAKVSDFGLSKQFTKAD-ASHITTVVKGTAGYLDPEYYI
 AT1G51790/1-273 116 CKPPIIHRNVKCTNVFLDESFNKLGFGLSRAFDAAE-GSHLNTAIAGTPGYLDPEYYT
 IOS1/1-274 117 CKPQIVHRDIKTTNILLNEKFQAKLADFGLSRSFPLGT-ETHVSTIVAGTPGYLDPEYYR
 AT1G51910/1-272 118 SRPPMVHRDVKTTNILLNELYQAKLADFGLSRSFVVDG-ESYVSTIVAGTPGYLDPE---
 AT1G51890/1-274 117 CRPPMVHRDVKPTNILLNERSQAKLADFGLSRSFVVDG-ESHVMTVVAGTPGYLDPEYYR
 AT1G51860/1-274 117 CRPPMVHRDVKTTNILLNERCGAKLADFGLSRSFPIDG-ECHVSTVVAGTPGYLDPEYYR
 AT1G51880/1-274 117 CTPPMVHRDVKTTNILLNERYGAKLADFGLSRSFVVDG-ESHVSTVVAGTPGYLDPEYYR
 AT1G07550/1-273 117 CKPKIVHRDVKSTNILLSEEFKAKIADFGLSRSFLIGN-EAQ-PTVVAGTFGYLDPEYHK
 AT2G14440/1-274 117 CKPPMVHRDVKSTNILLGLRFEAKLADFGLSRSFLVGS-QTHVSTNVAGTLGYLDPEYYQ
 AT2G14510/1-274 117 CQPPMVHRDVKSTNILLGLRFEAKLADFGLSRSFLVGS-QAHVSTNVAGTLGYLDPEYYR
 AT3G46350/1-275 118 CRPPSMVHRDVKSTNILLDNFMAKIAADFGLSRSFILGG-ESHVSTVVAGSLGYLDPEYYR
 AT3G46340/1-276 119 CRPSMVHRDVKSTNILLDQFTAKMADFGLSRSFQLGD-ESQVSTVVAGTPGYLDPEYYR
 AT3G46370/1-275 118 CRPSIVHRDVKSTNILLDQQLMAKIAADFGLSRSFKLGE-ESQASTVVAGTLGYLDPEYYR
 AT3G46400/1-275 118 CRPSMVHRDVKSTNILLDQFMAKIAADFGLSRSFKVGD-ESEISTVVAGTLGYLDPEYYL
 AT3G46330/1-276 118 CKPAMVHRDVKSTNILLDEFFKAKIADFGLSRSFQVGGQSQVSTVVAGTLGYLDPEYYL
 AT5G59670/1-275 118 CTPPMVHRDVKTANILLDENFKAKLADFGLSRSFQEG-ESQESTTIAGTLGYLDPECYH
 AT5G59680/1-275 118 CTPPMVHRDVKTTNILLDEHYKAKLADFGLSRSFPVGG-ESHVSTVIAGTPGYLDPEYYH
 AT5G59650/1-275 118 CTPIVHRDIKTTNILLDEQLKAKLADFGLSRSFPIGG-ETHISTVVAGTPGYLDPEYYQ
 AT5G16900/1-275 118 CKPLMVHRDVKSMNILLDEHFQAKLADFGLSRSFSVGE-ESHVSTGVVGTGGYLDPEYYR
 AT1G07560/1-272 115 CKPLIVHRDVKSSNILLDQLQAKLADFGLSRSFPIGD-ESHVSTLVAGTFGYLDHEYYQ
 AT4G20450/1-275 118 CKPPMIHRDIKSMNILLDNFQAKLGFGLSRSFPVGS-ETHVSTNVAGSPGYLDPEYYR
 AT2G28960/1-275 118 CKPPMVHRDVKTTNILLDEHFQAKLADFGLSRSFPVGG-ETHVSTAVAGTPGYLDPEYYR
 AT2G29000/1-275 118 CEPPMIHRDVKTTNILLDEHFHAKLADFGLSRSFPVGV-ESHVSTNVAGTPGYLDPEYYR
 AT2G28970/1-275 118 CKPPMVHRDIKSTNILLDERFQAKLADFGLSRSFPFTEEN-ETHVSTVVAGTPGYLDPEYYQ
 AT2G28990/1-275 118 CVPPMVHRDIKTTNILLDQHLQAKLADFGLSRSFPIGN-EKNVSTVVAGTPGYLDPEYYQ
 AT1G49100/1-275 118 CRPPIVHRDVKTANILLDEHFQAKLADFGLSRSFLNEG-ESHVSTVVAGTIGYLDPEYYR
 AT1G51810/1-275 118 CKPLMVHRDVKTTNILLNEHFDTKLADFGLSRSFPIEG-ETHVSTVVAGTIGYLDPEYYR
 AT1G51805/1-275 118 CKPLMVHRDVKTTNILLNEHFQAKLADFGLSRSFPIEG-ETHVSTVVAGTPGYLDPEYYK
 AT1G51830/1-275 118 CKPLMVHRDIKTTNILLNEQFADKADFGLSRSFPIEG-ETHVSTAVAGTPGYLDPEYYR
 AT1G51820/1-275 118 CKPPMVHRDVKTTNILLNEHFQAKLADFGLSRSFLIEG-ETHVSTVVAGTPGYLDPEYHR
 AT1G51850/1-275 118 CKPPMVHRDVKTTNILLNEHFQAKLADFGLSRSFPIEG-ETHVSTVVAGTPGYLDPEYYK
 AT2G04300/1-275 118 CKPPMVHRDVKTTNILLDEHFQAKLADFGLSRSFPLEG-ETRVDTVVAGTPGYLDPEYYR
 AT3G21340/1-275 118 CKPPMVHRDVKTTNILLNEHLHAKLADFGLSRSFPIEG-ETHVSTVVAGTPGYLDPEYYR
 AT1G05700/1-275 116 CKPPIVHRDVKTSNILLNEKNRAKADFGLSRSFHTES-RSHVSTLVAGTPGYLDPLCFE
 AT2G19210/1-276 116 CKPPIVQRDVKPANILINNEKLQAKIADFGLSRSVALDG-NNQDITAVAGTIGYLDPEYHL
 AT2G19230/1-275 116 CKPPIVHRDVKPTNILINNEKLQAKIADFGLSRSFTLEG-DSQVSTEVAGTIGYLDPEHYS
 AT2G19190/1-273 116 CKPPIVHRDVKPTNILLNEKLQAKMADFGLSRSFSVEG-SGQISTVVAGSIGYLDPEYYI
 AT4G29990/1-273 116 CKPPIVHRDVKPTNILLNENLQAKIADFGLSRSFPVEG-SSQVSTVVAGTIGYLDPEYYA
 AT4G29180/1-277 118 CRPPIVHRDVKTANILLNDNLEAKIADFGLSKVPEDD-LSHVVTAVMGTGGYLDPEYYN
 AT4G29450/1-277 118 CRPAIVHRDVKTANILINDNLEAKIADFGLSKVPEDD-LSHVVTVMGTGGYLDPEYYR

LjSYM RK/1-278 178 TQQLSEKSDVFSFGVVLLEIVSGREPLNIKR-PRTEWSLVEWATPYI-RGSKVDEIVDPG
 LjShRK1/1-278 177 SQQLTDKSDIYSFGVILLELISGQEAISNDSFGANCRNIVQWAKLHI-ESGDIQGIIDPA
 LjShRK2/1-272 174 SSRLNEKSDVFSFGVVLLELITGQPAVTKTE---DKIHIIQWVSSL-LQREVKDIVDP
 ShRK1/1-276 176 SQQLTEKSDVYSFGVVLLELISGQEAISNESFGVNCRNIVQWAKMHI-DNGDIRGIIDPA
 ShRK2/1-279 177 SQQLTEKSDVYSFGVILLELISGQEAISNESFGVNCRNIVQWAKMHI-DNGDIRGIIDPA
 AT5G48740/1-277 178 TLQLTEKSDVYSFGVVLLELISGQEAISNESFGVNCRNIVQWAKMHI-DNGDIRGIIDPA
 AT1G51790/1-273 175 SNMLTEKSDVYSFGVVLLEIVTAKPAIKNE---ERMHISQWVESLL-SRENIIVEILDPS
 IOS1/1-274 176 TNWLTEKSDVFSFGVVLLELVTNQPVIDMKR---EKSHIAEWVGLML-SRGDINSIVDPK
 AT1G51910/1-272 174 TNLLSEKTDVYSFGVVLLEIITNQPVIDTR---EKAHITDWVGFKL-MEGDIRNIIDPK
 AT1G51890/1-274 176 TNWLSEKSDVYSFGVVLLEIIVTNQPVIMKNR---ERPHINEWVFMML-TNGDIKSIVDPK
 AT1G51860/1-274 176 TNWLSEKSDVYSFGVVLLEIIVTNQPVIDKTR---ERPHINDWVGFML-TKGDIKSIVDPK
 AT1G51880/1-274 176 TNWLSEKSDVYSFGVVLLEIIVTNQPVTDKTR---ERTHINEWVGSML-TKGDIKSILDPK
 AT1G07550/1-273 175 TSLLSMKSDVYSFGVVLLEIISGQDVIDLSR---ENCNIVEWTSFIL-ENGDIESIVDPN
 AT2G14440/1-274 176 KNWLTEKSDVYSFGIVLLEIITGQPVIEQSR---DKSYIVEWAKSML-ANGDIESIMDRN
 AT2G14510/1-274 176 KNWLTEKSDVYSFGIVLLESITGQPVIEQSR---DKSYIVEWAKSML-ANGDIESIMDPN
 AT3G46350/1-275 177 TSRLAEMSDVYSFGIVLLEIITNQRVIDKTR---EKPHITEWTFMML-NRGDIRTRIMDPN
 AT3G46340/1-276 178 TGRLAEMSDVYSFGIVLLEIITNQRVIDPAR---EKSHITEWTFMML-NRGDIRTRIMDPN
 AT3G46370/1-275 177 TCRLAEMSDVYSFGIVLLEIITNQNVIDHAR---EKAHITEWVGLVL-KGGDVTRIVDPN
 AT3G46400/1-275 177 TSRLAEMSDVYSFGIVLLEIITNQRVDFDQAR---EKHITTEWVAFMML-NRGDIRTRIVDPN
 AT3G46330/1-276 178 TSELSEKSDVYSFGIVLLEIITNQRVIDQTR---ENPNIAEWVTFVI-KKGDTSQIVDPK
 AT5G59670/1-275 177 SGRLGEKSDVYSFGIVLLEMITNQPVINQTS---GDSHITQWVGFQM-NRGDILEIMDPN
 AT5G59680/1-275 177 TSRLSEKSDVYSFGIVLLEMITNQAVIDRNR---RKSHTQWVGFSEL-NGGDIKIMDLK
 AT5G59650/1-275 177 TTRLGEKSDVYSFGIVLLEIITNQPVIDQSR---SKSHISQWVGFEL-TRGDIKIMDPN
 AT5G16900/1-275 177 TYRLTEKSDVYSFGIVLLEIITNQPVLEQAN---ENRHIAERVMTML-TRSDISTIVDPN
 AT1G07560/1-272 174 TNRLSEKSDVYSFGVVLLEIITNKPVIDHNR---DMPHIAEWVKLML-TRGDISNIMDPK
 AT4G20450/1-275 177 TNWLTEKSDVFSFGVVLLEIITSQPVIDQTR---EKSHIGEWVGFKL-TNGDIRKNIVDP
 AT2G28960/1-275 177 TNRLNEKSDVYSFGIVLLEIITSRPVIQQTR---EKPHIAAWVGYML-TKGDIEENVVDP
 AT2G29000/1-275 177 TNWLTEKSDVYSFGIVLLEIITNQPVIQQVR---EKPHIAEWVGLML-TKGDIKSIMDPK
 AT2G28970/1-275 177 TNWLTEKSDVYSFGIVLLEIITNRPVIQQSR---EKPHILEWVGFIV-RTGDIGNIVDPN
 AT2G28990/1-275 177 TNWLTEKSDIYSFGIVLLEIITNRPVIQQSR---EKPHIVEWVGFMI-TKGDILRSIMDPN
 AT1G49100/1-275 177 TNWLTEKSDVYSFGVVLLEIITNQRVIERTR---EKPHIAEWNLMML-TKGDIRKIVDPN
 AT1G51810/1-275 177 TNWLTEKSDVYSFGVVLLEIITNQPVIDQNR---EKRHIAEWVGGML-TKGDIKSITDPN
 AT1G51805/1-275 177 TNRLTEKSDVYSFGIVLLEMITNRPVIDQSR---EKPYISEWVGIML-TKGDIIISIMDPS
 AT1G51830/1-275 177 TNWLTEKSDVYSFGVVLLEIITNQPVIDPRR---EKPHIAEWVGEVL-TKGDIIKNIMDPS
 AT1G51820/1-275 177 TNWLTEKSDVYSFGIVLLEIITNRHVIDQSR---EKPHIGEWVGVML-TKGDIIQSIMDPS
 AT1G51850/1-275 177 TNWLTEKSDVYSFGIVLLEIITNRPVIDKSR---EKPHIAEWVGVML-TKGDINSIMDPN
 AT2G04300/1-275 177 TNWLNEKSDVYSFGIVLLEIITNQHVINQSR---EKPHIAEWVGVML-TKGDIKSIVDPK
 AT3G21340/1-275 177 TNWLNEKSDVYSFGIVLLEIITNQLVINQSR---EKPHIAEWVGLML-TKGDIIQIMDPK
 AT1G05700/1-275 175 TNGLEKSDIYSFGVVLLEMITGKTVIKESQ--TKRVHVSDWVVISILRS-TNDVNNVIDSK
 AT2G19210/1-276 175 TQKLSEKSDIYSFGVVLLEIVVSGQPVARSRTAENIHITDRVDLML-STGDIRGIVDPK
 AT2G19230/1-275 175 MQQFSEKSDVYSFGVVLLEIVITGQPVISRSR-TEENRHISDRVSLML-SKGDIIKSIVDPK
 AT2G19190/1-273 175 TRQMNEKSDVYSFGVVLLEIVITGQPAIASSK--TEKVHISDHVRSIL-ANGDIRGIVDQR
 AT4G29990/1-273 175 TRQMNEKSDVYSFGVVLLEIVITGKPAIWHSR--TESVHLSQVGSML-ANGDIKIGIVDQR
 AT4G29180/1-277 177 TFKLNEKSDVYSFGIVLLELITGKRSMKTD-DGKMNVVHYVEPFL-KMGDIDGVVDP
 AT4G29450/1-277 177 TFVLEKSDVYSFGVVLLELITGQRAIKTE-EGDNISVIHYVWVFFF-EARELDGVVDP

LjSYMRK/1-278 236 IK-GGYHAEAMWRVVEVALQCLEPFSTYRPSMVAIVRELEDAI
 LjShRK1/1-278 236 LG-NDYDLQSMWKIAEKALMCVQPHGMRPSISEVLKEIQDAIA
 LjShRK2/1-272 230 LQ-GEFDIDSAKKALDTAMTCVAPTSINRPTMSHVVMELKLCPL
 ShRK1/1-276 234 IA-SNVKIESVWRVAEVANQCVQEQRGHNRPRMQEVIVAIQDAIR
 ShRK2/1-279 236 LAEDDYSLQSMWKIAEKALLCVKPHGNMRPSMSQVQDIQDAIR
 AT5G48740/1-277 235 LK-ETFDPASMKKAASIAIRCVGRDASGRPSIAEVLTKLKEAYS
 AT1G51790/1-273 231 LC-GDYDPNSAFKTVEIAVACVCRNSGDRPGMSQVVTALKESLA
 IOS1/1-274 232 LQ-GDFDPNTIWKVVEVETAMTCLNPSSRRPTMTQVVM DLKECLN
 AT1G51910/1-272 230 LI-KEFDTNGVWKAVELALSCVNPTSNHRPTMPHVVMELKECLD
 AT1G51890/1-274 232 LN-EDYDTNGVWKVVELALACVNPSSRRPTMPHVVMELNECLA
 AT1G51860/1-274 232 LM-GDYDTNGAWKIVELALACVNPSSNRPTMAHVVMELNDCVA
 AT1G51880/1-274 232 LM-GDYDTNGAWKIVELALACVNPSSNRPTMAHVVELNECVA
 AT1G07550/1-273 231 LH-QDYDTSSAWKVELAMSCVNRTSKERNMSQVHVHNECLE
 AT2G14440/1-274 232 LH-QDYDTSSWKALELAMLCINPSSTLRPNMTRVAHELNECLE
 AT1G14510/1-274 232 LH-QDYDSSSSWKALELAMLCINPSSQRPNMTRVAHELNECLE
 AT3G46350/1-275 233 LN-GDYNSHSVWRALELAMSCANPSSENRPSMSQVVAELKECLI
 AT3G46340/1-276 234 LQ-GDYNSRSVWRALELAMSCANPSSSEKRPMSQVVI ELKECIR
 AT3G46370/1-275 233 LD-GEYNSRSVWRALELAMSCANPSSSEHRPIMSQVVIDLKECLN
 AT3G46400/1-275 233 LH-GEYNSRSVWRALELAMSCANPSSSEYRPNMSQVVI ELKECLT
 AT3G46330/1-276 234 LH-GNYDTHSVWRALEVAMSCANPSSVSRPNMSQVI INLKECLA
 AT5G59670/1-275 233 LR-KDYNINSAWRALELAMSCAYPSSSKRPMSQVIHELKECIA
 AT5G59680/1-275 233 LN-GDYDSRSARWALELAMSCADPTSARRPTMSHVVI ELKECLV
 AT5G59650/1-275 233 LN-GDYERSVWRVLELAMSCANPSSVNRPNMSQVANELKECLV
 AT5G16900/1-275 233 LI-GEYDSGSVRKALKLAMSCVDPSPVARPDMSHVVQELKQCIK
 AT1G07560/1-272 230 LQ-GVYDSGSAWKALELAMTVCVNPSSSLKRPNMSSHVHELKECLV
 AT4G20450/1-275 233 MN-GDYDSSLWKALELAMSCVSPSSSGRPNMSQVANELQECLL
 AT2G28960/1-275 233 LN-RDYEP TSVWKALEIAMSCVNPSSSEKRP TMSQVTNELKQCLT
 AT2G29000/1-275 233 LN-GEYDSSVWKALELAMSCVNPSSGGRPTMSQVISELKECLI
 AT2G28970/1-275 233 LH-GAYDVGSVWKAIELAMSCVNISSARRPMSQVVS DLKECVI
 AT2G28990/1-275 233 LH-QDYDIGSVWKAIELAMSCVSLSSARRPNMSRVVNELKECLI
 AT1G49100/1-275 233 LK-GDYHSDSVWKFVVELAMTCVNDSSATRPMTQVVTETELTECVT
 AT1G51810/1-275 233 LL-GDYNSGSVWKAVELAMSCMNPSMTRPTMSQVVFELKECLA
 AT1G51805/1-275 233 LN-GDYDSGSVWKAVELAMSCVNPSSSTRPTMSQVLI ALNECLV
 AT1G51830/1-275 233 LN-GDYDSTSVWKAVELAMCCLNPSSARRPNMSQVVI ELNECLT
 AT1G51820/1-275 233 LN-EDYDSGSVWKAVELAMSCVNPSSARRPTMSQVVI ELNECLA
 AT1G51850/1-275 233 LN-EDYDSGSVWKAVELAMSCVNPSSARRPTMSQVVI ELNECIA
 AT2G04300/1-275 233 FS-GDYDAGSVWRAVELAMSCVNPSSSTGRPTMSQVVI ELNECLA
 AT3G21340/1-275 233 LY-GDYDSGSVWRAVELAMSCVNPSSARRPTMSQVVI ELNECLS
 AT1G05700/1-275 233 MA-KDFDVNSVWKVVELALSSVSNVSDRPNMPHIVRGLNECLQ
 AT2G19210/1-276 234 LG-ERFDAGSAWKITEVAMACASSSSKNRPTMSHVVAELKESVS
 AT2G19230/1-275 233 LG-ERFNAGLAWKITEVALACASESTKTRLTMSQVVAELKESLC
 AT2G19190/1-273 232 LR-ERYDVGSAAWKMEIALACTEHTSAQRPTMSQVVMELKQIV-
 AT4G29990/1-273 232 LG-DRFEVGSAAWKITELALACASESSEQRPTMSQVVMELKQSI-
 AT4G29180/1-277 235 LH-GDFSSNSAAWKFEVAMSCVDRDRGTNRPNNTNQIVSDLKQCLA
 AT4G29450/1-277 235 LR-GDFSQDSAAWKFVDVAMSCVDRDKGSNRPTMNQIVAEKQCLA

Appendix 2: Pruned alignment of extracytoplasmatic region of LjSYMRK (amino acid positions 30-517) and homologous stretches of related MLD-LRR-RLKs (sequences as above; conserved GPC motifs are at alignment positions 464-467).

```

LjSYMRK/1-509      1 --MMELPATRILLSQAVTCFLCLYIFIGSASATEGFESIACCAD----LNYTDPLTTLNYYTT
LjShRK1/1-522     1 PFFLSLTLTLLLR-----S--SA-QMK-GFVSLDCGGK----ENFTD-EIGLQWTF
LjShRK2/1-506     1 --MAGL-LLLLQVQLSWTLPPII--VHA-QDQSGFISIDCGLEDE--PSYDETTSIHYTS
ShRK1_(AT1/1-533  1 -----MGLCLAQLAVTCLFLVFPVLS-QVT-EFVSDICGCS----SNYDPRGTGLGWVS
ShRK2_(AT2/1-529  1 --MVRISLTLCLLVSTCLFTSS--SA-QAP-GFVSLDCGGA----EPFTD-ELGLKWSF
AT5G48740/1-492   1 -----MLFWVLLSSFCVFCF----SSPDGFLSLSCGG----SSYT-AAYNISWVS
AT1G51860/1-511   1 --MMSKAKALTF--ICCVALLNLAIA-QDQSGFISIDCGLPEN--PSYTEKSTDIKYVS
IOS1(AT1G5/1-513  1 --MAFSSCFLLVLLQIFSAALLCL--A-QDQSGFISLDCGSPRE--TSFREKTNITYIS
AT1G51910/1-508   1 --MKTMMNGFLLS--TIAFAVFLVQA-QSQSGFISLDCGLIPKD-TTYTEQITNITYIS
AT1G51890/1-499   1 --MR----FLSFL--IFVFAVLGLVQA-QDQSGFISLDCGLVPE-ITYVEKSTNITYRS
AT1G51880/1-511   1 --MKSILHGFLLFL--ITAYVILSVQA-QDQLGFISLDCGLVPKN-ATYTEKTNITYKS
AT1G07550/1-504   1 --MDTCTRLIFA-A-CATLSILHLVQS-QNQGGFISLDCGLASNE-SPYNEANSNLTYIS
AT2G14440/1-502   1 --METRSKMLL-A-CATFSIISLVKS-QNQGGFISLYCGLPSNE-SPYIEPLTNLTYIS
AT2G14510/1-508   1 --METRNKFMLL-A-CATFSIMSLVKS-QNQGGFISLDCGLPSKE-SYIEPSNLTFIS
AT3G46350/1-491   1 --MNSSHELLLTAL-IATFAIFHLVQA-QEQEGFISLDCGLAPTEPSPYTEPVTTLQYSS
AT3G46340/1-513   1 --MEFPHSVLLVLLIATFAISNLVQAEEDEQGFISLDCGLPPNEVSPYIEPTGLRFSS
AT3G46370/1-427   1 -----
AT3G46400/1-508   1 --MESSHRFLLVALTVA---SSIIHLVQAQAGFISLDCGLSPNEQSPYVELETGLQFLS
AT3G46330/1-516   1 --MKNLCWVFLSLFWFGVFLIRFAEG-QNQEFGFISLDCGLPLNEP-PYIESETGIQFSS
AT5G59670/1-499   1 --MESSFGLLLAL--L-TLTIHIVQA-QDPQGFISLDCGLPANETSPYTETQGLLFSS
AT5G59680/1-508   1 --MERSLELLLLL--IRTLAIIHISQA-QSQGGFISLDCGLPANEPSPYTEPRTGLQFSS
AT2G59650/1-509   1 --MDSPCWLLLLL--LGAFAIIGCVQA-QDQEGFISLDCGLPMTPESSYTESVTLRFSS
AT5G16900/1-505   1 --MEDRHRYLFFI----FAIIHYVQA---QQGFISLDCGLPSNE-PPYIEPVTLVLFSS
AT1G07560/1-512   1 --MKNLRGLLLAFL-VLSLGISDFLRA-QDQGGFISLDCGLQADE-SPYTEPLTKLTFSS
AT4G20450/1-530   1 --MEGIHKLIFLAL-IWIFLITNIVDA-QDQGGFISLDCGMPRNE-SSYDESTGLNLFSS
AT2G28960/1-507   1 --MEGRRQRLLVFI-FGALAIHVLVQA-QPPDGFISLDCGLPVNE-SPYDPRTGLTFSS
AT2G29000/1-506   1 --MEGHRGLLLALI-VNIFSIIVHLVHA-QNPEGFISLDCGLPAKE-SPYTESTSLVFTS
AT2G28970/1-409   1 -----MMSHLLLAIGTFAVI--VGA-QKQEGFISLDCGFPIEE-SPYSDPSTGLTFSS
AT2G28990/1-506   1 -----MKIHLLLAMIGTFVVI--IGA-QDQEGFISLDCGLPSDE-SPYDPSFNGLTFSS
AT1G49100/1-518   1 MEKYFHGVLVCFVITVAF--IHVVQA-QDPNGFITLDCGLLPDG-SPYTNPSTGLTFSS
AT1G51810/1-384   1 -----
AT1G51805/1-504   1 MESH-----RVFVATFMLILHLVQA-QDQPGFINVDCGLLPD-SPYNALGTGLVYTS
AT1G51830/1-315   1 -----MTVFFINDC-----
AT1G51820/1-504   1 MERH-----FVFIATYLLIFHLVQA-QNQTGFISVDCGLSLE-SPYDAPQTGLTYTS
AT1G51850/1-486   1 MERH-----CVLVATFLLMLHIVHA-QDQIGFISVDCGLAPRE-SPYNEAKTGLTYTS
AT2G04300/1-480   1 MKTHPQAILLCVLEFFITF-GLLHVVEA-GNQEFGFISLDCGLSPNE-PPYVDAATDLTYTT
AT3G21340/1-520   1 MEYHPQAIRLCALIFISFYALLHLVQA-QDQKGFISLDCGLSPNE-PPYNDPSTGLTYST
AT1G05700/1-507   1 --MEEFRFLYLIYSAAFALCLVSVLA-QDQSGFISIDCGIPSG--SSYKDDTTGINYVS
AT2G19210/1-516   1 --MVHYNFSLIIFACFFAVFVLLVRA-QDQSGFVSDICGIPED--SSYNDETDIKYVS
AT2G19230/1-516   1 --MGNFNFPLVVSFASVVLV-LVCA-QDQSGFVSDICGIPED--SSYDEKTDIKYIS
AT2G19190/1-516   1 --MAMKLSLSSILFTSFALLFF-LVHA-QDQSGFISIDCGIPED--SSYNDETGIKYVS
AT4G29990/1-511   1 --MTRLRLLSWISITS----CVCLVFA-QDQSGFISIDCGIPDD--SSYDEKTNMKYVS
AT4G29180/1-509   1 -----MGAHSVFLILFVSIAIAIVVHG-QQAGFISIDCGSPPN--INYVDTDTGISYTW
AT4G29450/1-513   1 -----MRANLVFGI-FCALVTTILVHG-QDQSGYISIDCGIPPY--DTPEDTMTNINYS

```

LjSYM RK/1-509 56 DYTWFSDKRSC-----RKIPETELRNRSNENVRLFDIDEGKRCYNLPTIKNG--VYLIRG
 LjShRK1/1-522 44 DDKMS-YGEISTI-----SVTNETRKYMTLRHFADSRKYCYTLDDVVSRT--RYLLRT
 LjShRK2/1-506 53 DVNFDTGVSHSI---SPKYE-ASLERQFNNVRSFPP-GGRRNCYTLVVPOGRSKYLVIRA
 ShRK1_ (AT1/1-533 49 DSEI IKQKPVTL-----ANTNWNMQYRRRRDFPTDNKICYRLSTKERR--RYIVRT
 ShRK2_ (AT2/1-529 50 DNHLI-YGETANI-----SSVNETRTQYTTLRHFADSRKYCYTLNVTSRN--RYLIRA
 AT5G48740/1-492 42 DNDYIETGNTTTVTYAEGNSTSSV-----PIRLFPDPQGRQCYKLPVRKDLN--SVLIRA
 AT1G51790/1-518 55 DSSYTDGTSYFV---APENRQNM-KQSMWSVRSFPP-EGIRNCYTIAVNSST--KYLIRA
 IOS1 (AT1G5/1-513 54 DANFINTGVGSGI---KQGYR-TQFQQQTWNLRSFPP-QGIRNCYTLNLTIGD--EYLIRA
 AT1G51910/1-508 55 DADYIDSGLTERI---SDSYKSQL-QQQTWTLSRFP-EGQRNCYNFNLKANL--KYLIRG
 AT1G51890/1-499 51 DATYIDSGVPGKI---NEVYRTQF-QQQIWAALRSFPP-EGQRNCYNFSLTAKR--KYLIRG
 AT1G51860/1-511 55 DVYIDSGLVGKI---NDAYKTQF-QQQVWAVRSFPP-VGQRNCYNVNLTANN--KYLIRG
 AT1G51880/1-511 55 DANYIDSGLVGRI---SAEYKAQL-QQQTWTVRSFPP-EGERNNCYNFNLAKS--RYLIRA
 AT1G07550/1-504 55 DADFIQGGKTGNV---QKDLLMKL-RKPYTVLRYFPP-DGIRNCYSLNVKQDT--NYLIRV
 AT2G14440/1-502 55 DVNFVRGKGTGNI---KNNSDIDFTSRPYKVLRYFPP-EGIRNCYLSVVKQT--KYLIRT
 AT2G14510/1-508 54 DVNFIRGGKTGNI---QNSRNFIFKPFKVLRYFPP-DGIRNCYLSVKQGT--KYLIRT
 AT3G46350/1-491 57 DSNFIQSGKLGRI---DTSLQTFP-LKQQTTLRYFPP-DGIRNCYNLTVKQGT--NYLIRA
 AT3G46340/1-513 59 DSSFIQSGKIGKV---DKSFEATT-LKSYMTLRYFPP-DGKRCYNLIVKQGT--TYMIRA
 AT3G46370/1-427 1 -----MRNCYNLSVHKET--KYLIRV
 AT3G46400/1-508 55 DSSFIQSGKIGRI---DASLESKY-PRSQTTLRYFPP-DGIRNCYNVNVYKGT--NYLIRA
 AT3G46330/1-516 57 DENFIQSGKTGRI---PKNLESEN-LKQYATLRYFPP-DGIRNCYDLRVEEGR--NYLIRA
 AT5G59670/1-499 55 DATFIQSGKTGRV---QANQESKF-LKPYRTLRYFPP-EGVRNCYNLVFKER--KYLIAA
 AT5G59680/1-508 56 DAAFIQSGKIGRI---QANLEADF-LKPSTTMRYPFPP-DGKRCYNLNVKGR--NHLIRA
 AT5G59650/1-509 56 DAEFIQTGESGKI---QASMENDY-LKPYTRLRYFPP-EBERRNCYSLVDKNR--KYLIRA
 AT5G16900/1-505 50 DADHIPSGISGRI---QKNLEAVH-IPYLFRLRYFPP-DGLRNCYTLVDLQNR--RYMIKA
 AT1G07560/1-512 56 DADFIKSGKSGKI---QNVPGMEY-IPYTVLRYFPP-DGVRNCYTLVIVQGT--NYLIVA
 AT4G20450/1-530 56 DADFISGKSGTIKTEDSDSGVKY-IPYKQLRYFPP-EGARNCYNLTVMQGT--HYLIRA
 AT2G28960/1-507 56 DADFILSGLRGE-----AGDDNTYIYRQYKDLRYFPP-DGIRNCYNLKVEQGI--NYLIRA
 AT2G29000/1-506 56 DANFISGISTKL---PKHDD---YKPYNFLRYFPP-DGTRHCYDLVVKQGT--NYLIRA
 AT2G28970/1-409 51 DSTFIQTGESGRV---DKELNKIF-RKPYLTLRYFPP-EGKRCN-----
 AT2G28990/1-506 51 DSTFIQTGKIDSV---DKDLNINL-SKQYTLRYFPP-EGKRCYNLSDVVRGT--TYLIVV
 AT1G49100/1-518 56 DSSFIESGKNGRV---SKDSERNF-EKAFVTLRYFPP-DGERNCYNLNVTOGT--NYLIRA
 AT1G51810/1-384 1 -----
 AT1G51805/1-504 52 DVGLVSSGKTGKI---AKEFEENN-STPNLTLRYFPP-DGARNCYNLNVSRTD--NYMIKA
 AT1G51830/1-315 10 -----
 AT1G51820/1-504 52 DADLVASGKTGRL---AKEFEPIV-DKPTLTLRYFPP-EGVRNCYNLNVTSDT--NYLIKA
 AT1G51850/1-486 52 DDGLVNVGKPGRI---AKEFEPLA-DKPTLTLRYFPP-EGVRNCYNLNVTSDT--NYLIKA
 AT2G04300/1-480 58 DNDFVQSGKTGTI---DKELESTY-NKPILQLRYFPP-EGVRNCYTLNVTLGT--NYLIRA
 AT3G21340/1-520 59 DDGFVQSGKTGRI---QKAFESIF-SKPSLKLRYFPP-DGFRNCYTLNVTQDT--NYLIKA
 AT1G05700/1-507 56 DSSFVETGVSKSI---PFTAQ----RQLQNLRSFPP-EGSRNCYTLIPIQGGKGYLIRA
 AT2G19210/1-516 56 DAAFVESGTIHSI---DPEFQTSLEKQFQNVRSFPP-EGNRNCYDVKPPQGGKGYLIRT
 AT2G19230/1-516 55 DAAFVESGTIHSI---DSKFQKKNLEKQFQKVRSFPP-EGKKNCYDVQPPQGGKGYLIRT
 AT2G19190/1-516 55 DSADFVDSGTTKRI---AAQFQSSGFDRLHLLNRSFPP-QSKRSCYDVPTPRGKGYLIRT
 AT4G29990/1-511 52 DLGFVESGTSHSI---VSDLQTTSLERQFQNVRSFPP-EGKRCYDIPRQGGKGYLIRT
 AT4G29180/1-509 53 DAFFINAGVNLNVSEEGYGPKNPVLPPPLADVRSFPP-QGNRNCYTLTPSDGKGNLYLIRA
 AT4G29450/1-513 52 DEAFITTVGNFKVSEEGYGPKNPVLPLAEVRAFPP-QGNRNCYTLKLSQGDHLYLIRA

LjSYMRK/1-509 109 TFPFDLSLN-----SSFNASIGVTQLGAVR-----SSRLQDLEIVFRATKDYI
 LjShRK1/1-522 95 TFLYGNFD-SNNVYPKFDISVGATHWSTIV-----IS----DANTIEVRELIFWASSPTV
 LjShRK2/1-506 108 RFVYGNVD-GNGSLPEFDIYLGDKWESLV-----FE----DASSVITKEIYYAASSDYV
 ShRK1_(AT1/1-533 101 TFLYGGGLG-SEEAYPKFQLYLDATKWATVT-----IQ----EVSRYVVEELIVRATSSYV
 ShRK2_(AT2/1-529 101 TFLYGNFDNSNNVYPKFDISLGATHWATIV-----IS----ETYI IETAEELVFLASSPTV
 AT5G48740/1-492 95 TFVYRNYD-SQNSPPAFHVSLGRRITSTVD-----LR----TNDPWIEELVWPVNNDL
 AT1G51790/1-518 108 DFMYGNVD-SRNEIPGFDLHLGPNKWDTVELVSPLQT-----VSKEIYYVLTDTI
 IOS1(AT1G5/1-513 107 NFLHGGYD-DKPST-QFELYLGPNLWSTVT-----TT----NETEASIFEMIHILTDRRL
 AT1G51910/1-508 108 TFVYGNVD-GLNQMPKFDLHIGPNKWTSVI-----LEGVANA--TIFEIHHVLTQDRL
 AT1G51890/1-499 104 TFIYGNVD-GLNQLPSFDLYIGPNKWTSVS-----IPGVRNG--SVSEMIHVLQRDHL
 AT1G51860/1-511 108 TFVYGNVD-GLNQFPPFDLHIGPNKWSVK-----ILGVTNT--SMHEIHHVLPQDSL
 AT1G51880/1-511 108 TFIYGNVD-GLRQVPKFDIHHIGPSKWTSVK-----LDGVGNG--AVLEMIHVLQDRL
 AT1G07550/1-504 108 MFRYGNVD-GLNNSPRFDLYLGNPIWTTID----MGKS--GDG--VLEEIIHITRNSL
 AT2G14440/1-502 109 LFFYGNVD-GLNTSPRFDLFLGPNIWTSVD----VQKVDGGDG--VIEEIIHVTRCNIL
 AT2G14510/1-508 108 FYYGNVD-GLNTSPRFDLHIGPNIWTSVD----VLIADVGDG--VVEEIVHFWRNSL
 AT3G46350/1-491 110 RFTYGNVD-GRNMSPTFDLYLGPNLWKRID----MTKL--QNKVSTLEEITYIPLNSL
 AT3G46340/1-513 112 TALLYGNVD-GLNISPKFDLYIGANFWTTID----AGEY--LSG--VVEEVNYIPRNSL
 AT3G46370/1-427 20 TSNYGNVD-GRNEPPRFDLYLGNPFWTTID----LGKH--VNG-DTWKEIHHIPKNSL
 AT3G46400/1-508 108 TINYGNVD-GLNISPRFDLYLGNPFWTTID----LEKH--VGG-DTWKEIHHIPKNSL
 AT3G46330/1-516 110 TFFYGNFD-GLNVSPFDMHIGPNKWTID----LQIV--PDG--TVKEIHHIPRNSL
 AT5G59670/1-499 108 SFLYGNVD-GHNIAPVFDLYLGPNLWAKID----L-QD--VNG--TGEIHLHPTNSL
 AT5G59680/1-508 109 RFVYGNVD-GRDTGPKFDLYLGNPWAATID----LAKQ--VNG--TRPEIHHIPTSNL
 AT5G59650/1-509 109 RFIYGNVD-GRNSNPIFELHLGPNLWATID----LQKF--VNG--TMEIHLHPTNSL
 AT5G16900/1-505 103 VFVYGNVD-GYNDYPSFDLYLGNPKWVRVD----LEGK--VNG--SVEEIIHIPSNSL
 AT1G07560/1-512 109 MFTYGNVD-NLNTHPKFDLYLGNPIWTTVD----LQRN--VNG--TRAIEIHHIPRSTSL
 AT4G20450/1-530 112 VFVYGNVD--LKQRPKFDLYLGNPFWTTIN----LQRIWLQDG--TVEEVIHMPKSNL
 AT2G28960/1-507 108 GFGYGNVD-GLNVYPKFDLHVGNMWAIV---DLEFGKD-----REIYYMTTSNLL
 AT2G29000/1-506 106 SFVYGNVD-GRNIMPRFDLYIGPNIWAVVSELDL-YSPE-----EIEIHMTKSTSL
 AT2G28970/1-409 89 -----
 AT2G28990/1-506 104 SFVYGNVD-GLNRDPNFDIHLGPNKWKRID----LDGE--KEG-TREEIHKARSNSL
 AT1G49100/1-518 109 AFLYGNVD-GLNTVFNFDLFIGPNKVTTVN----FNAT--GGG--VFVEIHHMSRSTPL
 AT1G51810/1-384 1 -----MWITVN-----TDN--TIKEIHLVSKNSL
 AT1G51805/1-504 105 TFVYGNVD-GHKDEPNFDLYLGPNLWATVS-----RSE--TVEEIIHVTKSDSL
 AT1G51830/1-315 10 -----
 AT1G51820/1-504 105 TFVYGNVD-GLNVGNPNLYLGPNLWTTVS-----SND--TIEEIIHVTRNSL
 AT1G51850/1-486 105 TFVYGNVD-GLNVGNPNFDLYFGPNLWTT-----
 AT2G04300/1-480 111 SFVYGNVD-GLNKELEFDLYLGPNLWANVNTAVYLMNGV--TT----EIEIHSKSKVL
 AT3G21340/1-520 112 VFVYGNVD-GLNNPPSFDLYLGNPLWVTVD----MNGR--TNG--TIQEIIHKTISKSL
 AT1G05700/1-507 107 SFMYGNVD-GENGSPFDLFLGNIWDTVL----LS----NGSSIVSKEVVYLSQSENI
 AT2G19210/1-516 112 RFMYGNVD-NLGKAPDFDLYLGFNIWDSVT-----ID----NATTIVTKEIHTLRSDHV
 AT2G19230/1-516 111 RFMYGNVD-NLGKAPDFDLYLGVNLWDSVT----LE----NSTTIVTKEIHYTLRSDKV
 AT2G19190/1-516 111 RFMYGNVD-DLGRVPEFDLYLGVNFWDSVK----LD----DATIILNKEIITIPLLDNV
 AT4G29990/1-511 108 RFMYGNVD-GFSKTPFDLYIGANLWESVV----LI----NETAIMTKEIHYTPPSDHI
 AT4G29180/1-509 112 SFMYGNVD-GKNALPEFDLYVNVNFWTSVK----LR----NASENVIKEILSFAESDTI
 AT4G29450/1-513 111 SFMYGNVD-GKKALPEFDLYVNVNFWSTVK----FK----NASDQVTKEILSFAESDTI

LjSYM RK/1-509 151 DFCLLK--GEVYPFISQLELRPSPEE-YLQDFPTS--VLKLI SRNNL-GDT----KDDIR
 LjShRK1/1-522 145 SVCLSN-ATTGQPFISTLELRQFNGSVYTDY-EEHFYLSVSARINFGAES----DAPIR
 LjShRK2/1-506 158 HVCLFN-TGKGTFFISVLELRVLNSDAYLV---N---SLELLARFDVGLRD----GEIIR
 ShRK1_ (AT1/1-533) 151 DVCVCC-AITGSPFMSTLELRPLNLSMYATDY-EDNFFLKVAARVNFAGPN----MDALR
 ShRK2_ (AT2/1-529) 152 SVCLSN-ATTGQPFISTLELRQLSGSMYGSMLEDRFYLSVAARINFGAES----EASVR
 AT5G48740/1-492 144 LLCLLAVKGRGIPVISSLEVRPLPLGSYKYSLEG---SPDIILRRSYRINSGYT-NGTIR
 AT1G51790/1-518 158 QVCLVN-TGNGTFFISVLELRQLPNSSYAAQS-E---SLQLFQRLDLFGSTT----NLTVR
 IOS1 (AT1G5/1-513) 156 QICLVK-TGNATPFISALELRKLMNNTTYLTRQ-G---SLQTFIRADVGATV----NQGYR
 AT1G51910/1-508 158 QVCLVK-TGQTPFISLELRPLNNDTYVTQG-G---SLMSFARIYF-PKT----AYFLR
 AT1G51890/1-499 154 QICLVK-TGETTPFISLELRPLNNTTYVTKS-G---SLIVVARLYF-SPT----PPFLR
 AT1G51860/1-511 158 EVCLVK-TGTPPFISLEVRPLNNSYLTQS-G---SLMLFARVYFPSSS----SSFIR
 AT1G51880/1-511 158 QICLVK-TGKGI PFISLELRPLNNTTYLTQS-G---SLIGFARVFF-SAT----PTFLR
 AT1G07550/1-504 158 DICLVK-TGTSTPMISSLELRPLLYDTYIAQT-G---SLRNYNRFYF-TDS----NNYIR
 AT2G14440/1-502 161 DICLVK-TGTTTPMISALELRPLRYDTYTART-G---SLKKILHFYF-TNS----GKEVR
 AT2G14510/1-508 160 DICLVK-TGTSTPMISALELRPLRYDTYTART-G---SLKSMAHFYF-TNS----DEAIR
 AT3G46350/1-491 162 DVCLVK-TNTTTPFISALELRPLPSNSYITTA-G---SLRTFVRFCF-SNS----VEDIR
 AT3G46340/1-513 162 DVCLVK-TDTSTPFLSLELRPLDNDSYLTGS-G---SLKTFRRYYL-SNS----ESVIA
 AT3G46370/1-427 71 DVCLIK-TGTTTPIISTLELRSLPKYSYNAIS-G---SLKSTLRAFL-SES----TEVIR
 AT3G46400/1-508 159 DVCLIK-TGTSTPIISVLELRPLNNTYITES-G---SLKSILRSYL-SVS----NLTVR
 AT3G46330/1-516 160 QICLVK-TGATTPMISALELRPLANDTYIAKS-G---SLKYYFRMYL-SNA----TVLLR
 AT5G59670/1-499 157 QICLVQ-TGETTPLISSLELRPLMRTGSYTTVS-G---SLKTYRRLYF-KKS----GSRLR
 AT5G59680/1-508 159 QVCLVK-TGETTPLISVLEVRPMGSGTYLTKS-G---SLKLYREYF-SKS----DSSLR
 AT5G59650/1-509 159 NVCLVK-TGTTPLISALELRPLGNNSYLT-D-G---SLNLFVRIYL-NKT----DGFLR
 AT5G16900/1-505 153 QICLVK-TGNSLPPISALELRPLRNDTYVVQD-V---SLKHLFRVYY-RQS----DRLIR
 AT1G07560/1-512 159 QICLVK-TGTTPLISALELRPLRNDTYIPQS-G---SLKTLFRVHL-TDS----KETVR
 AT4G20450/1-530 163 DICLVK-TGTTTPFISLELRPLRDDTYTTT-G---SLKLI SRWYF-RKPFPTLESIIR
 AT2G28960/1-507 155 QICLVK-TGSTIPMISTLELRPLRNDSYLTQF-G---PLDLIYRRAYSNS----TGFIR
 AT2G29000/1-506 155 QICLVK-TGPTTFFISTLELRPLRNDYITQS-G---SLKLMQRMCM-TET----VSTLR
 AT2G28970/1-409 89 -----SLRNSFRVHC-SIS----DSEIR
 AT2G28990/1-506 154 DICLVK-TGETLPIISAIEIRPLRNDTYVTQS-G---SLMMSFRVYL-SNS----DASIR
 AT1G49100/1-518 159 DICLVK-TGTTTPMISTLELRPLRSDTYISAI-GS--SLLLYFRGYL-NDS----GVVLR
 AT1G51810/1-384 24 QVCLVK-TGTSIPYINTLELRPLADDIYTNES-G---SLNYLFRVYY-SNL----KGYIE
 AT1G51805/1-504 151 QVCLAK-TGDFIPFINILELRPLKKNVYVTES-G---SLKLLFRKYF-SDS----GQDIR
 AT1G51830/1-315 10 -----VIR
 AT1G51820/1-504 151 QVCLVK-TGISIPFINMLELRPMKKNMYVTQS-G---SLKYLFRGYI-SNS----STRIR
 AT1G51850/1-486 132 -VCLIK-TGISIPFINVLELRPMKKNMYVTQG-E---SLNYLFRVYI-SNS----STRIR
 AT2G04300/1-480 163 QVCLIK-TGESIPINISLELRPLINDTYNTQS-G---SLKYLFRNYF-SIS----RRIIR
 AT3G21340/1-520 162 QVCLVK-TGTSSPMINTLELRPLKNDTYNTQS-G---SLKYFRYF-SGS----GQDIR
 AT1G05700/1-507 157 FVCLGN-KGKGTFFISTLELRFLGNDTYDSPN-G---ALFFSRRWDLRSLM----GSPVR
 AT2G19210/1-516 162 HVCLVD-KNRGTPFLSALEIRLLKSNTYETPY-D---SLILFKRWDLGGLG----ALPVR
 AT2G19230/1-516 161 HVCLVD-KERGTFFLSVLELRLLKNNIYETAS-D---SLMLYRRWDLGATG----DLPAR
 AT2G19190/1-516 161 QVCVVD-KNAGTPFLSVLEIRLLNNTYETPY-D---ALTLRLRLDYSKTG----KLPSR
 AT4G29990/1-511 158 HVCLVD-KNRGTPFLSVLEIRFLKNDTYETPY-E---ALMLGRRWDFGTAT----NLQIR
 AT4G29180/1-509 162 YVCLVN-KGKGTFFISALELRPMNSSIYGTEF-GRNVSLVLYQRWDT-GYL----NGTGR
 AT4G29450/1-513 161 YVCLVN-KGKGTFFISGLELRPLVNSSIYGTEF-GRNVSLVLYRWDI-GYL----NGTGR

LjSYM RK/1-509 201 **F**PVDQ**S**DR**I**W**K**A-----SSISSAVPLSSNVSNVDLNAVVT**P**PL**T**VL**Q**TAL**T**--
 LjShRK1/1-522 199 **Y**PDD**P**FDR**I**W**E**SDSVKKAN**Y**LVDVAPG**T**T**K**IS**T**KE**P**ID**V**---NR**D**EM**P**PGR**V**M**Q**TAV**V**GT
 LjShRK2/1-506 207 **Y**PDD**T**FDR**M**W**T**P-----Y**N**S**I**E--**W**K**L**M**N**T**S**L**T**I**D**Q**P**S**F**N**F**L**P**L**P**S**I**V**S**T**A**A**I**PA
 ShRK1_ (AT1/1-533 205 **Y**PDD**P****Y**DR**I**W**E**SD**I**N**K**R**P****N****Y**L**V**G**V**APG**T****T**R**I**N**T**S**K**T**I**N**T**---L**T**R**E****Y****P****P**M**K****V**M**Q**TAV**V**GT
 ShRK2_ (AT2/1-529 207 **Y**PDD**P****Y**DR**I**W**E**SD**L**Q**K**K**P****N****Y**L**V**D**V**AAG**T****V**R**V**S**T**L**P**I**E**S---R**V**D**D**R**P****P****Q****K****V**M**Q**TAV**V**GT
 AT5G48740/1-492 200 **Y**P**S**D**P**FDR**I**W**D**P**Q**S**Y**S**P**F**H**A**S**W**S**F**N**G**L**T**K**L**N**-----S**F**N**I**T**E**N**P****P**A**S****V**L**K**T**A**R**I**L**A**
 AT1G51790/1-518 209 **Y**P**N**D**V**FDR**I**W**F**P-----A**T**P**N**G**T**K**P**L**S**D**P**S**T**S**L**T**S**N**S**-**T**G**N**--**F**R**L**P**Q****V****V**M**R**T**G**I**V****D**
 IOS1 (AT1G5/1-513 207 **Y**G**I**D**V**FDR**V**W**T**P-----**Y**N**F**G**N**--**W**S**Q**I**S**T**N**Q**S****V**---N**I**N**N**D**Y****Q****P****E**I**A**M**V**T**A**S**V****P**T
 AT1G51910/1-508 208 **Y**S**D**D**L****L****D**R**V**W**V**P-----**F**S**Q**-**N**E--**T**V**S**L**S**T**N**L**P**V**D**T--**S**S**N**-**S****Y**N**V****P****Q****N****V**A**N**S**A**I**I**PA
 AT1G51890/1-499 204 **Y**D**E**D**V**HDR**I**W**I**P-----**F**L**D**-**N**K--**N**S**L**S**T**E**L**S**V**D**T**--**S**N-**F****Y**N**V****P****Q****T****V**A**K**T**A**A**V****P**L
 AT1G51860/1-511 209 **Y**D**E**D**I**HDR**V**W**N**S-----**F**T**D**-**D**E--**T**V**W**I**S**T**D**L**P**I**D**T--**S**N-**S****Y**D**M****P****Q****S****V**M**K**T**A**A**V****P**K
 AT1G51880/1-511 208 **Y**D**E**D**I**HDR**V**W**V**R-----**Q**F**G**-**N**G--**L**K**S**I**S**T**D**L**L**V**D**T--**S**N-**P****Y**D**V****P****Q**A**V**A**K**T**A**C**V****P**S
 AT1G07550/1-504 208 **Y**P**Q**D**V**HDR**I**W**V**P-----**L**I**L**-**P**E--**W**T**H**I**N**T**S**H**H**V**I**D-**S**I**D**-**G****Y**D**P****P****Q****D****V**L**R**T**G**A**M****P**A
 AT2G14440/1-502 211 **Y**P**E**D**V****Y**DR**V**W**I**P-----**H**S**Q**-**P**E--**W**T**Q**I**N**T**T**R**N**V**S**G-**F**S**D**-**G****Y**N**P****P****Q****D****V**I**K**T**A**S**I****P**T
 AT2G14510/1-508 210 **Y**P**E**D**V****Y**DR**V**W**M**P-----**Y**S**Q**-**P**E--**W**T**Q**I**N**T**T**R**N**V**S**G-**F**S**D**-**G****Y**N**P****P****Q****Q****V**I**Q**T**A**S**I****P**T
 AT3G46350/1-491 212 **F**P**M**D**V**HDR**M**W**E**S-----**Y**F**D**-**D**D--**W**T**Q**I**S**T**S**L**T**V**N**T-**S**-**D**-**S****F**R**L****P****Q**A**A**L**I**T**A**A**T****P**A
 AT3G46340/1-513 212 **Y**P**E**D**V****K**DR**I**W**E**P-----**T**F**D**-**S**E--**W**K**Q**I**W**T**L**K**P**N**N**-**S**-**N**-**G****Y**L**V****P****K****N****V**L**M**T**A**A**I**PA
 AT3G46370/1-427 121 **Y**P**N**D**F****Y**DR**M**W**V**P-----**H**F**E**-**T**E--**W**K**Q**I**S**T**N**L**K**V**N**S-**S**-**N**-**G****Y**L**L****P****Q****D****V**L**M**T**A**A**I****P**V
 AT3G46400/1-508 209 **Y**P**D**D**F****Y**DR**K**W**V**P-----**Y**F**E**-**S**E--**W**R**Q**I**S**T**L**L**K**V**N**N-**T**I**N**-**G****Y**D**M****P****Q****S****V**M**K**T**A**A**V****P**K
 AT3G46330/1-516 210 **Y**P**K**D**V****Y**DR**S**W**V**P-----**Y**I**Q**-**P**E--**W**N**Q**I**S**T**T**S**N**V**S**N-**K**-**N**-**H****Y**D**P****P****Q****V**A**L**K**M**A**A**T**P**T
 AT5G59670/1-499 207 **Y**S**K**D**V****Y**DR**S**W**F**P-----**R**F**M**-**D**E--**W**T**Q**I**S**T**A**L**G**V**I**N-**T**-**N**-**I****Y****Q****P****P****E**D**A**L**K**N**A**A**T****P**T
 AT5G59680/1-508 209 **Y**P**D**D**I****Y**DR**Q**W**T**S-----**F**F**D**-**T**E--**W**T**Q**I**N**T**T**S**D**V**G**N-**S**-**N**-**D****Y****K****P****P****K****V**A**L**T**A**A**I****P**T
 AT5G59650/1-509 208 **Y**P**D**D**I****Y**DR**R**W**H**N-----**Y**F**M****V****D**D--**W**T**Q**I**F**T**T**L**E**V**T**N-**D**-**N**-**N****Y****E****P****P****K****A**L**A**A**A**A**T****P**S
 AT5G16900/1-505 203 **Y**P**D**D**V****Y**DR**V**W**S**P-----**F**F**L**-**P**E--**W**T**Q**I**T**T**S**L**D**V**N**N-**S**-**N**-**N****Y****E****P****P****K****A**A**L**T**S**A**A**T**P**G
 AT1G07560/1-512 209 **Y**P**E**D**V**HDR**L**W**S**P-----**F**F**M**-**P**E--**W**R**L**L**R**T**S**L**T**V**N**T-**S**D**D****N****G****Y****D**I**P**E**D****V****V**T**A**A**T****P**A
 AT4G20450/1-530 217 **H**P**D**D**V**HDR**L**W**D**V-----**Y**H**A**D**E**E--**W**T**D**I**N**T**T****P**V**N**T-**T**V**N**-**A****F****D**L**P****Q**A**I**I**S****K**A**S**I**P**Q
 AT2G28960/1-507 206 **Y**P**D**D**I**FDR**K**W**D**R-----**Y**N**E**-**F**E--**T**D-**V**N**T**L**N**V**R**S-**S**-**S**-**P****F****Q****V****P****E**A**V**S**R**M**G**I**T****P**E
 AT2G29000/1-506 205 **Y**P**D**D**V****Y**DR**L**W**Y**T-----**D**G**I**-**Y**E--**T**K**A**V**K**T**A**L**S**V**N**S-**T**-**N**-**P****F****E**L**P****Q****V**I**I**R**S**A**A**T**P**V
 AT2G28970/1-409 107 **Y**D**D**S**S****Y**DR**V**W**Y**P-----**F**F**S**-**S**S--**F**S**Y**I**T**T**S**L**N**I**N**N-**S**-**D**-**T****F****E**I**P****K**A**A**L**K**S**A**A**T****P**K
 AT2G28990/1-506 204 **Y**A**D**D**V**HDR**I**W**S**P-----**F**N**G**-**S**S--**H**T**H**I**T****D**L**N**I**N**N-**S**-**N**-**A****Y****E**I**P****K****N**I**L****Q**T**A**A**I****P**R
 AT1G49100/1-518 210 **Y**P**D**D**V**NDR**R**W**F**P-----**F**S**Y****K**E--**W**K**I****V**T**T**L**N**V**N**T-**S**-**N**-**G****F****D**L**P****Q****G**A**M**A**S**A**A**T**R**V
 AT1G51810/1-384 74 **Y**P**D**D**V**HDR**I**W**K**Q-----**I**L**P****Y****Q**D--**W**Q**I**L**T**T**N**L**Q**I**N**V-**S**-**N**-**D****Y****D**L**P****Q****R****V**M**K**T**A**V**T****P**I
 AT1G51805/1-504 201 **Y**P**D**D**I****Y**DR**V**W**H**A-----**S**F**L**E**N**N--**W**A**Q****V**S**T**L**G**V**N**V-**T**-**D**-**N****Y****D**L**S****Q****D****V**M**A**T**G**A**T****P**L
 AT1G51830/1-315 12 **F**P**D**D**V****Y**DR**K**W**Y**P-----**I**F-**Q**N**S**--**W**T**Q****V**T**T**N**L**V**N**I-**S**-**T**-**I****Y****E**L**P****Q****S****V**M**S**T**A**A**T****P**L
 AT1G51820/1-504 201 **F**P**D**D**V****Y**DR**K**W**Y**P-----**L**F-**D**D**S**--**W**T**Q****V**T**T**N**L**K**V**N**T**-**S**-**I**-**T****Y****E**L**P****Q****S****V**M**A****K**A**A**T**P**I
 AT1G51850/1-486 181 **F**P**D**D**V****Y**DR**K**W**Y**P-----**Y**F-**D**N**S**--**W**T**Q****V**T**T**L**D**V**N**T-**S**-**L**-**T****Y****E**L**P****Q****S****V**M**A****K**A**A**T**P**I
 AT2G04300/1-480 213 **Y**P**N**D**V**NDR**H**W**Y**P-----**F**F**D**E**D**A--**W**T**E**L**T**T**N**L**N**V**N**S-**S**-**N**-**G****Y**D**P****P****K****F****V**M**A**S**A**S**T****P**I
 AT3G21340/1-520 212 **Y**P**D**D**V**NDR**K**W**Y**P-----**F**F**D**A**K**E--**W**T**E**L**T**T**N**L**N**I**N**S-**S**-**N**-**G****Y**A**P****P****E****V****V**M**A**S**A**S**T****P**I
 AT1G05700/1-507 208 **Y**D**D**D**V****Y**DR**I**W**I**P-----**R**N**F**G**Y**--**C**R**E**I**N**T**S**L**P**V-**T**-**S**D**N**N**S****Y**S**L**S**S**L**V**M**S**T**A**M**T****P**I
 AT2G19210/1-516 213 **Y**K**D**D**V**FDR**I**W**I**P-----**L**R**F****P****K**--**Y**T**I**F**N**A**S**L**T**I**D**S-**N**N**N**E**G****F****Q****P**A**R**F**V**M**N**T**A**T**S****P**E
 AT2G19230/1-516 212 **Y**K**D**D**I**FDR**F**W**M**P-----**L**M**F****P****N**--**F**L**I**L**N**T**S**L**M**I**D**P-**T**S**S****N**G**F**L**P****P****S****V**M**S**T**A**V**A****P**M
 AT2G19190/1-516 212 **Y**K**D**D**I****Y**DR**I**W**T**P-----**R**I**V**S**S**E--**Y**K**I**L**N**T**S**L**T**V**D**Q-**F**L**N****N****G****Y****Q****P**A**S**T**V**M**S**T**A**E**T**A**R**
 AT4G29990/1-511 209 **Y**K**D**D**F****Y**DR**I**W**M**P-----**Y**K**S****P****Y**--**Q****K**T**L**N**T****S**L**T**I**D**E-**T**N**H****N**G**F**R**P**A**S**I**V**M**R**S**A**I**A****P**G
 AT4G29180/1-509 215 **Y**Q**K**D**T****Y**DR**I**W**S**P-----**Y**S**P**-**V**S--**W**N**T****M****T****T****G****Y**I**D**I-**F****Q**S-**G****Y**R**P****P****D****E****V**I**K**T**A**A**S****P**K
 AT4G29450/1-513 214 **Y**Q**D**D**R**FDR**I**W**S**P-----**Y**S**S****N**I**S**--**W**N**S**I**I**T**S**G**Y**I**D**V-**F****Q**N-**G****Y****C****P****P****D****E****V**I**K**T**A**A**P**E

LjSYM RK/1-509 248 ----DPERLEFIHTDLETEDYGYRVFLYFLELDR-TLQAGQ-RVFDIY-----
 LjShRK1/1-522 256 NGS-----LTYRMNLDGFPGIGWAVCYFAEIE--DLPQNESRKFRLLVLPGQPDISKAVV
 LjShRK2/1-506 257 NVN-DN---IEFYHHPKYNASTYMYMYFDEIK--KLQANQIREFDIF-----
 ShRK1_ (AT1/1-533 262 QGL-----ISYRLNLEDFPANARAYAYFAEIE--ELGANETRKFKLVQPYFPDYSNAVV
 ShRK2_ (AT2/1-529 264 NGS-----LTYRMNLDGFPGFGWAFYFAEIE--DLAEDESRKFRLLVLPQPEYSKSVV
 AT5G48740/1-492 252 RKE-S---LSYTLSLHTPGD-YYIILYFAGIL--SL----SPSFSVT-----
 AT1G51790/1-518 259 NPR-GF---VDFGWIPDDPSLEFFFYLYFTELQQPNSGTVETREFVIL-----
 IOS1 (AT1G5/1-513 254 DPD-AA---MNISLVGVERTVQFYVFMHFAEIQ--ELKSNDTREFNIM-----
 AT1G51910/1-508 256 EAT-HP---LNIWWDLQINAPSYVVMHFAEIQ--NLKANDIREFNIT-----
 AT1G51890/1-499 251 NAT-QP---LKNINWLDLITSQSYIYM HFAEIE--NLEANETREFNIT-----
 AT1G51860/1-511 256 NAS-EP---WLLWWTLDENTASQSYVVMHFAEVQ--NLTANETREFNIT-----
 AT1G51880/1-511 255 NAS-QP---LIFDWTLDNITSQSYVVMHFAEIQ--TLKDNIREFNIT-----
 AT1G07550/1-504 256 NAS-DP---MTITWNLKTATDQVYGYIYIAEIM--EVQANETREFEVV-----
 AT2G14440/1-502 259 NVS-EP---LFTWMSSESSDDETYAYLYFAEIQ--QLKANETRFKIL-----
 AT1G514510/1-508 258 NGS-EP---LFTWNLDESDDDETYAYLFFAEIQ--QLKVNNTREFNIT-----
 AT3G46350/1-491 259 KDG-PSYIGITFSTSSSE---ERFFIYLHFSEVQ--ALRANETREFNIS-----
 AT3G46340/1-513 259 NDS-AP---FRFTEELDSPTDELYVYLHFSEVQ--SLQANESREFDIL-----
 AT3G46370/1-427 168 NTS-AR---LSFTENLEFPDDELHYLYHFSEVQ--VLQANQSREFSIL-----
 AT3G46400/1-508 257 NAS-VP---LSFTKDLDFPKDKLYFYHFSEIQ--PLQANQSREFSIL-----
 AT3G46330/1-516 257 NLD-AA---LTMVWRLENPDDQIYLYMHFSEIQ--VLKANDTREFDII-----
 AT5G59670/1-499 254 DAS-AP---LTFKWNSEKLDVQYFYAHYAEIQ--DLQANDTREFNIL-----
 AT5G59680/1-508 256 NAS-AP---LTNEWSSVNPDEQYVYAHFSEIQ--ELQANETREFNML-----
 AT5G59650/1-509 256 NAS-AP---LTI SWPPDNFGDQYLYSHFSEIQ--DLQNTDREFDIL-----
 AT5G16900/1-505 250 DNG-TR---LTI IWTLDNPDDEQIHLVYHFAELE--PVGENTTRTFYFV-----
 AT1G07560/1-512 258 NVS-SP---LTI SWNLETPDDLVYAYLHVAEIQ--SLRENDTREFNIS-----
 AT4G20450/1-530 266 VAS-DT---WSTTWSIQNPDDDVHLYLHFAEIQ--ALKPSDTREFSIL-----
 AT2G28960/1-507 252 NAS-LP---LRFYVSLDDSDKVN VFHFAEIQ--ALRCNETREFDIE-----
 AT2G29000/1-506 252 NSS-EP---ITVEYGGYSSGDQVYLYLHFAEIQ--TLKASDNREFDIV-----
 AT2G28970/1-409 154 NAS-AP---LIITWKPRPSNAEVYFYLHFAEIQ--TLAANETREFDIV-----
 AT2G28990/1-506 251 NAS-AP---LIITWDPLPINAEVYLYMHFAEIQ--TLEANETRFQDVI-----
 AT1G49100/1-518 257 NDN-GT---WEFPWSLEDSTTRFHIYLFHFAELQ--TLLANETREFNVL-----
 AT1G51810/1-384 122 KASTTT---MEFPWNLEPPTSQFYFLHFAELQ--SLQANETREFNVV-----
 AT1G51805/1-504 249 NDS-ET---LNI TWNVEPPTTKVYSYMHFAELE--TLRANDTREFNVM-----
 AT1G51830/1-315 59 NAN-AT---LNI TWTEIEPPTTFYSYIHFAELQ--SLRANDTREFNVT-----
 AT1G51820/1-504 248 KAN-DT---LNI TWTEVEPPTTFYSYVHIAEIQ--ALRANETREFNVT-----
 AT1G51850/1-486 228 KAN-DT---LNI TWTEVEPPTTFYSYMHFAELQ--TLRANDAREFNVT-----
 AT2G04300/1-480 261 SKN-AP---FNF TWSLIPSTAKFYSYMHFAEQ--TLQANETREFDMM-----
 AT3G21340/1-520 260 STF-GT---WNFSWLLPSTTQFYVVMHFAEIQ--TLRSLDTREFKVT-----
 AT1G05700/1-507 256 NTT-RP---ITMTLENSDPNVRFYVVMHFAEVEDLSLKPNTREFDIS-----
 AT2G19210/1-516 262 DLS-QD---IIFSWEPKDPWKYFVVMHFAEVV--ELPSNETREFKIL-----
 AT2G19230/1-516 261 NSSIEQ---IMVYWEPRDPNWKFYIYIHFAEVE--KLPSNETREFSVF-----
 AT2G19190/1-516 262 NES-LY---LTL SFRPPDPNAKFYVVMHFAEIE--VLKSNQTFREFSIW-----
 AT4G29990/1-511 258 NES-NP---LKFNWAPDDPRSKFYIYM HFAEVR--ELQRNETREFDIY-----
 AT4G29180/1-509 263 SDD-EP---LELSWTSSDPDRFYAYLYFAELE--NLKRNESREIKIF-----
 AT4G29450/1-513 263 NVD-DP---LELFWTSDDPNVRFYAYLYFAELE--TLEKNETR KIKIL-----

LjSYMRK/1-509 290 -VNSEIK-----KESFDVLAGGSNYRYDVLDISASGSLNV-TLVKASKS-EFGPLL
 LjShRK1/1-522 308 NIEENALGKYRLYEPGYTNLSLPPVLSF-----RFGKTSDS-TRGPLL
 LjShRK2/1-506 299 -VN-GKLFNNDPVNPVYL--KSLYYISAI-----AKPHLEL-WINRTSRs-TLPLLI
 ShRK1_ (AT1/1-533 314 NIAENANGSYTLYEPSYMNVTLDLFTF-----SFGKTKDS-TQGPLL
 ShRK2_ (AT2/1-529 316 NIKENTQRPYRVYAPGYFNITLPPVLFN-----RFAKTADS-SRGPIL
 AT5G48740/1-492 288 -INDEVK--QSDYTVTSSEAGTLYFTQKG-----ISKLNITLRIKIFNPQV
 AT1G51790/1-518 303 -LNGKSF--GEPKSLNYFRTLALFTSNPL-----KAESFQF-SLRQTQSS-SLPLLI
 IOS1 (AT1G5/1-513 296 -YN--NKHIYGPFRPLNFTSSVFTPTTEV-----VADANGQYIF-SLQRTGNS-TLPLLI
 AT1G51910/1-508 298 -YNGGQVW-ESSIRPHNLSITTISSPTAL-----NS-SDGFFNF-TFTMTTTS-TLPLLI
 AT1G51890/1-499 293 -YNGGENW-FSYFRPPKFRITTVYNPAAV-----SS-LDGNFNF-TFSMTGNS-THPLLI
 AT1G51860/1-511 298 -YNGGLRW-FSYLRPNLSISTLYFNPRAV-----SS-SNGIFNF-TFAMTGNS-TLPLLI
 AT1G51880/1-511 297 -YNGGQNV-YSYLRPEKFEISTLFDKPL-----SS-PDGSFSL-SFTKTGNS-TLPLLI
 AT1G07550/1-504 298 -VNNKVH--FDPFRPTREFAQVMFNVP-----TC-EGGFCLL-QLIKTPKS-TLPLLM
 AT2G14440/1-502 301 -VNGVYV--ID-YIPRKFEAETLITPAAL-----KC-GGGVCRV-QLSKTPKS-TLPPQM
 AT2G14510/1-508 300 -ANGVDV--ID-YTPWKFEARTLNPAAL-----KC-EGGVCRV-QLSKTPKS-TLPLLM
 AT3G46350/1-491 301 -INGESV--ADLYRP-----LSRTQSS-THPPMI
 AT3G46340/1-513 301 -WSGEVA--YEAFIPEYLNITTIQTNTPV-----TC-PGKCNL-ELKRTKNS-THPLLI
 AT3G46370/1-427 210 -WNGMVI--YPDFIPDYLGAAVYVNPSPS-----LC-EVGKCLL-ELERTQKS-TLPLLI
 AT3G46400/1-508 299 -WNGELI--IPTLSPKYLKASTLYSVSPF-----VC-EVGKCLL-QLVRSR-TPPLLI
 AT3G46330/1-516 299 -LNGETIN-TRGVTPKYLEIMTWLTTNPR-----QC-NGGICRM-QLTKTQKS-TLPLLI
 AT5G59670/1-499 296 -LNGQNSVGTGPEVPDKLSIKTFQSSSPI-----SC-NGWACNF-QLIRTKRS-TLPLLI
 AT5G59680/1-508 298 -LNGKLF--FGPVVPPKLAISTLISVSPN-----TC-EGGECNL-QLIRTNRS-TLPLLI
 AT5G59650/1-509 298 -WDGAVV--EEGFIPPKLGVTTIHNLSPV-----TC-KGENCIY-QLIKTSRS-TLPSLL
 AT5G16900/1-505 292 -VNGKISY-DESIPLDLAVSTVETVVN-----KC-DGGNCSL-QLVRSR-TPPLLI
 AT1G07560/1-512 300 -AGQDVN--YGPVSPDEFVLVGTTFNTSPV-----KC-EGGTCHL-QLIKTPKS-TLPLLI
 AT4G20450/1-530 308 -WKNKNTII-RDYSPLEFMADTVPIRTSS-----KCGDDGFCSL-DLRTKSS-TLPPYC
 AT2G28960/1-507 294 -LEEDII--QSAYSPTMLQSDTKYNLSPH-----KC-SSGLCYL-KLVRTPRS-TLPLLI
 AT2G29000/1-506 294 -WANNIK--KLAYPKVQIDTLNLTSPN-----KC-DNTFCKA-FLVRTQRS-TLPLLI
 AT2G28970/1-409 196 -FKGNFN--YSAFSPKLELLTFFTSGPV-----QC-DSGKCNL-QLVTPNS-TLPLLI
 AT2G28990/1-506 293 -LRGNFN--HSGFSPKLVKVFVLYTEEPM-----KC-GSEGCYL-QLVTPNS-TLPLLI
 AT1G49100/1-518 299 -LNGKVY--YGPYSPKMLSIDTM---SQPDSLTLC-KGGSCCL-QLVTKS-TLPLLI
 AT1G51810/1-384 165 -LNGNVT--FKYSPPKLEMQTVYSTAPK-----QC-DGGKCLL-QLVTKS-TLPLLI
 AT1G51805/1-504 291 -LNGNDL--FGPYSPKPLKTEETNLKPE-----EC-EDGACIL-QLVTKS-TLPLLI
 AT1G51830/1-315 101 -LNGEYT--IGPYSPKPLKTEETQDLSPE-----QC-NGGACIL-QLVTKS-TLPLLI
 AT1G51820/1-504 290 -LNGEYT--FGPSPKPLKTEETQDLSPE-----QC-DGGRCIL-QVVKTLKS-TLPLLI
 AT1G51850/1-486 270 -MNGIYT--YGPYSPKPLKTEETIYDKIPE-----QC-DGGACIL-QVVKTLKS-TLPLLI
 AT2G04300/1-480 303 -LNGNLA--LER-----
 AT3G21340/1-520 302 -LNGKLA--YERYSPKTLATETIFYSTPQ-----QC-EDGTCLL-ELTKTPKS-TLPLLM
 AT1G05700/1-507 300 -IN--GVTVAAGFSPKYLQNTFFL-NPE-----SQSKIAF-SLVRTPKS-TLPPIV
 AT2G19210/1-516 304 -LNEKEINMSS-FSPRYLYTDTLTVQNPV-----SGPKLEF-RLQQTPRS-TLPPII
 AT2G19230/1-516 304 -LNKEQIDTTSVFRPSYLYTDTLTVQNPV-----SGPFLEF-VLRQGVKS-TRPPIM
 AT2G19190/1-516 304 -LN--EDVISPSFKLRYLLTDTFVTPDPV-----SGITINFSLLQPPGEF-VLPPII
 AT4G29990/1-511 300 -IN--DVILAENFRPFYLFDTTRSTVDPV-----GRKMNEI-VLQRTGVS-TLPPII
 AT4G29180/1-509 305 -WNGSPVS--GAFNPSPEYSMTVSNRAF-----TGKDHWI-SVQKTAES-TRPPIL
 AT4G29450/1-513 305 -WNGSPVS-ETSFEPSKYSTTFSNPRAF-----TGKDHWI-SIQKTVDS-TLPLLI

LjSYM RK/1-509 338 NAYEILQVRPWIEE-TNQT DVGVIQKMREELLQNSEWS---GDPCI--LLPWKGIACD
 LjShRK1/1-522 350 NAMEINKYLE---KNGGSPDGEAISSVLSHY---SSADWAQEGGDPCLPVPWSW--IRCS
 LjShRK2/1-506 345 NAIEIYMTKDFLQSQTYQTADAIINVKSIYGI-KR-NWQ---GDPCIPLAYLWDGLNCS
 ShRK1_(AT1/1-533 356 NAIEISKYLPISVKTDRS-DVSVLDAIRSMS---PDSWASEGGDPCIPVLSWSW--VNCS
 ShRK2_(AT2/1-529 358 NAMEISKYLR---KSDGSDATVMANVASLY---SSTEWAEQGGDPCSPSPWSW--VQCN
 AT5G48740/1-492 331 SALEVVEILQIPPE-ASSTVSAKLVIEQFTG--QDLGWQ---DDPCTP--LPWNHIECE
 AT1G51790/1-518 350 NAMEYFVNKLQSSDTPNDLSAMRNIKSAYKV-KR-NWE---GDPCVPQAYTWEGLNCS
 IOS1(AT1G5/1-513 346 NAMEIYSVNLLPQOETDRKEVDAMNNIKSA YGV-NKIDWE---GDPCVPLDYKWSGVNCT
 AT1G51910/1-508 348 NALEVYTLVENLLETTYQDEVSAMNIIKTYGLSKKISWQ---GDPCSPQIYRWEGLNCL
 AT1G51890/1-499 343 NGLIYQVLELPLQDITYQDEVSAMNIIKTYGLSKRSSWQ---GDPCAPELYRWEGLNCS
 AT1G51860/1-511 348 NALEIYTVVDILQOSETNPDVSAAMNIIKTYGLSKKISWQ---GDPCAPQLYRWEGLNCS
 AT1G51880/1-511 347 NGLIYKVLDLLELETDQDEVSAMINIKATYDLSKKVSWQ---GDPCAPKSYQWEGLNCS
 AT1G07550/1-504 347 NAEIFFTGIEFPQSETNQNDVIAVKNIQAS YGL-NRISWQ---GDPCVPKQFLWTGLSCN
 AT2G14440/1-502 349 NAEIFSVIQFPQSDTNTDEVI AIKNIQSTYKV-SRISWQ---GDPCVPIQFSWMGVSN
 AT2G14510/1-508 348 NAEIFSVIQFPQSDTNTDEVI AIKNIQSTYQL-SRISWQ---GDPCVPKQFLWSWGLNCS
 AT3G46350/1-491 326 NAEIFLVSELLQSETYENDVIAIKKIKD TYGL-QLISWQ---GDPCVPRLYKWDGLDCT
 AT3G46340/1-513 350 NAIEFYTVVNFQOETNEDDVVAIKDIKATYEL-NRITWQ---GDPCVPKQFIWEGLDCN
 AT3G46370/1-427 259 NAIEVFTVMNFPQSETNDDDVIAITKI KDTHRL-NRTSWQ---GDPCVPLQFWSWGLSCI
 AT3G46400/1-508 348 NAIEVFTVIDFPQSETNDDVIAIKNIKDT HGL-SRVSWQ---GDPCVPRQFLWTGLNCS
 AT3G46330/1-516 349 NAIEVYVSVLQLPQSQTNEDVVAIKNI RTTYGL-SRISWQ---GDPCVPLQFLWDGLNCS
 AT5G59670/1-499 347 NALEVYTVIQFPRSETDES DVVAMKNISASYGL-SRINWQ---GDPCFPQQLRWDALDCT
 AT5G59680/1-508 347 NAYEVYKVIQFPQOETNEDDVSAVKNIQAT YEL-SRINWQ---SDPCVPPQFMWDGLNCS
 AT5G59650/1-509 347 NALEIYTVIQFPRNQLSSTSVVAVKNIEAAYKL-SRIRWQ---GDPCVPPQYAWDGLNCS
 AT5G16900/1-505 342 NAMEAFTAIFPHSETNPDVIAIKNIQATYEL-SRVDWQ---GDPCLPQQLWDTGLNCS
 AT1G07560/1-512 349 NAIEAFITVEFPQSETNANDVLAIKS IETSYGL-SRISWQ---GDPCVPPQLLWDGLTCE
 AT4G20450/1-530 359 NAMEVFGLLQLLQOETEDNDVTTLKNIQATYRI-QKTNWQ---GDPCVPIQFIWTGLNCS
 AT2G28960/1-507 343 SAIEAFKVVDFPYAETNPNDVAA MKDIEAFYGL-KMISWQ---GDPCVPELLKWEGLKCS
 AT2G29000/1-506 343 NAYEVYILVEFPYSETHPDVVAIKKIKAA YGL-KIISWQ---GDPCLPREYKWEYIECS
 AT2G28970/1-409 245 NAIEAYTIEFPQLETSLSDVNAIKNIKATYRL-SKTSWQ---GDPCLPQELSWENLRCS
 AT2G28990/1-506 342 NAIEAYSVIEFSQLETSLSDVDAIKNIKN TYKL-NKITWQ---GDPCLPQDLSWESIRCT
 AT1G49100/1-518 350 NAIELEFTVVEFPQSETNQDEVI AIKKIQLTYGL-SRINWQ---GDPCVPEQFLWAGLKCS
 AT1G51810/1-384 214 NAMEAYTVLDFFQIETNVDEVI AIKNIQSTYGL-SKTTWQ---GDPCVPKFLWDLNCS
 AT1G51805/1-504 340 NAIEAFTVIDFLQVETDEDDAAAIKNVQ NAYGLINRSSWQ---GDPCVPKQYSWDGLKCS
 AT1G51830/1-315 150 NAIEAFTVIDFPQMETNEDDVTGINDVQNTYGL-NRISWQ---GDPCVPKQYSWDGLNCS
 AT1G51820/1-504 339 NAIEAFTVIDFPQMETNENDVAGIKNVQGT YGL-SRISWQ---GDPCVPPQLLWDGLNCK
 AT1G51850/1-486 319 NAIEAFTVIDFPQMETNGDDVDAIKNVQD TYGI-SRISWQ---GDPCVPKFLWDLNCS
 AT2G04300/1-480 312 -ALEVFTVIDFPELETNQDDVIAIKNIQNTYGV-SKTSWQ---GDPCVPPKRFMWDLNCS
 AT3G21340/1-520 351 NALEVFTVIDFPQMETNPDVVAIKSIQSTYGL-SKISWQ---GDPCVPKQFLWEGLNCS
 AT1G05700/1-507 346 NALEIYVANSFSQSLTNQEDGDAVTS LKTSYKV-KK-NWH---GDPCLPNDYIWEGLNCS
 AT2G19210/1-516 352 NAIEYRVNEFLQSPDQDDVDAIMRIKSKYGV-KK-SWL---GDPCAPVKYPWKDINCS
 AT2G19230/1-516 353 NAIEYRTNEFLDLPTDQNDVDAIMKIKTKYKV-KK-NWL---GDPCAPFGYPWQGINCS
 AT2G19190/1-516 352 NALEVYQVNEFLQIPTHPQDAMDAMRRIKATYRV-KK-NWQ---GDPCVVDYSWEGIDCI
 AT4G29990/1-511 347 NAIEIYQINEFLQLPTDQDDVDAMTKIKFKYRV-KK-NWQ---GDPCVVDNSWEGLECL
 AT4G29180/1-509 352 NAIEIFSAQSLDEFYTRIDDVQAIESIKSTYKV-NKI-WT---GDPCSPRLFPWEGIGCS
 AT4G29450/1-513 353 NAIEIFTAQSLEDFSTTIEDIHAIESIKATYKV-NKV-WS---GDPCSPRLFPWEGVGCS

LjSYM RK/1-509 392 --GSNGSSVITKLDLSSSNLKGILPSSIAEMTNLETLNISHNSFDG-SVPSFLLS--LI
 LjShRK1/1-522 402 ---SDIQPRIVSILSSKNLTGNIPLDITKLTGLVELWLDGNMLTG-PIPDF-TGCM DLK
 LjShRK2/1-506 400 YAESD-SPRIIYLNLS SSGLTGNIAPSI SNMKSIEYLDLSNNLTG-ALPDFLSQLRFLR
 ShRK1_ (AT1/1-533 410 ---S TSPRVTKIALSRKNLRGEIPPGIN YMEALTE LWLDDNE LTG-TLPDM-SKLVNLK
 ShRK2_ (AT2/1-529 410 ---SDQP RVVAIKLSSMNL TGNI PSDLVKLTGLVELWLDGNSFTG-PIPDF-SRCPNLE
 AT5G48740/1-492 383 -----GNRVTSLDLHNTSLTGAIQSELEDLVNLEVL D LQNNSLQG-SVPE TLGK LK LKLR
 AT1G51790/1-518 405 -FNGT NMPRVIALNLS SAGLTG EITSDI SRLSQLQILDLSNNLSGPAVPAFLAQLQFLR
 IOS1 (AT1G5/1-513 402 YVDNE-TPKII SLDLSTSGLTGEILEFISDLTSL EVLDLSNNSLTG-SVPEFLANMETLK
 AT1G51910/1-508 405 YLD-SDQPLITSLNLR TSGLTG IITHDISNLIQLRELDLSDNLSG-EI PDFLADMKMLT
 AT1G51890/1-499 400 YPN-FAPPQIISLNLSSGNSLGTITSDISKLTHLRELDLSNNLSG-DIPFVFS DMKNLT
 AT1G51860/1-511 405 YPD-SEGSRIISLNLNGSGLTGTITSDISKLTLTVLDLSNNLSG-DIPFFAEMKSLK
 AT1G51880/1-511 404 YPN-SDQPRIISLNLAEKLTGTITPEISKLTQLIELDLSKNLSG-EIPEFFADMKLLK
 AT1G07550/1-504 403 VIDVSTPPRIKLDLSSSGLNGVI PPSIQNLTLQELDLSQNNLTG-KVPEFLAKMKYLL
 AT2G14440/1-502 405 VIDISTPPRIISL DLS SSGLTGVI T P S I Q N L T M L R E L D L S N N N L T G - E V P E F L A T I K P L L
 AT2G14510/1-508 404 VIDISTPPRIISL DLS SGLTGVISPSIQNLTLRELDLSNNLTG-EVPEFLATIKPLL
 AT3G46350/1-491 382 DTDYIAPRITSLKLSKGLTGTIAADIQYLTSL EKLDLSDNKLVG-VVPEFLANMKSLM
 AT3G46340/1-513 406 SKDALTLPRITSLNLSSTGLTGNIAAGIQNLTHL DKLDLSNNLTG-GVPEFLASMKLS
 AT3G46370/1-427 315 DTNVSTPPRIISLNLSSSGLTGN IATG IQNLTKLQKLDLSNNLTG-VVPEFLANMKSL
 AT3G46400/1-508 404 DKNVSAPRITSLNLS SGLTGVIPSGIQNF TLEKLDLSNNLTG-LVPEFLAKMKSL
 AT3G46330/1-516 405 ITDISAPPRIISLNLSSSGLSGTIVSNFQNL AHLES LDLSNNSLSG-IVPEFLATMKSL
 AT5G59670/1-499 403 NRNISQPPRITSLNLS S R L N G T I A A A I Q S I T Q L E T L D L S Y N N L T G - E V P E F L G K M K S L S
 AT5G59680/1-508 403 ITDITTPPRIITLNLSSSGLTGTITAAIQNLTL EKLDLSNNLTG-EVPEFLSNMKSL
 AT5G59650/1-509 403 NTDVSKPPRVLSLNLSSSGLTGI AAAIQNLTHLEKLDLSNNLTG-VVPEFLAQMKSLV
 AT5G16900/1-505 398 YMNMTSPRIISL DLS S H K L T G R I V P D I Q N L T Q L Q K L D L S N N K L T G - G V P E F L A N M K S L L
 AT1G07560/1-512 405 YTNMSTPPRIHSLDLSSELTGIVPEIQNLTELK K L D F S N N N L T G - G V P E F L A K M K S L L
 AT4G20450/1-530 415 NMFPSIPPRITSIDFSNGLNGTITSDIQYLNQLQKLDLSNNLTG-KVPEFLAKMKMLT
 AT2G28960/1-507 399 YTNKSTPPRIISL DLS S R L K G V I A P A F Q N L T E L R K L D L S N N S F T G - G V P E F L A S M K S L S
 AT2G29000/1-506 399 YTNNSIPPRII SL D L S N R G L K G I I E P V L Q N L T Q L E K L D L S I N R L S G - E V P E F L A N M K S L S
 AT2G28970/1-409 301 YTNSS T P P K I I S L N L S A S G L T G S L P S V F Q N L T Q I Q E L D L S N N S L T G - L V P S F L A N I K S L S
 AT2G28990/1-506 398 YVDGSTSPITISL D L S K S G L N G S I P Q I L Q N F T Q L Q E L D L S N N S L T G - P V P I F L A N M K T L S
 AT1G49100/1-518 406 NINSSTPPTITFLNLSSSGLTGIISPSIQNLTHLQELDLSNNDLTG-DVPEFLADIKSLL
 AT1G51810/1-384 270 NSDDSTPPIITSLNLSSSGLTGIIVLTIQNLANLQELDLSNNLSG-GVPEFLADMKSLL
 AT1G51805/1-504 397 YSD-STPPIINFLDLSASGLTGIIPAIQNLTHLEILALSNNLTG-EVPEFLADLKSIM
 AT1G51830/1-315 206 NSDISIPPIIISL DLS S S G L N G V I T Q G I Q N L T H L Q Y L D L S D N N L T G - D I P K F L A D I Q S L L
 AT1G51820/1-504 395 NSDISTPPIITSL DLS S S G L T G I I T Q A I K N L T H L Q I L D L S D N N L T G - E V P E F L A D I K S L L
 AT1G51850/1-486 375 NSDNSTSPITISL DLS S S G L T G S I T Q A I Q N L T N L Q E L D L S D N N L T G - E I P D F L G D I K S L L
 AT2G04300/1-480 367 NSYISTPPTITFLNLSSSHLTGIIASA IQNLTHLQNL D L S N N N L T G - G V P E F L A G L K S L L
 AT3G21340/1-520 407 NLDNSTPPIVTSLNLSSSHLTGIIAAGIQNLTHLQELDLSNNLTG-GIPEFLADIKSLL
 AT1G05700/1-507 401 Y-DSLTPPRITSLNLSSSGLTGHIS S S F S N L T M I Q E L D L S N N L T G - D I P E F L S K L K F L R
 AT2G19210/1-516 407 YVDNE-SPRIISVNLSSSGLTGEIDA A F S N L T L L H I L D L S N N S L T G - K I P D F L G N L H N L T
 AT2G19230/1-516 408 YTANN-PPRIISVNL S F S G L T G Q I D P V F I T L T P L Q K L D L S N N R L T G - T V P D F L A N L P D L T
 AT2G19190/1-516 407 QSDNTTNPRVSLNLSFSELRGQIDPAFNSLTSIRKLDLSGNTLTG-EIPAFLANLPNLT
 AT4G29990/1-511 402 HSDNNTSPKSIALNLS S S G L T G Q I D P A F A N L T S I N K L D L S N N S L T G - K V P D F L A S L P N L T
 AT4G29180/1-509 407 --YNTSSYQIKSLNLSSSGLHGPIAFAFRNLSLLES LDLSNNLKG-IVPEFLADLKYLK
 AT4G29450/1-513 408 --DNNNNHQIKSLNLSSSGLLGPVLAFRNLSLLES LDLSNNDLQQ-NVPEFLADLKHK

LjSYMRK/1-509 447 SVDLSYN-DLM-GKLPESIVKL---PHLKSLYFGCNEHMSPEDPANMNSSLINTDYGRCK
 LjShRK1/1-522 457 IHLNENN-QFS-GALPTSLVNL---PKLRELWVQNNMLSGTVPSDLKSLVLVNYSGNVK
 LjShRK2/1-506 458 VLNLEGN-QLS-GTIPMPLTVRSKNDLL-ESNFGGN-----PDLCSPGSCN--
 ShRK1_ (AT1/1-533) 465 IMHLENN-QLS-GSLPPYLAHL---PNLQELSIENNSEFKGKIPSAALLKGVLFKYNNNPE
 ShRK2_ (AT2/1-529) 465 IHLNENN-RLT-GKIPSSLTKL---PNLKELYLQNNVLTGTIPSD-LAKDVISNFSGNLN
 AT5G48740/1-492 436 LLNLENN-NLV-GPLPQSL---NITGL-EVRI TGN-----PCLSFSSSCNNV
 AT1G51790/1-518 464 VLHLANN-QLS-GPIPSLIER-----LDSFSGN-----PSICSANACEEV
 IOS1 (AT1G5/1-513) 460 LINLSGN-ELN-GSIPATLLDKERRGSI-TLSIEGN-----TGLCSSTSCA--
 AT1G51910/1-508 463 LVNLKGNPKLN-LTVPDSIKHRINNKS L-KLIIDEN-----Q----SSE
 AT1G51890/1-499 458 LINLSGNKLN-RSVPETLQKRIDNKS L-TLIRDE-----TGK
 AT1G51860/1-511 463 LINLSGNPNLNLTAIPDSLQQRVNSKSL-TLILGEN-----L--TLTPK
 AT1G51880/1-511 462 LINLSGNLGLN-STIPDSIQRLDSKSL-ILILSKT-----VTKTVTLK
 AT1G07550/1-504 462 VINLSGN-KLS-GLVPQALLDRK-EGE-KLLV--D-----ENMICVSCGTRF
 AT2G14440/1-502 464 VIHLRGN-NLR-GSVPQALQDRENNNDGL-KLL-----RGKHQ
 AT2G14510/1-508 463 VIHLRGN-NLR-GSVPQALQDRKNDGL-KLFV--D-----PNI--TRRGKHQ
 AT3G46350/1-491 441 FNLTKN-DLH-GSIPQALDREREK-KGL-KILFDGDK-----NDPCLSTSCN--
 AT3G46340/1-513 465 FINLSKN-NLN-GSIPQALLKREK-DGL-KLSVDE-----QIRCFPGSCV--
 AT3G46370/1-427 374 FIDLKRN-KLN-GSIPKTLDRK-KGL-QLFVDGDDDK-----GDDNKCLSGSCV--
 AT3G46400/1-508 463 FIDLKRN-KLN-GSIPQALLDRK-KGL-QLFVDG-----PDLCLSTSCV--
 AT3G46330/1-516 464 VINLSGN-KLS-GAIPQALRDREK-EGE-KLNVLGN-----KELCSSSTCIDK
 AT5G59670/1-499 462 VINLSGN-NLN-GSIPQALR---K-KRL-KLYLEGN-----PRL-IPPKK--
 AT5G59680/1-508 462 VINLSGN-DLN-GTIPQSLQ---R-KGL-ELLYQGN-----PRL-ISPSTET
 AT5G59650/1-509 462 IINLSGN-NLS-GPLPQGLR---R-EGE-ELLYQGN-----PRLCLSGSCTEK
 AT5G16900/1-505 457 FINLSNN-NLV-GSIPQALLDR---KNL-KLEFEGN-----PKLCATGPCNNS
 AT1G07560/1-512 464 VINLSGN-NLS-GSVPQALLNKVK-NGL-KLNIQGN-----PNLCFSSSCNKK
 AT4G20450/1-530 474 FINLSGN-NLS-GSIPQSLNMEK-NGLITLLYNGN-----NLCLDPSCSE
 AT2G28960/1-507 458 IINLNWN-DLT-GPLPKLLLDREK-NGL-KLTIQGN-----PKLCNDASCNN
 AT2G29000/1-506 458 NINLSWN-NLK-GLIPPALEEKRK-NGL-KLNTQGN-----QNLCPGDECKRS
 AT2G28970/1-409 360 LLDLSGN-NFT-GSVPQTLDRK-EGE-VLKLEGN-----PELCKFSSCNPK
 AT2G28990/1-506 457 LINLSGN-NLS-GSVPQALLDREK-EGE-VLKLEGN-----PDLCKSSFCNTE
 AT1G49100/1-518 465 IINLSGN-NFS-GQLPKLIDK---KRL-KLNVEGNP-----KLLCTKGPCGNK
 AT1G51810/1-384 329 VINLSGN-NLS-GVVPQKLIK---KML-KLNIEGNP-----KLNCTVESC VNK
 AT1G51805/1-504 455 VIDLRGN-NLS-GVPASLLQK---KGL-MLHLDDNP-----HILCTTGSCMHK
 AT1G51830/1-315 265 VINLSGN-NLT-GSVPPLLLQK---KGL-KLNVEGNP-----HLLCTDGLCVNK
 AT1G51820/1-504 454 VINLSGN-NLS-GSVPPLLLQK---KGM-KLNVEGNP-----HILCTTGSCVKK
 AT1G51850/1-486 434 VINLSGN-NLS-GSVPPLLLQK---KGM-KLNVEGNP-----HLLCTADSCVKK
 AT2G04300/1-480 426 VINLSGN-NLS-GSVPQTLQK---KGL-KLNLEGNI-----YLNCPDGSCVSK
 AT3G21340/1-520 466 VINLSGN-NFN-GSIPQILLQK---KGL-KLILEGNA-----NLICPDGLCVNK
 AT1G05700/1-507 459 VLNLENN-TLT-GSVPSELLERSNTGSF-SLRIGEN-----PGLCTEISCR--
 AT2G19210/1-516 465 ELNLEGN-KLS-GAIPVKLLERSNKLI-LLRIDGN-----PDLCVSASCQIS
 AT2G19230/1-516 466 ELNLEEN-KLT-GILPEKLLERSKDGSL-SLRVGGN-----PDLCVSDSCR--
 AT2G19190/1-516 466 ELNVEGN-KLT-GIVPQLHERSKNGSL-SLRFGRN-----PDLCLSDSCS--
 AT4G29990/1-511 461 ELNLEGN-KLT-GSIPAKLLEKSKDGSL-SLRFGGN-----PDLQSPSCQ--
 AT4G29180/1-509 464 SLNLKGN-NLT-GFIPRSLRKRATANGL-ALSVDE-----QNIHSRSCRDR
 AT4G29450/1-513 465 VLNLKGN-NFT-GFIPKSLMKKLAGLL-TLSADE-----QNLN--SCQEK

LjSYMRK/1-509	502	GKESRFGQ-----
LjShRK1/1-522	512	LHKGSRRKSHM-----
LjShRK2/1-506	501	QKNGNK-----
ShRK1_(AT1/1-533	520	LQNEAQRKHFQIL-
ShRK2_(AT2/1-529	519	LEKSGDKGKKL-----
AT5G48740/1-492	477	SSTIDTPQVTIPINKK
AT1G51790/1-518	503	SQNRSKKNKLPSFVIP
IOS1(AT1G5/1-513	503	--TTKKKKKNTVI---
AT1G51910/1-508	501	KHGKIFPL-----
AT1G51890/1-499	494	NSTNVV-----
AT1G51860/1-511	504	KESKKVPM-----
AT1G51880/1-511	504	GKSKKVPM-----
AT1G07550/1-504	504	P-----
AT2G14440/1-502	498	PKSWL-----
AT2G14510/1-508	504	PKSWL-----
AT3G46350/1-491	484	-PKKKFSVM-----
AT3G46340/1-513	506	ITKKKFPV-----
AT3G46370/1-427	421	-PKMKFPL-----
AT3G46400/1-508	502	-PKNKFP-----
AT3G46330/1-516	508	PKKKVAVKV-----
AT5G59670/1-499		-----
AT5G59680/1-508	502	KSGKSFP-----
AT5G59650/1-509	503	NSKKKFP-----
AT5G16900/1-505	499	SGNKETT-----
AT1G07560/1-512	508	KNSIM-----
AT4G20450/1-530	518	TGPGNNKKLLVP---
AT2G28960/1-507	502	NNQTYI-----
AT2G29000/1-506	502	IPKFP-----
AT2G28970/1-409	404	KKKGLL-----
AT2G28990/1-506	501	KKNKFL-----
AT1G49100/1-518	508	PGEKGHPKCSI-----
AT1G51810/1-384	372	DEEGGRQIKSMTI---
AT1G51805/1-504	498	G-EGEKKS-----
AT1G51830/1-315	308	G-DGHKKKS-----
AT1G51820/1-504	497	KEDGHKKK-----
AT1G51850/1-486	477	GEDGHKKKSV-----
AT2G04300/1-480	469	DGNGGAKKKNV-----
AT3G21340/1-520	509	AGNGGAKKMN-----
AT1G05700/1-507	502	KSNSKK-----
AT2G19210/1-516	510	DEKTKKN-----
AT2G19230/1-516	509	NKKTERKE-----
AT2G19190/1-516	509	--NTKKKNKN-----
AT4G29990/1-511	504	-TTTKKKIG-----
AT4G29180/1-509	508	NR-----
AT4G29450/1-513	507	KKKKSMV-----

Appendix 3: Host-related metabolic cues affect colonization strategies of a root endophyte

Lahrman U, Ding Y, **Banhara A**, Rath M, Hajirezaei MR, Dohlemann S, Von Wiren N, Parniske M & Zuccaro A (2013) Host-related metabolic cues affect colonization strategies of a root endophyte. *Proc Natl Acad Sci U S A* **110**:13965-70. doi: 10.1073/pnas.1301653110.

Contributions by Aline Banhara:

- Designed, performed, and analysed all experiments concerning growth promotion of *A. thaliana* by *Piriformospora indica* on plants grown on a modified Hoagland's medium;
- Analysed results of *P. indica* on biomass of plants grown on 1/10 PNM medium;
- Prepared the corresponding Figure S4.

Host-related metabolic cues affect colonization strategies of a root endophyte

Urs Lahrmann^{a,1}, Yi Ding^{a,1}, Aline Banhara^b, Magnus Rath^c, Mohammad R. Hajirezaei^d, Stefanie Döhlemann^a, Nicolaus von Wirén^d, Martin Parniske^b, and Alga Zuccaro^{a,2}

^aMax-Planck-Institut für terrestrische Mikrobiologie, D-35043 Marburg, Germany; ^bLudwig-Maximilians-Universität München, D-82152 Martinsried, Germany; ^cSpezielle Botanik und Mykologie, Philipps-Universität Marburg, D-35032 Marburg, Germany; and ^dLeibniz-Institut für Pflanzengenetik und Kulturpflanzenforschung, OT Gatersleben, D-06466 Stadt Seeland, Germany

Edited by Paul Schulze-Lefert, Max Planck Institute for Plant Breeding Research, Cologne, Germany, and approved July 8, 2013 (received for review January 25, 2013)

The mechanisms underpinning broad compatibility in root symbiosis are largely unexplored. The generalist root endophyte *Piriformospora indica* establishes long-lasting interactions with morphologically and biochemically different hosts, stimulating their growth, alleviating salt stress, and inducing local and systemic resistance to pathogens. Cytological studies and global investigations of fungal transcriptional responses to colonization of barley and *Arabidopsis* at different symbiotic stages identified host-dependent colonization strategies and host-specifically induced effector candidates. Here, we show that in *Arabidopsis*, *P. indica* establishes and maintains biotrophic nutrition within living epidermal cells, whereas in barley the symbiont undergoes a nutritional switch to saprotrophy that is associated with the production of secondary thinner hyphae in dead cortex cells. Consistent with a diversified trophic behavior and with the occurrence of nitrogen deficiency at the onset of saprotrophy in barley, fungal genes encoding hydrolytic enzymes and nutrient transporters were highly induced in this host but not in *Arabidopsis*. Silencing of the high-affinity ammonium transporter *PiAMT1* gene, whose transcripts are accumulating during nitrogen starvation and in barley, resulted in enhanced colonization of this host, whereas it had no effect on the colonization of *Arabidopsis*. Increased levels of free amino acids and reduced enzymatic activity for the cell-death marker VPE (vacuolar-processing enzyme) in colonized barley roots coincided with an extended biotrophic lifestyle of *P. indica* upon silencing of *PiAMT1*. This suggests that *PiAMT1* functions as a nitrogen sensor mediating the signal that triggers the *in planta* activation of the saprotrophic program. Thus, host-related metabolic cues affect the expression of *P. indica*'s alternative lifestyles.

root cortical cell death | RCD | broad-host range | biotrophy | mutualism

Upon plant colonization, fungi adopt different strategies to gain access to host nutrients. Whereas necrotrophs kill plant cells with subsequent saprotrophic nutrition, other fungi maintain biotrophic relationships with their hosts either transiently (hemibiotrophs) or as lifelong interactions. The degree of specialization to a particular host and the host's metabolic status may greatly influence plant colonization (1–4). Broad-host range root endophytes undergo long-term interactions with a large variety of plants, thereby playing a significant role in natural and managed ecosystems and in the evolution of land plants. To establish and maintain a compatible interaction with different hosts, these endophytes must respond and adapt to host-specific signals. Alternative lifestyles and colonization strategies in different host species thus may be a consequence of this adaptation to highly variable host environments. In this study, we addressed the question of whether endophytes adopt different strategies during colonization of distinct hosts or whether their success resides in a general colonization strategy. An interesting system to explore this issue is the mutualistic root endophyte *Piriformospora indica* (Basidiomycota, Sebaciniales), with its large number of plant hosts. Among others, this generalist can establish a mutualistic interaction with roots of the agriculturally important monocot barley (*Hordeum vulgare*) and the dicot model

plant *Arabidopsis thaliana* (5, 6), two biochemically and morphologically distinct plants. Cytological studies in both hosts have shown that *P. indica* has a biphasic colonization strategy (7–10). Initial root cell invasions are biotrophic where colonized host cells maintain membrane integrity and invasive hyphae of *P. indica* remain enveloped by the host plasma membrane and thus are not accessible to cell wall stains such as wheat germ agglutinin–Alexa Fluor 488 (WGA-AF488) conjugate (7, 8, 11). Later, *P. indica* is found more often in dead or dying host cells (9, 10), especially in the root cortex of barley. Colonization at later stages was shown to be reduced by overexpression of the negative cell death regulator BAX inhibitor 1 in barley and to be mediated by an endoplasmic reticulum stress-triggered caspase-dependent cell death in *Arabidopsis* (9, 10). Questions arise as to what extent hemibiotrophy in a root endophyte reflects a general colonization strategy of the symbiont to benefit from different plants and how the symbiotic lifestyle is influenced by the hosts. Our study reveals that broad compatibility in a root endophyte is associated with phenotypic plasticity of the symbiont and with the expression of alternative lifestyle strategies in a host-dependent way.

Results

Expression Patterns of *P. indica* Genes, Including Those Encoding Small Secreted Proteins, Support a Diversified Colonization Strategy for Barley and *Arabidopsis*. We hypothesized that successful root colonization of different hosts would require host-related colonization strategies, and we studied these differences by a global characterization of fungal transcriptional responses to barley and *Arabidopsis* at different developmental stages. A customized Agilent microarray was designed to monitor *P. indica*'s gene expression during root colonization of plants grown on sugar-free minimal medium (PNM) at 3 (early biotrophic phase) and 14 (late saprotrophic phase) days post inoculation (dpi). Fungal material grown on PNM was used as a control. From the 11,463 *P. indica* genes represented on the microarray chip, about 2,400 genes were differentially regulated at the early biotrophic phase in barley compared with the control. At this time point, 3,162 fungal genes were differentially regulated in colonized *Arabidopsis* compared with the control. At the late colonization stage, far more genes (4,482 genes) were differentially regulated in

Author contributions: N.v.W., M.P., and A.Z. designed research; U.L., Y.D., A.B., M.R., M.R.H., S.D., and A.Z. performed research; U.L., Y.D., and A.Z. analyzed data; and U.L., N.v.W., and A.Z. wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission.

Freely available online through the PNAS open access option.

Data deposition: The data reported in this paper has been deposited in the Gene Expression Omnibus (GEO) database, www.ncbi.nlm.nih.gov/geo (accession no. GSE47775).

¹U.L. and Y.D. contributed equally to this work.

²To whom correspondence should be addressed. E-mail: zuccaro.alga@mpi-marburg.mpg.de.

This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10.1073/pnas.1301653110/-DCSupplemental.

colonized barley than in colonized *Arabidopsis* roots (1,948 genes; Dataset S1). In total, about 70% (2,023) of the *in planta*-induced genes (2,861 genes) encoding intracellular proteins and 60% (277 of 463 induced genes) of those encoding putative secreted proteins had host-specific expression profiles (SI Appendix, Fig. S1). In particular, 123 of the 216 induced genes encoding small secreted proteins (SSPs; <300 amino acids), also known as putative effectors (12), were either *Arabidopsis* or barley responsive (Fig. 1A), suggesting that colonization of different hosts may require exploitation of distinct effectors that can interact with elements characteristic to each host. SSPs recently were shown to facilitate colonization by manipulating host defense and reprogramming plant metabolism during symbiosis (13, 14). Many of these proteins are cysteine rich (15) or possess distinctive features, such as a regular pattern of histidine and alanine residues found in all members of the *P. indica*-specific DELD family (7, 8). Eighteen of the 29 genes encoding *P. indica* DELD proteins were plant responsive and largely induced in barley but to a lesser extent in *Arabidopsis* (Fig. 1B and Dataset S2). Additionally, a small set of 21 SSPs were identified that are expressed in both hosts at comparable symbiotic stages (Fig. 1A). These SSPs may represent general determinants that target conserved recognition and signaling pathways in roots. Congruent with the broad definition of effectors (12), at the early symbiotic stage, 16 and 14 of *P. indica*'s top 20 up-regulated SSPs in *Arabidopsis* and in barley, respectively, encoded proteins with no known functional domains. A different situation was found at 14 dpi, when 50% of *P. indica*'s SSPs induced during colonization of barley, but not of *Arabidopsis*, encoded putative hydrolases (mainly glycoside hydrolase families GH10, GH11, and GH61; Dataset S3). Taken together, these data are consistent with a diversified colonization strategy for barley and for *Arabidopsis*, especially at 14 dpi, and prompted us to further characterize the two interactions at this stage.




Fig. 1. Host-dependent expression profiles of *in planta*-induced *P. indica* genes encoding SSPs (<300 amino acids). (A) Number of SSPs significantly induced at 3 and 14 dpi during *Arabidopsis* (PI_AT) and barley (PI_HV) colonization, calculated vs. PNM control. (B) RT-qPCR analysis of *PiDLD1* (PiIN_05872), a member of the *P. indica*-specific putative effector family DELD, during colonization of *Arabidopsis* (Left) and barley (Right) at different time points from three independent experiments.

***P. indica* Undergoes Major Trophic, Phenotypic, and Transcriptional Rearrangements During Colonization of Barley, Whereas It Maintains a Predominant Biotrophic Nutrition in *Arabidopsis*.** Transcriptional data, together with cytological analyses, show that in barley, in response to a progressively increasing natural root cortical cell death (RCD) in older/basal root zones (16), *P. indica* undergoes a distinct nutritional shift from biotrophic to saprotrophic nutrition associated with the production of secondary thinner hyphae and with the secretion of hydrolytic enzymes (Fig. 2A–D and SI Appendix, Fig. S2). Thus, after biotrophic colonization of the outermost cell layer (3–5 dpi), on its route toward the endodermis, the fungus progresses inter- and intracellularly by digestion of barley cortical cell walls, which became most evident at 30 dpi (Fig. 2C), but without visible root necrosis at the macroscopic level (10, 17). Host cell wall appositions, named papillae, often are visible beneath the site of fungal penetration attempts of living host barley cells (8), and at later colonization stages, hyphal contact with the endodermis, which is not penetrated (10), results in a strong autofluorescence of the host cell wall, indicating activation of plant defense upon attempted penetration of the root vasculature (SI Appendix, Fig. S3). Despite a predominant saprotrophic lifestyle, the presence of *P. indica* in barley leads to beneficial effects (6, 17).

Colonization of *Arabidopsis* by *P. indica* results in growth promotion (5) (SI Appendix, Fig. S4) and is characterized by a long-term feeding relationship with living host cells via the production of thicker bulbous invasive hyphae in epidermal cells and no sign for papillae formation upon penetration (Fig. 2F and G and SI Appendix, Fig. S5). We show here that these intracellular, non-WGA-stainable multilobed invasive hyphae are present throughout the *Arabidopsis* colonization process, leading to a nondestructive progression within the epidermis and cortex layers, as demonstrated by colonized living root cells capable of endocytosis also at later time points (Fig. 2F). Consistent with this observation is the reduced expression of *P. indica* genes involved in host cell wall and lipid degradation at 3 and 14 dpi in this host compared with the situation in barley (Fig. 2D and SI Appendix, Fig. S2). The occurrence of a long-term biotrophic nutrition in *Arabidopsis* and of a switch to a saprotrophic nutrition in barley is supported further by comparative expression analyses of fungal genes involved in primary metabolism and nutrient transport. Whereas in colonized *Arabidopsis* fungal transcripts for amino acid biosynthetic processes and glycolysis are abundant, they show lower expression values in colonized barley at 14 dpi (SI Appendix, Fig. S6 and Dataset S4A). Conversely, at this time point, transcripts encoding fungal carbohydrate and nitrogen transporters are strongly induced during colonization of barley (Fig. 2E and SI Appendix, Fig. S7 and Dataset S4B). In particular, the high-affinity ammonium transporter PiAmt1 (PiIN_02036), whose transcripts are accumulating upon nitrogen starvation in axenic culture (SI Appendix, Fig. S8A), is induced during the late saprotrophic phase in colonized barley but to a much lower extent in colonized *Arabidopsis* (SI Appendix, Fig. S9A and B), strongly indicating that in barley, but not in *Arabidopsis*, a status of nitrogen depletion is reached.

Transcripts for *P. indica* ABC transporters and other transporters most probably implicated in detoxification processes are well represented in both hosts, but to a greater extent in colonized *Arabidopsis* at 14 dpi (SI Appendix, Figs. S2 and S10). ABC transporters are efflux pumps that have been implicated in resistance to antifungal compounds in various host–fungal interactions, but little is known about their substrate specificities (18–20). Their host-dependent expression profiles therefore might represent a fungal stress response matched to distinct phytoalexins and other antifungal compounds produced by different hosts (21).

The Switch to Saprotrophic Nutrition in Barley Is Affected by Nitrogen Availability in a PiAMT1-Dependent Manner. To address whether nitrogen availability is one of the key switches to saprotrophic nutrition *in planta*, *P. indica* RNAi strains carrying a silencing




Fig. 2. Colonization patterns and *P. indica* gene expression polymorphisms correlate with extended biotrophy in *Arabidopsis* (AT) and with a switch from biotrophic to saprotrophic nutrition during colonization of barley (HV). (A) Maximum projection of a barley root colonized by *P. indica* at 30 dpi. Broad extraradical hyphae are visible at the boundary of the epidermis, whereas thin secondary hyphae are filling the cortical cells. Host nuclei are absent in the cortex cells while the cylinder is undamaged and preserves intact nuclei. The root was stained with acid fuchsin. (B) Closer view of biotrophic broad invasive hyphae within a barley epidermal cell and secondary hyphae in cortical cells (*). (C) Transverse 4- μ m sections of barley roots inoculated with *P. indica* (30 dpi), stained with toluidine blue. (Upper) Heavily colonized cortex cells. (Lower) A noncolonized part of the root. (D and E) Heat map showing \log_2 -fold expression changes of *P. indica* genes encoding (D) hydrolytic enzymes and (E) transporters. Significant (t test, $P < 0.05$) \log_2 -fold expression changes were calculated vs. PNM control. A consistent divergence is observed at 14 dpi, where a strong induction is visible for most of these genes during colonization of barley only (for a detailed overview, see Dataset S1). ¹Raw expression data for barley 3 dpi were retrieved from ref. 8. Color coding indicates up-regulation (red) and down-regulation (blue) of genes. (F) Biotrophic broad invasive hyphae (white *) in *Arabidopsis* epidermal cell at 14 dpi. In contrast to extracellular hyphae, invasive hyphae are not stainable with WGA-AF488 (green) because of the presence of a plant-derived membrane. Endomembrane structures stained by FM4-64 (red) are visible inside the plant cells, indicating cell viability. (G) *Arabidopsis* epidermal cell with biotrophic invasive hyphae (white *) of *P. indica* GFP strain (14 dpi). The scale bars represent 25 μ m.

construct targeting the high-affinity ammonium transporter *PiAMT1* were generated. The success of transformation for selected *P. indica* RNAi strains was confirmed by Southern blot, and the efficiency of silencing was verified by quantitative PCR (qPCR) experiments (Fig. 3A and SI Appendix, Fig. S11A). *PiAMT1* displays strong homology (75% amino acid residues identity) to the high-affinity ammonium transporters AMT1 and AMT2 from *Hebeloma cylindrosporium* and to the high-affinity ammonium permease MEP2 from *Saccharomyces cerevisiae* (SI Appendix, Fig. S8B). These transporters were proposed to function as ammonium sensors, generating downstream signals in response to nitrogen starvation (22, 23). In yeast, MEP2 is

required for pseudohyphal growth, which is the outgrowth of nuclear-free hyphae to better forage the medium at low N availability (22, 24, 25). The ammonium import function of *PiAMT1* was verified by yeast complementation (SI Appendix, Fig. S8C), and the predicted topological structure of the *P. indica* Amt1 polypeptide demonstrated the presence of a long cytoplasmic tail at the C-terminus, which might be involved in downstream signaling (SI Appendix, Fig. S8D). To verify that ammonium uptake by the *P. indica* RNAi strains is reduced relative to wild-type (WT) and empty-vector (EV) controls, these strains were analyzed for growth on minimal medium with a low concentration of ammonium as the sole nitrogen source. Their growth on complex




Fig. 3. Silencing of *PiAMT1* affects *P. indica* colonization of barley but not of *Arabidopsis*. (A) Relative expression of *PiAMT1* in *P. indica* WT and EV controls and RNAi strains grown in liquid complete medium (CM) for 7 d. **ANOVA, $P < 0.01$. (B) Colony phenotype of *P. indica* WT and transformants on yeast nitrogen base medium supplemented with 2 mM NH_4Cl as the sole nitrogen source, 14 dpi. (C and D) Relative amount of fungal DNA in (C) barley and (D) in *Arabidopsis* roots colonized by *P. indica* WT or transformants at 14 dpi. Plants were grown on 1/10 PNM. Error bars represent SE of the mean from three independent biological repetitions. Grouping was done by ANOVA. *Southern blot analyses showed that all transformed strains had one to two integrations of the plasmids, with the exception of EV2, which had multiple integrations (SI Appendix, Fig. S11).

media and media supplied with large amounts of ammonium was not affected (SI Appendix, Fig. S11B), whereas hyphal growth at low N was suppressed (Fig. 3B). The RNAi strains were investigated further for altered root colonization. Despite a substantial reduction in growth under a low supply of ammonium as the sole nitrogen source and the absence of any evidence for a compensatory

up-regulation of the second ammonium transporter *PiAMT2* (SI Appendix, Fig. S9C), the RNAi strains displayed a significantly increased colonization of barley roots at 14 dpi compared with the WT and EV controls, calculated as the ratio of fungal DNA to plant DNA (elongation factor *PiTEF*/ubiquitin *HvUBI*) (Fig. 3C) or as the ratio of *PiTEF* to *HvUBI* transcripts (SI




Fig. 4. Gene expression profiles, VPE-like enzymatic activity, and amino acid levels in barley roots suggest extended biotrophy in *P. indica* AMT1-RNAi strains compared with WT. (A) Relative expression of the plant defense-related gene *PR10* and (B) of the *P. indica* gene *PIIN_05889* encoding a putative xylanase, during colonization of barley by the *P. indica* WT and RNAi strain Amt28 at different time points. Expression data are standardized relative to *HvUBI* or to *PiTEF*. SEs are calculated from three independent biological repetitions. (C) VPE-like enzymatic activities during biotrophic (≤ 5 dpi) and cell death-associated colonization (≥ 10 dpi) of barley roots colonized by *P. indica* WT and RNAi strain Amt28 or mock treated. For the assay, the fluorescent VPE-specific substrate Ac-ESEN-MCA was added to the root extracts for spectrophotometric determination of enzymatic activities (for details, see SI Appendix). (D) VPE-like enzymatic activities during early (3 and 5 dpi) and late (10 and 14 dpi) colonization of *Arabidopsis* roots. (E) VPE-like enzymatic activities at the onset of saprotrophy (10 dpi) in barley roots and (F) in *Arabidopsis* roots. (G) Concentrations of free amino acids in the roots of barley noncolonized or colonized by *P. indica* WT and RNAi strain Amt28 at 7 and 14 dpi. (H) Concentrations of free amino acids in the roots of *Arabidopsis* at 7 and 14 dpi. Error bars represent SEM from three to four independent biological repetitions. **ANOVA, $P < 0.01$. Columns not sharing a letter are significantly different (ANOVA, $P < 0.01$).

Appendix, Fig. S9D). The temporal expression pattern for transcripts encoding the barley pathogenesis-related gene PR10 was examined to determine the plant's response to colonization by the *P. indica* WT and the RNAi strain Amt28. A decreased transcript accumulation for the PR gene in roots colonized by the RNAi strain compared with the WT strain at the onset of saprotrophy was observed (Fig. 4A). Consistent with this, expression of the *P. indica* saprotrophic marker gene, PIIN_05889 (encoding a putative xylanase), was reduced in the RNAi strain compared with *P. indica* WT (Fig. 4B). This indicates an extended biotrophic phase of the RNAi strain Amt28, which likely contributes to the enhanced colonization of barley roots at 14 dpi. To verify this hypothesis, we measured the activity of the vacuolar processing enzyme (VPE). This is a cysteine protease with caspase-like activity responsible for the maturation of various vacuolar proteins in higher plants and reported to be induced in dying cells (9, 26). VPE-like enzymatic activity increased significantly in barley roots colonized by *P. indica* WT compared with barley mock treated and barley roots colonized by the RNAi strain Amt28, especially at the onset of saprotrophy (Fig. 4C). Analyses of the VPE-like enzymatic activity performed at 10 dpi with the other RNAi strains and EV controls confirmed this finding (Fig. 4E). Remarkably, in the RNAi strains, elicitation of growth promotion was not affected compared with the WT situation (SI Appendix, Fig. S12). This indicates that an extended biotrophic phase does not seem to influence *P. indica*'s beneficial effects on barley.

Consistent with low expression of the *PiAMT1* gene in *Arabidopsis* at 14 dpi, analyses of the *P. indica* RNAi strains impaired in ammonium uptake and possibly ammonium sensing displayed no difference in their colonization patterns compared with the WT and EV controls (Fig. 3D and SI Appendix, Fig. S9E). Similarly, no differences in the VPE-like enzymatic activity were observed in *Arabidopsis* roots colonized by *P. indica* WT or RNAi strain Amt28, in which both displayed moderate increased activity compared with mock control at later stages (Fig. 4D and F). We therefore concluded that *PiAMT1* is not required for biotrophic growth but is involved in the lifestyle switch of *P. indica* to saprotrophic growth as observed in barley.

Quantification of free amino acid levels in barley roots by ultra-pressure reversed-phase chromatography (see SI Appendix for details on this method) showed that amino acid concentrations decreased significantly in colonized roots by *P. indica* WT compared with roots colonized by the RNAi strain Amt28 and control roots at the onset of the saprotrophic stage/late biotrophic stage (Fig. 4G and SI Appendix, Fig. S13). At 14 dpi, the concentrations of free amino acids were remarkably lower than at 7 dpi in the older/basal root zone of barley, irrespective of colonization (Fig. 4G and SI Appendix, Fig. S13). In particular, asparagine and glutamine decreased by approximately five- and sevenfold, respectively, in control roots (SI Appendix, Fig. S13). This might be the result of reallocation of nitrogen to younger root zones in response to developmental RCD (16) and is in agreement with the microarray data, which indicate limited nitrogen availability to *P. indica* in the basal root zone of barley at 14 dpi. In *Arabidopsis*, colonization by *P. indica* WT decreased amino acid concentrations 7 dpi (Fig. 4H and SI Appendix, Fig. S13), whereas the level of free amino acids increased significantly 14 dpi (Fig. 4H and SI Appendix, Fig. S13). The altered organic nitrogen allocation upon *P. indica* colonization at 14 dpi was mainly the result of changes in asparagine and glutamine (SI Appendix, Fig. S13) and suggests that nitrogen supply to the fungus is not limited in this phase. Biotrophic and hemibiotrophic fungi have been shown to induce plant nitrogen mobilization and accumulation at the site of infection. In particular, nitrogen-rich amino acids such as glutamine and asparagine have been identified as the major forms of nitrogen reallocation during infection of different plant hosts (27–29). Changes in free amino acid pools during biotrophy have been speculated to reflect the demand for organic nitrogen by the fungus or to be required to launch defense responses by the host (27, 30).

Asparagine and glutamine are the preferred nitrogen source of *P. indica* in axenic culture, and when grown on these amino acids as the sole nitrogen source, *P. indica* produces enlarged hyphae that resemble the multilobed biotrophic hyphae *in planta* (SI Appendix, Fig. S14). On nitrate-containing medium, known to be a poor nitrogen source for *P. indica* (8), or on medium without nitrogen, *P. indica* hyphae are thin and less branched, similar to the secondary hyphae found in barley at later colonization stages (SI Appendix, Fig. S14). Asparagine and glutamine therefore may represent a ready source of organic nitrogen during biotrophy.

Taken together, these results strongly suggest that host-related nutritional cues affect *P. indica*'s lifestyle and that the switch from biotrophic to saprotrophic nutrition during colonization of barley is affected by *PiAMT1* and by nitrogen availability.

Discussion

Establishment of biotrophy during colonization of *Arabidopsis* and barley by *P. indica* is an important feature of the symbioses with both plant hosts, and it implies a strong interdependence between host metabolism and fungal nutrient uptake in this endophyte. Transcriptional profiling revealed that far fewer genes need to be induced to maintain biotrophy compared with those necessary for coordinating the switch to saprotrophy and the manipulation of RCD. Evolution of obligate biotrophy was shown to correlate, among others, with the loss of genes involved in nitrate metabolism, and it is speculated that host plants provide a ready source of organic nitrogen in the form of amino acids (15). In the draft genome of *P. indica*, no nitrate transporters or reductases were found (8), indicating that nitrate is not essential for this symbiont. *P. indica* can grow on ammonium and on glutamine and asparagine, which are its favorite nitrogen sources. The low expression of the *P. indica* high-affinity ammonium transporter in *Arabidopsis* at later colonization stages suggests that




Fig. 5. Schematic representation of *P. indica* colonization strategies at different symbiotic stages in barley and in *Arabidopsis*. (Upper) Invasive hyphae (IH) and secondary thin hyphae (SH) of *P. indica* in barley dead cells (10 dpi). (Lower) *P. indica* non-WGA-stainable biotrophic broad invasive hyphae in an *Arabidopsis* epidermal cell (10 dpi). Fungal structures were stained with WGA-AF488 (green); membranes were stained with FM4-64 (red). Staining of fungal structures by WGA-AF488 is enhanced in dead host cells. In *H. vulgare* roots, cell death is initiated a few days after germination by developmental RCD (16) and is increased by *P. indica* colonization.

an adequate source of nitrogen is provided by this host during colonization. Increased concentrations of free amino acids, principally of glutamine and asparagine upon *P. indica* colonization of this host at 14 dpi, strongly support this conclusion. The induction of *PiAMT1* and other fungal nitrogen transporters in barley leads to the assumption that this plant cannot provide *P. indica* with sufficient organic nitrogen during the onset of the RCD program and that nitrogen depletion therefore may function as a trigger for the *in planta* expression of fungal genes encoding hydrolytic enzymes and for the activation of the saprotrophic program. Silencing of the high-affinity ammonium transporter *PiAMT1* by RNAi resulted in reduced expression of fungal xylanase, barley defense response, and VPE activity in colonized roots, supporting the hypothesis that PiAmt1 is involved in sensing the N status and in downstream signaling upon nitrogen depletion in *P. indica*. Because biotrophic growth of *P. indica* in *Arabidopsis* was independent of PiAmt1, we conclude that expression and a signaling function of PiAmt1 are needed for the switch of *P. indica*'s lifestyle to saprotrophy. Loss or suppression of the expression of *PiAMT1* would represent a step toward a progressively more intimate biotrophic association with its hosts. The maintenance of this gene at the expense of biotrophy might benefit *P. indica* during prolonged saprotrophic growth on decaying plant material, making this fungus also able to survive in the absence of living hosts or on dying host cells.

Plant-associated fungi are either specialists, which are adapted to one or a few distinct hosts, or generalists that can thrive in highly variable host environments. Specialists and their hosts are in an evolutionary arms race that leads to the development of fungal tools and colonization strategies that are efficiently tailored to the respective host. Conversely, broad-host range species must evolve adaptations to cope with a plethora of different host-associated signals and host-specific defense mechanisms. The evolutionary force, in this case, possibly drives the expansion and diversification of the fungal toolkit and the host-adapted gene expression to better suit different plants. Recently it was shown that the obligate biotrophic ascomycete powdery mildew

pathogen, *Blumeria graminis* f. sp. *hordei*, which can grow and reproduce only on living cells of its natural host, barley, displays a conserved transcriptional program during early pathogenesis on barley and on immunocompromised *Arabidopsis* (31). Although we cannot exclude that at early time points (<2 dpi, prepenetration stage) a certain conservation of the transcriptional program may be present, our data show that the broad-host range fungal root symbiont *P. indica* responds differently to divergent hosts, especially at later time points during establishment and maintenance of the intracellular biotrophic interaction.

In conclusion, the transcriptional and phenotypic plasticity of *P. indica* during symbioses (summarized in Fig. 5) establishes a highly adaptive capacity in a root endophyte with broad compatibility that can reconfigure itself and its lifestyle in response to different environmental and host signals.

Material and Methods

Microarray Analyses. Microarray experiments were performed with total RNA extracted from *P. indica*-inoculated barley and *Arabidopsis* roots. As a control, total RNA from *P. indica* grown on 1/10 PNM-agar was used. Root samples from three independent biological replicates were labeled and hybridized according to Agilent's One-Color Microarray-Based Gene Expression Analysis Low Input Quick Amp Labeling protocol. For details, see *SI Appendix*.

RNAi Vector Construction and *P. indica* Transformation. A 570-bp fragment of the *PiAMT1* gene was amplified by PCR (*Dataset S5*) from cDNA and inserted in the EcoRV site of the convergent dual-promoter vector pPiRNAi (17). *P. indica* was transformed with vector pPiRNAi-AMT1 and the EV control as described in ref. 17. For details, see *SI Appendix*.

ACKNOWLEDGMENTS. We thank Elmar Meyer for technical support and Regine Kahmann and Gregor Langen are gratefully acknowledged for reading the manuscript prior to submission. A.Z. acknowledges support from Max-Planck-Gesellschaft and Deutsche Forschungsgemeinschaft Grant ZU263/2-1. U.L. acknowledges support from the International Max Planck Research School for Environmental, Cellular and Molecular Microbiology (IMPRS-MIC) Marburg.

- López-Berges MS, Rispaill N, Prados-Rosales RC, Di Pietro A (2010) A nitrogen response pathway regulates virulence functions in *Fusarium oxysporum* via the protein kinase TOR and the bZIP protein MeaB. *Plant Cell* 22(7):2459–2475.
- Stuttman J, et al. (2011) Perturbation of *Arabidopsis* amino acid metabolism causes incompatibility with the adapted biotrophic pathogen *Hyaloperonospora arabidopsidis*. *Plant Cell* 23(7):2788–2803.
- Kämper J, et al. (2006) Insights from the genome of the biotrophic fungal plant pathogen *Ustilago maydis*. *Nature* 444(7115):97–101.
- Tyler BM, et al. (2006) Phytophthora genome sequences uncover evolutionary origins and mechanisms of pathogenesis. *Science* 313(5791):1261–1266.
- Peskan-Berghofer T, et al. (2004) Association of *Piriformospora indica* with *Arabidopsis thaliana* roots represents a novel system to study beneficial plant-microbe interactions and involves early plant protein modifications in the endoplasmic reticulum and at the plasma membrane. *Physiol Plant* 122(4):465–477.
- Waller F, et al. (2005) The endophytic fungus *Piriformospora indica* reprograms barley to salt-stress tolerance, disease resistance, and higher yield. *Proc Natl Acad Sci USA* 102(38):13386–13391.
- Lahrman U, Zuccaro A (2012) Opprimo ergo sum—evolution and suppression in the root endophytic fungus *Piriformospora indica*. *Mol Plant Microbe Interact* 25(6):727–737.
- Zuccaro A, et al. (2011) Endophytic life strategies decoded by genome and transcriptome analyses of the mutualistic root symbiont *Piriformospora indica*. *PLoS Pathog* 7(10):e1002290.
- Qiang X, Zechmann B, Reitz MU, Kogel KH, Schäfer P (2012) The mutualistic fungus *Piriformospora indica* colonizes *Arabidopsis* roots by inducing an endoplasmic reticulum stress-triggered caspase-dependent cell death. *Plant Cell* 24(2):794–809.
- Deshmukh S, et al. (2006) The root endophytic fungus *Piriformospora indica* requires host cell death for proliferation during mutualistic symbiosis with barley. *Proc Natl Acad Sci USA* 103(49):18450–18457.
- Jacobs S, et al. (2011) Broad-spectrum suppression of innate immunity is required for colonization of *Arabidopsis* roots by the fungus *Piriformospora indica*. *Plant Physiol* 156(2):726–740.
- Stergiopoulos I, de Wit PJ (2009) Fungal effector proteins. *Annu Rev Phytopathol* 47: 233–263.
- Plett JM, et al. (2011) A secreted effector protein of *Laccaria bicolor* is required for symbiosis development. *Curr Biol* 21(14):1197–1203.
- Kloppholz S, Kuhn H, Requena N (2011) A secreted fungal effector of *Glomus intraradices* promotes symbiotic biotrophy. *Curr Biol* 21(14):1204–1209.
- Spanu PD, et al. (2010) Genome expansion and gene loss in powdery mildew fungi reveal tradeoffs in extreme parasitism. *Science* 330(6010):1543–1546.
- Liljeroth E, Bryngelsson T (2001) DNA fragmentation in cereal roots indicative of programmed root cortical cell death. *Physiol Plant* 111(3):365–372.
- Hilbert M, et al. (2012) Indole derivative production by the root endophyte *Piriformospora indica* is not required for growth promotion but for biotrophic colonization of barley roots. *New Phytol* 196(2):520–534.
- Schoonbeek H, Del Sorbo G, De Waard MA (2001) The ABC transporter BcatrB affects the sensitivity of *Botrytis cinerea* to the phytoalexin resveratrol and the fungicide fenpiclonil. *Mol Plant Microbe Interact* 14(4):562–571.
- Fleissner A, Sopalla C, Weltring KM (2002) An ATP-binding cassette multidrug-resistance transporter is necessary for tolerance of *Gibberella pulicaris* to phytoalexins and virulence on potato tubers. *Mol Plant Microbe Interact* 15(2):102–108.
- Urban M, Bhargava T, Hamer JE (1999) An ATP-driven efflux pump is a novel pathogenicity factor in rice blast disease. *EMBO J* 18(3):512–521.
- Ahuja I, Kissen R, Bones AM (2012) Phytoalexins in defense against pathogens. *Trends Plant Sci* 17(2):73–90.
- Lorenz MC, Heitman J (1998) The MEP2 ammonium permease regulates pseudohyphal differentiation in *Saccharomyces cerevisiae*. *EMBO J* 17(5):1236–1247.
- Javelle A, et al. (2003) Molecular characterization, function and regulation of ammonium transporters (Amt) and ammonium-metabolizing enzymes (GS, NADP-GDH) in the ectomycorrhizal fungus *Hebeloma cylindrosporum*. *Mol Microbiol* 47(2):411–430.
- Marini AM, André B (2000) In vivo N-glycosylation of the mep2 high-affinity ammonium transporter of *Saccharomyces cerevisiae* reveals an extracytosolic N-terminus. *Mol Microbiol* 38(3):552–564.
- Marini AM, Soussi-Boudekou S, Vissers S, Andre B (1997) A family of ammonium transporters in *Saccharomyces cerevisiae*. *Mol Cell Biol* 17(8):4282–4293.
- Hara-Nishimura I, Hatsugai N, Nakaune S, Kuroyanagi M, Nishimura M (2005) Vacuolar processing enzyme: An executor of plant cell death. *Curr Opin Plant Biol* 8(4):404–408.
- Horst RJ, et al. (2010) *Ustilago maydis* infection strongly alters organic nitrogen allocation in maize and stimulates productivity of systemic source leaves. *Plant Physiol* 152(1):293–308.
- Solomon PS, Oliver RP (2001) The nitrogen content of the tomato leaf apoplast increases during infection by *Cladosporium fulvum*. *Planta* 213(2):241–249.
- Tavernier V, et al. (2007) The plant nitrogen mobilization promoted by *Colletotrichum lindemuthianum* in *Phaseolus* leaves depends on fungus pathogenicity. *J Exp Bot* 58(12):3351–3360.
- Liu G, et al. (2010) Amino acid homeostasis modulates salicylic acid-associated redox status and defense responses in *Arabidopsis*. *Plant Cell* 22(11):3845–3863.
- Hacquard S, et al. (2013) Mosaic genome structure of the barley powdery mildew pathogen and conservation of transcriptional programs in divergent hosts. *Proc Natl Acad Sci USA* 110(24):E2219–E2228.

1 **Supplementary Materials:**

2 Material and Methods

3 Supplementary Figures 1 to 15

4 Supplementary Datasets 1 to 5

5 **Material and methods**

6 **1. Fungal and plant material**

7 *Piriformospora indica* Sav. Verma, Aj. Varma, Rexer, G. Kost & P. Franken (DSM11827, Deutsche Sammlung von
8 Mikroorganismen und Zellkulturen, DSMZ, Braunschweig, Germany) cultures were propagated at 28°C in liquid
9 complete medium (CM) (1) with 2% glucose and 130 rpm shaking or on solid medium with 1.5% agar. Hygromycin
10 B (80 µg/ml) was supplemented for growth of *P. indica* transformants. For microarray experiments *P. indica* was
11 cultivated on 1/10 PNM plates (2) for 3 days. 1/10 PNM final concentrations: 0.5 mM KNO₃, 0.367 mM KH₂PO₄,
12 0.144 mM K₂HPO₄, 2 mM MgSO₄ x H₂O, 0.2 mM Ca(NO₃)₂, 0.25% (v/v) Fe-EDTA (0.56% w/v FeSO₄ x 7H₂O and
13 0.8% w/v Na₂EDTA x 2H₂O), 0.428 mM NaCl; pH-adjusted to 6.0 and buffered with 10 mM MES. For solid media,
14 0.4% (w/v) GELRITE (Duchefa) was added.

15 Barley seeds (*Hordeum vulgare* cv. Golden Promise, Ingrid or Ingrid *mlo-5* mutants I22, kindly provided by Gregor
16 Langen) were surface sterilized with 70% ethanol for 1 minute, 12% sodium hypochlorite for 1.5 hours and washed
17 with sterile distilled water for 3 hours. Sterilized seeds were kept in the dark for 3 days on sterile wet filter paper at
18 room temperature. For colonization studies, three-day-old barley seedlings were transferred into sterile jars
19 containing 1/10 PNM medium and inoculated with 3 ml chlamyospore suspension (5×10⁵ ml⁻¹ in 0.002%
20 TWEEN20). Incubation was performed in a Conviron phytochamber with a day/night cycle of 16/8 hours (light
21 intensity: 110 µmol m⁻² s⁻¹) at a temperature of 22/18°C. Tween water (0.002% TWEEN20) treated seedlings were
22 used as control if required. Root samples were collected at different time points as described in the figure legends
23 and carefully washed in distilled water. The first 4 cm of the roots, starting from the seed, were excised and
24 immediately frozen in liquid nitrogen or used for microscopy. All experiments were prepared in 3 to 4 independent
25 biological repetitions of four plants per jar.

26 *Arabidopsis thaliana* seeds (ecotype Columbia-0) were incubated for 5 min in 70% ethanol, surface sterilized for 5
27 min with 6% sodium hypochlorite and washed 6 times for 5 min in sterile water. After stratification for 3 days at
28 4°C in the dark on 1/10 PNM medium, Arabidopsis seedlings were grown for 14 days under sterile conditions in a
29 phytochamber (Vötsch, Balingen-Frommern, Germany) at long day conditions (day: 16 h, 23°C, 350 µmol m⁻² s⁻¹;
30 night: 8 h, 18°C; 60% humidity). For inoculation of Arabidopsis roots with *P. indica* chlamyospores, plants of
31 roughly the same size were first transferred to square petri dishes containing 1/10 PNM and then inoculated directly
32 with either 5 x 10⁵ *P. indica* spores per 20 seedlings or mock treated as described for barley. Root material was
33 harvested, washed carefully with distilled water and frozen liquid nitrogen after 3, 7 and 14 days. For each time
34 point roots from 80 to 100 plants were pooled and the experiments were performed in 3 to 4 independent biological
35 repetitions.

36 Chlamyospores were collected from 3 to 4 week-old CM plates using 0.002% TWEEN20 for all described
37 experiments.

38 **2. Growth promotion assays**

39 For co-culture of *A. thaliana* with *P. indica* or *P. williamsii* and tests for plant growth promotion, seeds were
40 sterilized by incubation for 5 min in 70% ethanol followed by 2 min in 100% ethanol, left to dry and stratified as

1 described before either on a modified Hoagland's medium with four times more phosphate: 5mM KNO₃; 5mM
2 Ca(NO₃)₂; 2mM MgSO₄; 4mM KH₂PO₄; 0.03g/L Sprint 138 iron chelate; 0,1% micronutrients solution containing
3 2.86g/L H₃BO₃; 1.81 g/L MnCl₂·4H₂O; 0.08g/L CuSO₄·5H₂O; 0.02 g/L 85% MoO₃·H₂O, based on (3) or on 1/10
4 PNM as described above. Media were solidified with 4g/L GELRITE (Carl Roth, Karlsruhe, Germany). For assays
5 on the modified Hoagland's medium, plants were grown for 7 to 10 days under long day conditions (16h light, at
6 23°C, 85 μmol m⁻² s⁻¹; 60% humidity), and on 1/10 PNM medium plants grew for 10 days (16h light, at 23°C, 350
7 μmol m⁻² s⁻¹; 60% humidity; night: 8h, 18°C) before mock-inoculation with tween water or inoculation with either
8 *P. indica* or *P. williamsii* (when indicated) chlamydospores (5 x 10⁵ ml⁻¹). The seeds that failed to germinate and the
9 seedlings that did not look healthy were removed from the plates preceding tween water or spore application. Plants
10 grown on modified Hoagland's were inspected for biomass 7 days post inoculation (dpi), and for seedlings grown on
11 1/10 PNM biomass was inspected 14 dpi. Statistical analyses of the results were performed in R 2.15.1 (4)
12 (<http://www.R-project.org/>). Pairwise comparisons of the different subsets of data were performed using ANOVA
13 followed by Tukey's HSD (honestly significant difference) on 95% family-wise confidence level. Homogeneous
14 subsets were identified using the package multcompView (5) (<http://CRAN.R-project.org/package=multcompView>).

15 Barley growth promotion experiments with *P. indica* WT, empty vector control and RNAi strains were carried out
16 as described in (6).

17 **3. DNA, RNA extraction and real time qPCR analyses**

18 DNA from 200 mg of ground fungal or plant material was isolated using the protocol of Doyle and Doyle (7). Total
19 RNA from 200 mg of ground material was extracted using TRIzol (Invitrogen, Karlsruhe, Germany) following the
20 manufacturer's instructions. The amount and quality of the extracted RNA were estimated by using an agarose gel
21 and the NanoDrop-1000 Spectrophotometer (Agilent) or a 2100 Bioanalyzer (Agilent, Santa Clara, USA) for the
22 microarray experiments. For real-time qPCR analyses, two μg of RNA were used for DNase I digestion (Fermentas)
23 followed by cDNA synthesis (First Strand cDNA Synthesis Kit, Fermentas). Real-time qPCR analyses were
24 performed from 10 ng DNA or cDNA mixed with the appropriate primers (Dataset S5) in 10 μl SYBRgreen
25 Supermix (BIORAD) using the following amplification protocol: initial denaturation for 5 min at 95°C, followed by
26 40 cycles with 30 s at 95°C, 30 s at 59°C, 30 s at 72°C and a melt curve analysis. Relative DNA amount or relative
27 expression and its fold change values were calculated using the 2^{-ΔΔC_i} method (8).

28 **4. Microarray experiments**

29 Microarray analyses were performed with 200 ng total RNA extracted from *P. indica* inoculated barley and
30 Arabidopsis roots at 14 dpi or from *P. indica* inoculated Arabidopsis roots at 3 dpi. As control total RNA from *P.*
31 *indica* grown on 1/10 PNM-agar was used. Here, 3 days old hyphae were thoroughly scratched from the agar surface
32 with a sterile scalpel and immediately frozen in liquid nitrogen. For the 3 dpi barley time point, data from Gene
33 Expression Omnibus (<http://www.ncbi.nlm.nih.gov/geo/>), accession number GSE31266 was used (9). For each
34 treatment, samples from three independent biological replicates were labeled and hybridized according to Agilent's
35 One-Color Microarray-Based Gene Expression Analysis Low Input Quick Amp Labeling protocol (version 6.5).
36 Cy5-3-labeled probes were hybridized to 8x60k custom-designed Agilent microarray chips. The microarray design
37 was performed based on a previous 2x105k custom-designed Agilent microarray described in (9). For this, the best
38 performing probe (BP) per *P. indica* gene was selected from the existing probes based on the following criteria: 1)
39 Probes with a cross hybridization potential were discarded if an alternative probe for the gene could be used. If this
40 was not possible, three probes with different cross hybridization targets were chosen and used (XP1-XP3). 2) The
41 signal intensity of each probe was compared to its position in the gene. The probe was defined to be a BP if its
42 signal intensity was high and its position was close to the 3' prime end of the respective gene. Thus, the BP probe for
43 each *P. indica* gene was selected as such if it had either the strongest signal of all probes of the respective gene and
44 was located at most 300 bp apart from the 3' prime end or if it was the closest to the 3' prime end and had at least
45 90% signal intensity of the strongest probe. 3) Non-uniformity outliers were detected by comparing the signal

1 intensity between biological replicates. The signal intensity of the best probe defined in step 2 was compared against
2 those of its intra- and inter-array replicates. If the signal intensity ranged between +/- 1.42 of the interquartile range,
3 it was accepted as the best probe. Alternatively, step 3 was repeated with the second (third, etc.) best probe. 4) New
4 probes (NP) were calculated via the eArray program (earray.chem.agilent.com/earray/) for those genes that had no
5 suitable BP probe. Control probes were additionally loaded on the chip as described before (9). Microarray image
6 files were analyzed using Agilent's Feature Extraction software v. 10.5 which calculates for each spot a background
7 corrected signal intensity value (gProcessedSignal) that was used for further analysis. For statistical analysis of the
8 raw data, the R environment (www.r-project.org; version 2.15.1) including the Bioconductor package 'Limma' was
9 used. In summary, raw data were standardized by quantile normalization, intensity values from replicate probes
10 were averaged, log₂-ratios between experiments were calculated and Student's t-statistic applied to test for
11 significance. Quality and suitability of the applied statistics was estimated by generating density and MA plots. The
12 degree of variability between the experiments was shown by principle component analysis (Figure S15). Selection
13 of differentially expressed genes was based on a fold change of 2 and a false discovery rate-adjusted significance
14 level (adj. p-value) of less than 0.05. Expression data from all experiments are stored in the GEO database
15 (accession number GSE47775). Microarray data were verified by quantitative real-time PCR (qRT-PCR) as
16 described previously (9) from three biologically independent kinetics for each host (time points: 3, 7 and 14 dpi for
17 Arabidopsis; 3, 5, 10 and 14 dpi for barley).

18 **5. Enrichment analysis**

19 To identify significantly enriched gene ontology (GO) terms from the performed microarray experiments the Gene
20 Ontology Enrichment Analysis Software Toolkit (GOEAST) was used
21 (omicslab.genetics.ac.cn/GOEAST/index.php) with settings for customized microarray platform. GO's were
22 assigned to *P. indica* genes by using the Blast2GO suite (version 2.6.0) (10). Significantly higher expressed genes in
23 Arabidopsis roots colonized by *P. indica* at 14 dpi compared to barley roots colonized by *P. indica* at 14 dpi (and
24 *vice versa*) were analyzed using the recommended parameter settings. Datasets S4a and S4b, summarizing all
25 enriched GO terms (Arabidopsis: GO_AT14up, Barley: GO_HV14up), were prepared from the GOEAST outputs.

26 **6. Confocal laser microscopy**

27 *Cell viability*

28 Root colonization and Arabidopsis/barley epidermal and cortex cells viability were analyzed by confocal
29 microscopy. Colonized roots were stained for 10 min with 10 µg/ml WGA-AF488 (Molecular Probes, Karlsruhe,
30 Germany) to visualize fungal structures. Membranes were stained with 3 µM FM4-64 (Probes, Karlsruhe, Germany)
31 for 5 min. Root samples were imaged with a TCS-SP5 confocal microscope (Leica, Bensheim, Germany) using an
32 excitation at 488 nm for WGA-AF488 and detection at 500–540 nm. Excitation of FM4-64 was performed at 633
33 nm and detection at 650–690 nm.

34 *Maximum projection and serial slices of barley root material*

35 Root segments of *Hordeum vulgare* were fixed in FPA (formalin (37%): propionic acid (> 99%): ethanol (50%),
36 0.5:0.5:9) for three days and subsequently stored in 70% ethanol for further use. For microtomy the root segments
37 were dehydrated in an ascending ethanol series and embedded in Unicryl™ (British Biocell Int.). For serial slices,
38 sequential series of 4 µm longitudinal and transversal sections (Leica Supercut 2065) were stained with toluidine
39 blue O (1 g toluidine blue O + 1 g sodium tetraborate in 100 ml distilled H₂O) and mounted in Corbit balsam.
40 Pictures were taken with a Leica DMRB or TCS-SP5 microscope, equipped with a digital camera (Moticam 2300).
41 For maximum projections the root of barley was stained with acid fuchsin and depicted by confocal laser scanning
42 microscopy using an Argon blue light laser (488 nm) as described in (11). The z-stack was used to compute a
43 Maximum Projection (LCS Version 2).

1 7. Yeast complementation assay

2 The ammonium import function of PiAmt1 (PIIN_02036) was verified by yeast complementation analysis. For this,
3 full length cDNA of *PiAMT1* was cloned into the *NotI* site of the pDR195 vector (kindly provided by Dr. Mike
4 Guether, Karlsruhe Institute of Technology, KIT) with primers PiAMT1_SC_notI_F and PiAMT1_SC_notI_R
5 (Dataset S5) and the resulting plasmid was then transformed by heat shock into the ura-AMT-defective yeast
6 (*Saccharomyces cerevisiae*) strain 31019b; $\Delta\Delta\Delta\text{mep1;2;3}$. Growth complementation assays were performed on solid
7 YNB medium (Yeast Nitrogen Base w/o N, Bacto) supplemented with 2% glucose and 1 mM or 5 mM $(\text{NH}_4)_2\text{SO}_4$
8 as sole nitrogen source.

9 8. RNAi vector construction and *P. indica* transformation

10 To perform RNAi silencing experiments, a 570 bp fragment of the *PiAMT1* (PIIN_02036) cDNA was amplified by
11 PCR with primers PiAMT1_RNAi_ecoRV_F and PiAMT1_RNAi_ecoRV_R (Dataset S5) and then inserted in the
12 *EcoRV* site of the convergent dual promoter vector pPiRNAi (6). PCR reactions were performed using a proof
13 reading Pfu Polymerase (Promega). The vector pPiRNAi-AMT1 and empty vector control were sequenced and
14 subsequently transformed into *P. indica* by PEG-mediated transformation as described in (6). Putative transformants
15 generated from three independent transformations were transferred onto new CM plates with 80 $\mu\text{g/ml}$ hygromycin
16 B. Cloning was performed using T4 ligase (NEB), Antarctic phosphatase (NEB) and *E. coli* Top10 cells. Vector
17 sequencing was done by Eurofins MWG Operon. The success of transformation was confirmed by southern blot and
18 the efficiency of silencing was verified by qPCR experiments. All RNAi strains were analyzed for reduced growth
19 on minimal medium with ammonium as sole nitrogen source as described in the next chapter. Selected silenced
20 strains with a reduced growth phenotype on minimal medium supplemented with ammonium as sole nitrogen source
21 and with a normal growth phenotype on CM were chosen for *in planta* assays.

22 9. Ammonium feeding test of *P. indica* transformants

23 *P. indica* transformants harboring the RNAi construct were analyzed for ammonium uptake by using wild-type and
24 empty vector transformed strains as controls. Ten μl of chlamydo-spores suspension with 10-fold serial dilutions
25 starting from $5 \times 10^5 \text{ ml}^{-1}$ were inoculated onto YNB medium (Yeast Nitrogen Base w/o N, Bacto) supplemented with
26 2 mM NH_4Cl (Roth) for 14 days. The experiments were carried out in three replicates.

27 10. Measurement of caspase activity in roots

28 Barley and Arabidopsis roots were inoculated with *P. indica* WT, RNAi strains, empty vector controls or mock
29 treated. The first 4 cm of the roots, starting from the seed, were excised and immediately frozen in liquid nitrogen
30 from plant samples harvested at 3, 5, 10 and 14 dpi. One hundred mg ground roots were extracted with a buffer
31 containing 100 mM sodium acetate (pH 5.5), 100 mM NaCl, 1 mM EDTA, 2mM DTT and 1mM
32 phenylmethylsulfonyl fluoride. To measure caspase activities, 100 μM fluorogenic VPE substrate (Ac-ESEN-MCA,
33 Peptide Institute) was added to the root extracts. Fluorescence intensities were measured at 465 nm after excitation
34 at 360 nm using a fluorescence microplate reader (TECAN Safire). The enzyme activity was calculated as the
35 increasing unit of fluorescent intensity compared to the buffer+substrate control after 60 minutes incubation.

36 11. Preparation and assay of samples for amino acid concentration

37 Plant material was incubated for 60 minutes at 80°C in 80% ethanol and thereafter centrifuged for 10 minutes at
38 14000 rpm and 4°C. Supernatant were evaporated to dryness, re-suspended in purest water and used for HPLC
39 analysis. Prior to HPLC analysis samples were derivatized using a fluorescing reagent AQC (6-aminoquinolyl-N-
40 hydroxysuccinimidylcarbamate). 3 mg of self-made AQC (IPK, Germany) was dissolved in 1 ml acetonitrile and
41 incubated exactly for 10 minutes at 55°C. The prepared reagent was stored at 4°C and used up to four weeks. For
42 derivatization of the sample 0.01 ml of the prepared reagent solution was used for each sample which contained 0.8

1 ml of a buffer (0.2 M boric acid, pH 8.8) and 0.01 ml of the supernatant. Separation of soluble amino acids was performed on a newly developed UPLC-based method using Ultra pressure reversed phase chromatography (AcQuity H-Class, Waters GmbH, Germany). UPLC system consisted of a quaternary solvent manager, a sample manager-FTN, a column manager and a fluorescent detector (PDA eλ Detector). The separation was carried out on a C18 reversed phase column (ACCQ Tag Ultra C18, 1.7 μm, 2.1x100 mm) with a flow rate of 0.7 ml per min and duration of 10.2 min. The column was heated at 50°C during the whole run. The detection wavelengths were 266 nm for excitation and 473 nm as emission. The gradient was accomplished with four solutions prepared from two different buffers purchased from Waters GmbH (eluent A concentrate and eluent B for amino acid analysis, Germany). Eluent A was pure concentrate, eluent B was a mixture of 90% LCMS water (The Geyer GmbH, Germany) and 10 % eluent B concentrate, eluent C was pure concentrate (eluent B for amino acid analysis) and eluent D was LCMS water (The Geyer GmbH, Germany). The column was equilibrated with eluent A (10%) and eluent C (90%). The gradient was produced as indicated in the table below.

Retention time in min	Eluent A (%)	Eluent B (%)	Eluent C (%)	Eluent D (%)
0	10	0	90	0
0.29	9.9	0	90.1	0
5.49	9	80	11	0
7.10	8	15.6	57.9	18.5
7.30	8	15.6	57.9	18.5
7.69	7.8	0	70.9	21.3
7.99	4	0	36.3	59.7
8.59	4	0	36.3	59.7
8.68	10	0	90	0
10.2	10	0	90	0

13

14 12. Phylogeny

15 Phylogenetic relations of ammonium transporters from *Piriformospora indica*, *Saccharomyces cerevisiae* and
 16 *Hebeloma cylindrosporum* were calculated using seaview (version 4) (12). Protein alignment was generated by
 17 MUSCLE (v3.8.31) (13) and phylogenetic tree by PHYML (v3.0) (14) using standard settings. An approximate
 18 *likelihood-ratio* test (aLRT) was performed to calculate branch support.

19 Legends for the supplementary figures

20 **Figure S1.** Host-dependent expression profiles of *in planta*-induced *P. indica* genes encoding intracellular (A) and
 21 secreted (B) proteins at 3 and 14 days post inoculation (dpi) during Arabidopsis (PI_AT) and barley (PI_HV)
 22 colonization, calculated versus PNM-control.

23 **Figure S2.** Heatmaps showing normalized, background-corrected log₂ signal intensity values of 305 hydrolytic
 24 enzymes (left) and 252 transporters (right) as previously classified (9). Columns represent biological independent
 25 replicates of *P. indica* colonized barley roots 3 (PI_HV_3) and 14 dpi (PI_HV_14), *P. indica* colonized Arabidopsis
 26 roots 3 (PI_AT_3) and 14 dpi (PI_AT_14) and *P. indica* on 1/10 PNM-agar alone 3dpi (PI_PNM_3). Signal
 27 intensities (gProcessedSignal) from Agilent's feature extraction software (v10.5) were quantile normalized and intra-
 28 array replicates were averaged using R (www.r-project.org; version 2.15.1). Only those genes are shown for which a
 29 best performing probe could be designed (see microarrays section in material and methods for details). Data from
 30 the time point PI_HV_3 is from Gene Expression Omnibus (<http://www.ncbi.nlm.nih.gov/geo/>), accession number
 31 GSE31266.

1 **Figure S3.** Barley root colonized by *P. indica* 30 dpi. A colonized epidermal cell with larger hyphae and thinner
2 hyphae in the cortex cells are visible. Hyphal contact with the endodermis layer results in autofluorescence which is
3 visible as a diffuse white signal on the host cell wall. For maximum projections the root of barley was stained with
4 acid fuchsin and depicted by confocal laser scanning microscopy using an Argon blue light laser (488 nm) as
5 described in (11).

6 **Figure S4.** Growth promotion effect on Arabidopsis plants ecotype Col-0 colonized by *P. indica* or by the closely
7 related sebacinoid fungus *P. williamsii* (2). (A-C) Box-plots of the fresh weight of *A. thaliana* plants grown in the
8 presence or absence of *P. indica* or *P. williamsii*. In all 3 experiments, treatment had a significant effect on the mean
9 fresh weight of the plants (plot A: ANOVA, $F_{2,74}=64.45$, $p<0.001$; plot B: ANOVA, $F_{2,28}=44.63$, $p<0.001$; plot C:
10 ANOVA, $F_{2,48}=14.07$, $p<0.001$). Treatment with *P. indica* chlamydo-spores resulted in a significant increase on the
11 mean fresh weight of the plants in all treatments (Tukey's HSD, $p<0.001$), but treatment with *P. williamsii*
12 chlamydo-spores had no significant effect (A, C) or a small but significant effect on the mean fresh weight in
13 comparison with the mock-treatment (B) (Tukey's HSD for plot A: $p=0.51$; for plot B: $p=0.012$; for plot C:
14 $p=0.985$). (D) Box-plot of the fresh weight of *A. thaliana* plants grown in the presence or absence of *P. indica* on
15 1/10 PNM. The mean fresh weight of *P. indica* treated plants was significantly increased compared to that of mock-
16 treated plants (two-sided t-test, unpaired, equal variance: $t = -3.04$, $df = 48$, $p\text{-value} < 0.01$). In all plots, boxes not
17 sharing a letter are significantly different (based on Tukey's HSD). Culture media (HO: modified Hoagland's; PNM:
18 1/10 PNM), plant age (dps = days post-sowing) and co-culture times on the respective plant media (dpi = days post
19 inoculation) are indicated in the upper left corner of each plot. (E) Representative plates containing sets of 14-day
20 old plants 7 days after mock-treatment with tween water (left) or inoculation with either *P. indica* (center) or *P.*
21 *williamsii* (right) chlamydo-spores.

22 **Figure S5.** *Piriformospora indica* biotrophic broad invasive hyphae (white asterisk) in Arabidopsis epidermal cell
23 (A) 10 and (B) 21 dpi. In contrast to extracellular hyphae, IH are not stainable with WGA-AF488 (green) due to the
24 presence of a plant-derived membrane. (C) After cooking with KOH for 5 minutes and infiltrating WGA for 10
25 minutes the hyphae of *P. indica* can be stained, indicating that chitin is still present in the cell wall of the fungus.

26 **Figure S6.** Gene Ontology Enrichment Analyses for the GO category biological process performed using GOEAST
27 (<http://omicslab.genetics.ac.cn/GOEAST/index.php>). Shown is the graphical output for the significantly enriched
28 GO terms, as well as their relationships in the whole GO hierarchy, among the *P. indica* genes induced during
29 colonization of Arabidopsis compared to barley at 14 dpi. Data displayed in this graphic show enrichment for genes
30 involved in tRNA aminoacylation for protein translation, amino acid biosynthesis, glycolysis, tetracycline transport
31 and mitochondrion organization.

32 **Figure S7.** Gene Ontology Enrichment Analyses for the GO category biological process performed using GOEAST
33 (<http://omicslab.genetics.ac.cn/GOEAST/index.php>). Shown is the graphical output for the significantly enriched
34 GO terms among the *P. indica* genes induced during colonization of barley compared to Arabidopsis at 14 dpi. Data
35 displayed in this graphic show enrichment for genes involved in chitin catabolic process, L-arabinose metabolic
36 process, xylan and cellulose catabolic processes and carbohydrate transport.

37 **Figure S8.** Analyses of *P. indica* ammonium transporter Amt1. (A) Expression of *PiAMT1* is ammonium depletion
38 responsive. *P. indica* was pre-grown on CM for one week (T_0) and then transferred to YNB medium supplemented
39 with different concentrations of ammonium: without nitrogen (w/o N); with 0.5 mM $(\text{NH}_4)_2\text{SO}_4$; with 2 mM
40 $(\text{NH}_4)_2\text{SO}_4$; with 10 mM $(\text{NH}_4)_2\text{SO}_4$. Mycelium was collected after 12 and 24 hours and immediately frozen in
41 liquid nitrogen prior to RNA extraction and cDNA synthesis for real time qPCR analyses. (B) *PiAmt1* is related to
42 high affinity ammonium transporters. Phylogenetic relationships between the deduced amino acid sequence of *P.*
43 *indica* Amt1 and Amt2 and *Hebeloma cylindrosporium* and *Saccharomyces cerevisiae* ammonium
44 transporters/permeases. Sequences were obtained from the GenBank database. ScMEP2, HcAmt1, HcAmt2 are

1 functionally characterized as high affinity ammonium transporters, whereas ScMEP1, ScMEP3 and HcAmt3 are low
2 affinity ammonium transporters (15-16). (C) Yeast complementation assay for the high affinity ammonium
3 transporter PiAmt1. Growth of ammonium uptake-deficient yeast strain $\Delta\Delta\Delta\text{mep1;2;3}$ transformed with the empty
4 control plasmid pDR195 or with pDR195::PiAmt1 on YNB medium supplemented with 5 mM or 1 mM $(\text{NH}_4)_2\text{SO}_4$
5 as the sole nitrogen source. Shown are serial dilutions of yeast cell suspensions ranging from 1 to 1×10^{-3} .
6 Experiments were performed in three independent biological repetitions. (D) Predicted topological structure of the
7 *P. indica* Amt1.

8 **Figure S9.** Analyses of selected *PiAMT1* RNAi strains and controls *in planta*. (A-B) Relative expression of the *P.*
9 *indica* high affinity ammonium transporter *AMT1* compared to *PiTEF* during colonization of barley and Arabidopsis
10 at 14 days post inoculation (dpi) in the wild type (WT), empty vector controls (EV) and RNAi strains (Amt9 and
11 Amt28). (C) Relative expression of the *P. indica* ammonium transporter *PiAMT2* compared to *PiTEF* during
12 colonization of barley at 14 days post inoculation (dpi) in the wild type (WT), empty vector controls (EV) and RNAi
13 strains (Amt9 and Amt28). Error bars represent SE of the mean from three independent biological repetitions. No
14 compensatory upregulation of *PiAMT2* was observed in the *PiAMT1* RNAi strains. (D-E) Relative abundance of
15 fungal *PiTEF* transcripts compared to plant *UBI* transcripts in barley and Arabidopsis roots colonized by *P. indica*
16 wild type (WT), empty vector controls (EV) and RNAi strains (Amt9 and Amt28) at 14 dpi. Silencing of *PiAMT1*
17 results in a significant increased colonization of barley roots compared to WT and EV controls (t-test $P < 0.05$). Error
18 bars represent SE of the mean from three independent biological repetitions. *P. indica* colonized Arabidopsis and
19 barley plants were grown on 1/10 PNM medium.

20 **Figure S10.** Host-dependent expression profiles of *in planta*-induced *P. indica* genes encoding ABC transporters
21 identified using the TCDB (<http://www.tcdb.org/>) and Blast2Go as described in (9).

22 **Figure S11.** Southern blot and colonies phenotype of the *P. indica* ammonium-uptake deficient RNAi strains. (A)
23 Southern blot analysis of *P. indica* WT and EV controls and RNAi strains. Genomic DNA from seven days old *P.*
24 *indica* cultures grown on CM was extracted and digested overnight with *SacI* (NEB). The digested DNA was
25 separated on 0.8% TAE agarose gel for 3 h at 80V. Blotting was performed using the DIG-labeling and detection kit
26 following the manufacturer's instructions (Roche Biochemicals). Hybridization was performed with a HYG probe.
27 (B) Colonies phenotype of *P. indica* WT and EV controls and RNAi strains on YNB medium supplemented with 10
28 mM NH_4Cl as sole nitrogen source. Plates were supplemented with 1% glucose and buffered with MES pH 5.6. Ten
29 microliters of chlamydo spores suspension with 10-fold serial dilutions starting from $5 \times 10^5 \text{ ml}^{-1}$ were used for the
30 growth assay. The experiments were carried out in three replicates. Representative photographs were taken at 7 days
31 after inoculation.

32 **Figure S12.** Barley growth promotion experiment with *P. indica* WT, empty vector control and RNAi strains. Plants
33 were grown as described in (6) and harvested at 30 days post inoculation. (A) Length of leaves in cm. (B) Leaves
34 dry weight in grams. (C) Root dry weight (in grams) in colonized or non-colonized barley plants. Error bars
35 represent standard errors of the mean ($n=20$). Asterisks represent significant differences (ANOVA, $P < 0.01$). (D)
36 Growth promotion in representative non-colonized (left) and colonized barley plants by *P. indica* WT (middle) and
37 RNAi strain Amt28 (right).

38 **Figure S13.** Concentrations of free amino acids in barley and Arabidopsis roots (4 cm below the seed). (A)
39 Concentrations of the most abundant free amino acids in the roots of barley non-colonized or colonized by *P. indica*
40 WT and RNAi strain Amt28 at 7 and (B) 14 dpi. Error bars represent SEM from three independent biological
41 repetitions with two to three technical repetitions. (C) Concentrations of the most abundant free amino acids in roots
42 of Arabidopsis non-colonized and colonized by *P. indica* WT and RNAi strain Amt28 at 7 and (D) 14 dpi. Error
43 bars represent SEM from four independent biological repetitions. *T-test $P < 0.05$, **T-test $P < 0.01$.

1 **Figure S14.** Morphology of *P. indica* hyphae stained with WGA-AF488 grown on minimal medium containing (A)
2 nitrate NaNO₃ 2mM, (B) asparagine Asn 1mM, or (C) glutamine Gln 1mM as sole nitrogen source. (D) Average of
3 hyphal size on different nitrogen sources. (E-F) Enlarged, multilobed hyphae from *P. indica* grown on asparagine.

4 **Figure S15.** Principle component analysis (PCA) performed using R (www.r-project.org; version 2.15.1) with
5 normalized, background-corrected log₂ signal intensity values. Shown are three main principle components (90.06%
6 of the total variability) in a x-y- (PC1 vs PC2) and z-y- (PC3 vs PC2) projection. The PCA shows that the whole
7 transcriptome of *P. indica* colonizing barley at 14 dpi (PI_HV_14) is most different from the other conditions
8 supporting the overall switch of the fungal transcriptome to a different feeding state. The second highest variation is
9 shown for *P. indica* colonizing Arabidopsis at 3 dpi (PI_AT_3).

10 **Dataset S1.** Significantly differentially regulated genes from microarray data. AT, *Arabidopsis thaliana*; PNM,
11 plant minimal medium; HV, *Hordeum vulgare*; dpi, days post inoculation.

12 **Dataset S2.** Expression data for the DELD protein family.

13 **Dataset S3.** Top20 induced *P. indica* SSPs during colonization of Arabidopsis and barley roots.

14 **Dataset S4.** Enrichment analysis performed using GOEAST for upregulated genes.


15 **Dataset S5.** List of primers used in this study.

16 **References**


- 17 1. Pham G. H. KR, Singh A., Malla R., Prasad R., Sachdev M., Kaldorf M., Buscot F., Oelmuller R.,
18 Hampp R., Saxena A. K., Rexer K.-H., Kost G., Varma A. (2008) Axenic Culture of Symbiotic
19 Fungus *Piriformospora indica*. *Plant Surface Microbiology* 593–613
- 20 2. Basiewicz M, *et al.* (2012) Molecular and phenotypic characterization of *Sebacina vermifera*
21 strains associated with orchids, and the description of *Piriformospora williamsii* sp. nov. *Fungal*
22 *Biol* 116(2):204-213.
- 23 3. Hoagland DR & Arnon DI (1950) The Water-Culture Method for Growing Plants without Soil.
24 *Calif. Agr. Expt. Sta. Circ.* 347
- 25 4. R-Core-Team (2012) R: A language and environment for statistical computing. R Foundation for
26 Statistical Computing, Vienna, Austria. ISBN 3-900051-07-0, .
- 27 5. Graves S, Piepho HP, & Selzer L (2012) multcompView: Visualizations of Paired Comparisons. R
28 package version 0.1-5. <http://CRAN.R-project.org/package=multcompView>.
- 29 6. Hilbert M, *et al.* (2012) Indole derivative production by the root endophyte *Piriformospora*
30 *indica* is not required for growth promotion but for biotrophic colonization of barley roots. *New*
31 *Phytol* 196(2):520-534.
- 32 7. Doyle JJ & Doyle JL (1987) A rapid DNA isolation procedure for small quantities of fresh leaf
33 tissue. *Phytochemical bulletin* 19(1):11-15.
- 34 8. Livak KJ & Schmittgen TD (2001) Analysis of relative gene expression data using real-time
35 quantitative PCR and the 2(T)^{-Delta Delta C} method. *Methods* 25(4):402-408.
- 36 9. Zuccaro A, *et al.* (2011) Endophytic Life Strategies Decoded by Genome and Transcriptome
37 Analyses of the Mutualistic Root Symbiont *Piriformospora indica*. *PLoS Pathog* 7(10):e1002290.
- 38 10. Conesa A, *et al.* (2005) Blast2GO: a universal tool for annotation, visualization and analysis in
39 functional genomics research. *Bioinformatics* 21(18):3674-3676.
- 40 11. Rath M (2013) Combining Microtomy and Confocal Laser Scanning Microscopy for structural
41 analyses of plant-fungus associations. *PhD thesis*.

- 1 12. Gouy M, Guindon S, & Gascuel O (2010) SeaView version 4: A multiplatform graphical user
2 interface for sequence alignment and phylogenetic tree building. *Mol Biol Evol* 27(2):221-224.
- 3 13. Edgar RC (2004) MUSCLE: a multiple sequence alignment method with reduced time and space
4 complexity. *Bmc Bioinformatics* 5:113.
- 5 14. Guindon S & Gascuel O (2003) A simple, fast, and accurate algorithm to estimate large
6 phylogenies by maximum likelihood. *Syst Biol* 52(5):696-704.
- 7 15. Lorenz MC & Heitman J (1998) The MEP2 ammonium permease regulates pseudohyphal
8 differentiation in *Saccharomyces cerevisiae*. *EMBO J* 17(5):1236-1247.
- 9 16. Javelle A, *et al.* (2003) Molecular characterization, function and regulation of ammonium
10 transporters (Amt) and ammonium-metabolizing enzymes (GS, NADP-GDH) in the
11 ectomycorrhizal fungus *Hebeloma cylindrosporum*. *Mol Microbiol* 47(2):411-430.
- 12
13

A








B




0 μm 25










(A) 10 dpi








(B) 21 dpi




(C) 10 dpi






A**B**PhyML $\ln(L) = -6516.5$ 525 sites LG 4 rate classes**C****D**


A *PiAMT1* expression in *H. vulgare* 14 dpi




B *PiAMT1* expression in *A. thaliana* 14 dpi




C *PiAMT2* expression in *H. vulgare* 14 dpi




D Colonization of *H. vulgare* 14 dpi (RNA)




E Colonization of *A. thaliana* 14 dpi (RNA)






A




Southern blot













B

P. indica strains on YNB + 10 mM ammonium




A**B****C****D**





PC1 vs PC2



PC3 vs PC2

